

Collection: COST Action E52 Meeting 2008 - Florence (Italy)
 "Evaluation of beech genetic resources for sustainable forestry"
 Guest Editors: Georg von Wühlisch, Raffaello Giannini

Effects of abiotic stress on gene transcription in European beech: ozone affects ethylene biosynthesis in saplings of *Fagus sylvatica* L.

Betz GA⁽¹⁻²⁾, Gerstner E⁽¹⁾, Olbrich M⁽¹⁾, Winkler JB⁽³⁾,
 Langebartels C⁽¹⁾, Heller W⁽¹⁾, Sandermann H⁽¹⁻⁴⁾, Ernst D*⁽¹⁾

The influence of ozone (150-190 nl L⁻¹; 8h/d) on transcription levels of genes involved in the biosynthesis of the stress hormone ethylene, and its precursor 1-aminocyclopropane-1-carboxylate (ACC), was analysed in leaves of European beech saplings. Ozone-induced leaf lesions appeared 7 weeks after onset of ozone exposure. Cell lesion formation was preceded by persistent increases in ethylene emission, in the level of its malonylated precursor ACC, and in the transcript levels of specific ACC synthase 1 (ACS1), ACS2, ACC oxidase 1 (ACO1), and ACO2. Our results demonstrate that mechanisms similar to those operating in herbaceous plants may determine beech saplings responses to ozone exposure.

Keywords: Abiotic stress, Ethylene biosynthesis, *Fagus sylvatica*, Gene expression, Ozone

Introduction

European beech is the most abundant broadleaf tree in Germany (Schütt et al. 1992). It is also of major importance for the European forest industry and was therefore chosen as an experimental species for our

studies, to obtain deeper molecular insights into the potentially detrimental effects of ozone on European broadleaf forest ecosystems. Plants have long been known to exhibit responses to ozone exposure resembling the hypersensitive response following pathogen attack. Ozone has been qualified as an abiotic elicitor of plant defence reactions (Sandermann et al. 1998). These responses include phytoalexin production and synthesis of pathogenesis-related (PR) proteins, accompanied by the formation of various aromatic metabolites like lignins and flavonoids (Langebartels et al. 2002).

Another well-known plant response to ozone exposure is the stimulation of ethylene biosynthesis (Tuomainen et al. 1997, Rao et al. 2002, Langebartels & Kangasjärvi 2004). Activation of ethylene biosynthesis is one of the fastest ozone-dependent biochemical responses so far observed in herbaceous plants, occurring within 5 hours, or even more rapidly (Schlagghauser et al. 1997, Nakajima et al. 2001, Moeder et al. 2002). This response was similarly rapid in birch (Vahala et al. 2003) and in an ozone-sensitive poplar clone (Diara et al. 2005). ACC synthase (ACS) and ACC oxidase (ACO), the two enzymes of ethylene biosynthesis, are encoded by gene families in herbaceous plants; several ACS and ACO isoforms have been reported to be induced by ozone (Nakajima et al. 2001,

Moeder et al. 2002). In ozone-sensitive as well as ozone-tolerant birch clones, ozone-induced accumulation of ACS and ACO transcripts was found (Vahala et al. 2003). Similarly, ozone fumigation of beech saplings (180-200 nl L⁻¹) showed enhanced transcript levels of ACS2 and ACO1 (Nunn et al. 2005a). We therefore extended our study by reporting the cloning of ACS2 and ACO2, and analysing over a period of 3 month an ozone-induced expression of the ethylene biosynthesis genes (ACS1, ACS2, ACO1, ACO2), the accumulation of the ethylene precursor free ACC and conjugated ACC, and the emission of the end product of this pathway, ethylene, using beech saplings.

Materials and methods

Plant material and growth conditions

Three-year-old European beech saplings (*Fagus sylvatica* L., provenance 81024; Schlegel Baumschulen, Riedlingen, Germany) were planted in 14-L pots filled with natural forest soil (site Höglwald, Bavaria, Germany - Betz et al. 2009b). Saplings were maintained during winter under a wooden pergola. Before bud burst, the plants were treated with Promanal[®] (Neudorff, Emmerthal, Germany), to prevent insect infestation. In spring the 4-year-old saplings were transferred to climate-controlled cabinets (39 m²) of a greenhouse covered with UV-permeable glass sheets (<http://www.helmholtz-muenchen.de/eus/index.php> - Olbrich et al. 2005, Betz et al. 2009a). Each cabinet contained 8 tables (2 m² per table) and 12 saplings were placed on a single table. The saplings were cultivated until full leaf development and were kept at a relative humidity of 70-80% and temperatures of 22-25 °C during day hours (6:00 to 20:00) and 17-20 °C during the night (20:00 to 6:00). At the beginning of June, saplings were exposed to ozone for 8 h/d (150 nl L⁻¹) or pollutant-free air, and after 13 d the ozone concentration was increased to 190 nl L⁻¹ for a further 70 d (Betz et al. 2009b). For pollutant treatments ozone was added to conditioned air by mass flow controllers and injected into the cabinet by a fan (1180 m³ h⁻¹). Additional 6 fans within the cabinet provided an even distribution of the pollutant. Ozone was generated by electrical discharge in dry oxygen (Fischer Ozon Generator 500, Neckenheim, Germany) and continuously monitored in the middle of the cabinet using a computer-controlled system and an UV-type ozone analyser CSI 3100 (Columbia Scientific Industries, USA). Ozone concentrations, temperature, humidity and light intensities were continuously measured and adjusted automatically to the climate setpoints by external shading, water nebulisation and aeration. Temperature and

✉ (1) Institute of Biochemical Plant Pathology "Helmholtz Zentrum München, German Research Center for Environmental Health, D-85764 Neuherberg Germany; (2) Olympus Life Science Research Europa GmbH, Sauerbruchstraße 50, D-81377 München (Germany); (3) Department of Environmental Engineering, Helmholtz Zentrum München - German Research Center for Environmental Health, D-85764 Neuherberg (Germany); (4) Ecotox Freiburg, Schubertstr. 1, D-79104 Freiburg (Germany)

@ Dieter Ernst (ernst@helmholtz-muenchen.de)

Received: Feb 04, 2009 - Accepted: Mar 12, 2009

Citation: Betz GA, Gerstner E, Olbrich M, Winkler JB, Langebartels C, Heller W, Sandermann H, Ernst D, 2009. Effects of abiotic stress on gene transcription in European beech: ozone affects ethylene biosynthesis in saplings of *Fagus sylvatica* L. iForest 2: 114-118 [online: 2009-06-10] URL: <http://www.sisef.it/iforest/show.php?id=495>

Tab. 1 - Sequences of primers (5'-3') used for quantitative real-time RT-PCR.

Gene	Acc. nr.	Forward primer	Reverse primer
<i>ACS1</i>	AJ420188	GGTCTTCACTGAAGGTCTTG	CTGCTCTTTGAGTAGGTGAC
<i>ACS2</i>	AY705445	TCCAAACCGCATAGTCATGG	TGGTGAGGGCACAAGAAAAG
<i>ACO1</i>	AJ420190	CTGGTGGGATCATCTTACTC	CAATAGAATGGCGCATAGGG
<i>ACO2</i>	AM062963	GAAGGAAGTGTGACATAGCT	GAAAACCCCTTCTTCTGTAG
<i>26S rRNA</i>	-	CGGCTCTTCTATCATTGTG	AACCTGTCTCACGACGGTCT

humidity were as described above, and no artificial light exposure was used. At sunny days maximum light intensities were 30 kLux. To minimize changes of ozone concentrations aeration was only opened at a temperature of 28 °C. In addition to the water nebulisation two tables were flooded with water up to 4 cm. Watering of saplings was carried out with deionized water using an automatically droplet watering system (250 ml/d/pot). No additional nutrient supply was carried out. Pest control against beech scale was carried out weekly by hand to spare beneficial insects like larvae of lacewings and hover flies. Thrips were fought off with predatory mites and predatory bugs (Sautter & Stepper, Ammerbuch, Germany). Two to three leaves were collected for RNA analysis from four ozone-treated and four control saplings at 15 different time points beginning at the onset of the ozone treatment (days: 0, 2, 6, 11, 16, 21, 27, 34, 41, 48, 55, 62, 69, 76, 83). The leaves were immediately frozen in liquid N₂ and stored at -80 °C until further analyses.

Isolation of RNA and cDNA transcription

Total RNA, was isolated according to the protocol described by Kiefer et al. 2000, treated with RQ1 DNase (Promega, Mannheim, Germany) and quantified photometrically (NanoDrop system; Kisker, Steinfurt, Germany). For cDNA synthesis, 5 µg of total RNA (14.5 µL), 1 µL of dNTP (10

mM) and 1 µL of oligo-dT₍₁₅₎ primer (5 µg µL⁻¹) was used for first-strand cDNA synthesis. Reverse transcription was carried out for 4 h at 42 °C using Superscript II reverse transcriptase, according to the manufacturer's instructions (Invitrogen, Karlsruhe, Germany).

Cloning of homologous beech cDNA for ACS2 and ACO2

To obtain the putative cDNA sequence *ACS2* from beech, degenerated primers were designed based on conserved coding regions identified from a multiple sequence alignment from different plants using MULTALIN (Corpet 1988). Eight identical PCR reactions (25 µL) were prepared containing 2.5 µL reaction buffer (10x), 1 µL MgCl₂ (50 mM), 0.5 µL dNTP (10 mM), 0.5 µL (10 µM) of degenerate forward (5'-TCTTTCAR-GAYTATCATGGMTTGCC-3') and reverse (5'-GGGVTTCCCTGGMYTYAGRGTYG-GC-3') primer, 0.2 µL cDNA, and 0.1 µL Taq Polymerase (Invitrogen). In the subsequent PCR reaction (1 cycle at 94 °C for 3 min; 35 cycles at 94 °C for 30 s; 45-60 °C for 50 s; 72 °C for 1 min), a temperature gradient was applied to optimise primer annealing. PCR products were cloned into pGEM-T Easy (Promega) and sequenced using an ABI3100 sequencer (Applied Biosystems, Darmstadt, Germany). The resulting cDNA sequence was published under the NCBI accession number AY705445. The cloned *ACO2* (Acc. Nr. AM062963) was de-

rived from a recently published ozone-induced cDNA library of beech leaves (Olbrich et al. 2005).


Quantitative real-time RT-PCR (qRT-PCR)

qRT-PCR was performed in a volume of 12.5 µL of SYBR Green ROX kit (ABgene, Hamburg, Germany), 0.5 µL gene-specific forward primer (10 µM), 0.5 µL gene-specific reverse primer (10 µM - Tab. 1), 0.5 µL cDNA and 11 µL H₂O, using an ABIPrism 7700 Sequence Detector (Applied Biosystems, Darmstadt, Germany). The PCR conditions were as follows: 1 cycle at 50 °C for 2 min and at 95 °C for 10 min; 40 cycles at 95 °C for 15 sec and at 60 °C for 1 min (Betz et al. 2009a). As an internal reference, *26S rRNA* was used, and the relative expression ratio (R) was calculated according to the "Delta-delta" method (eqn. 1):

$$R = \frac{2^{\Delta CT_{\text{target}}(\text{control} - \text{sample})}}{2^{\Delta CT_{\text{reference}}(\text{control} - \text{sample})}}$$


Determination of ethylene emission and concentrations of ACC

For ethylene determination, single fresh leaves from 12 saplings were analysed directly upon harvest (days: 1, 3, 6, 15, 20, 24, 29, 35, 42, 49, 65, 63, 70, 77, 84) as described previously (Nunn et al. 2005a) and determination of free and conjugated ACC from leaf material of 9 pooled saplings (days: 0, 2, 6, 11, 16, 21, 27, 34, 41, 48, 55,

**Fig. 1** - Ozone-induced leaf lesions in leaves of European beech saplings. Classification is as in Tab. 2.

Tab. 2 - Classification of ozone symptoms in leaves of European beech saplings.

Level	Symptoms
L1	No visible lesions
L2	Single punctual lesions
L3	Many small lesions all over the lamina
L4	About 50% lesions of the lamina
L5	About 70% lesions of the lamina (leaves started to roll up)
L6	More than 90% lesions of the lamina (leaves were shed)

**Fig. 2** - Percentage of leaves with ozone-induced lesions corresponding to the classification in Tab. 2 (L1: light green; L2: dark green; L3: yellow; L4: orange; L5: red; L6: dark red).

62, 59, 76, 83) was performed according to Tuomainen et al. (1997).

Results


Treatment of beech saplings with ozone (150-190 nl L^{-1}) led to visible leaf symptoms that were classified into six symptom levels (Fig. 1, Tab. 2). Lesions occurred on the upper leaf surface 42-49 d after the onset of treatment (Fig. 2). At the end of the experiment, ozone-treated saplings started to shed the leaves. Nine out of the 12 treated saplings developed severe symptoms, whereas control saplings did not show any damage. Ethylene emission is a frequent ozone response in plants, and it correlates with leaf injury and cell death (Moeder et al. 2002, Nunn et al. 2005a). To analyse details of ozone-induced ethylene biosynthesis in our experiment, we started the analysis using clones putatively assigned to *ACS2* and *ACO2*. Pairwise comparison of *ACS2* with nine *ACS* isoforms of tomato showed an identity of only 70%. However, CLUSTALW analysis with a homologous region of 600 bp indicated 82% identity with *ACS2* from birch, but only 70% with *ACSI*

from the same species (Vahala et al. 2003). The cloned *ACO2* from beech showed about 97% identity to the 3'-open reading frame of *ACO1* from beech (Calvo et al. 2004). *ACO2* had a 73 bp insertion in the C-terminal coding region compared to *ACO1*, which resulted in a protein sequence extension of 12 amino acids. This allowed us to design a specific primer for *ACO2* for qRT-PCR studies. For the known *ACO1* of beech we did not find such isoform specificity. Therefore, we could not exclude the possibility that the expression data for *ACO1* may mirror the more strongly expressed *ACO2*.

ACSI expression increased with the onset of ozone fumigation, then decreased, and increased again with increasing ozone concentration from 150 nl L^{-1} to 190 nl L^{-1} (Fig. 3). *ACS2* showed the highest expression level of all the genes analysed in this study, about 20 orders of magnitude higher than *ACSI* (Fig. 3 - please note the different graduation for the relative gene expression charts). *ACO1* showed the lowest induction rate of all the genes related to ethylene biosynthesis, and a second increase was observed for *ACO1* and *ACO2* after the ozone

concentration was increased (Fig. 3). The time-dependence of the expression levels of *ACO2* and *ACO1* were comparable, although the *ACO2* levels were much higher (Fig. 3). In contrast to what was observed with *ACS* isoforms, levels of *ACO* isoform transcripts did not increase when the macroscopic lesions appeared (Fig. 3).

Free and conjugated ACC, the precursor molecules of ethylene, were regularly analysed at 15 time points throughout the 83 d ozone fumigation period (Fig. 4a). Leaf material of 9 ozone-treated and 9 control saplings were pooled and analysed. As the small saplings should not lose more than 1/10 of their leaves, which might mimic pest attack and no ozone effects, single sapling analyses could not be carried out. Levels of both metabolites increased in the course of fumigation, although to different extents. The highest concentration of free ACC was about 20 nmol g^{-1} fw, whereas conjugated ACC reached 175 nmol g^{-1} fw on day 62, and then decreased. Control values of free and conjugated ACC were below 0.56 and 6.9 nmol g^{-1} fw, respectively, throughout the experiment (Fig. 4a). Interestingly, conjugated

**Fig. 3** - Quantitative real-time RT-PCR analysis of *ACSI*, *ACS2*, *ACO1* and *ACO2* gene transcripts in total RNA isolated from ozone-treated leaves of European beech. 4-year-old European beech saplings were fumigated with ozone (150-190 nl L^{-1} , 8 h/d, 83 d); n: 4 saplings, \pm SEM; (*): increasing the ozone concentration from 150 nl L^{-1} to 190 nl L^{-1} ; (**): first ozone-induced leaf lesions.




Fig. 4 - (a) Accumulation of free and conjugated ACC in leaves of control and ozone-treated 4-year-old European beech saplings (150-190 nL L⁻¹, 8 h/d, 83 d); leaf material of 9 ozone-treated and 9 control saplings was pooled; ↓ = first ozone-induced leaf lesions. (b) Boxplot diagram of the mean ethylene emission of ozone-treated 4-year-old European beech saplings (150-190 nL L⁻¹, 8 h/d, 83 d). The upper, middle and lower margins of the plot correspond to 75%, 50% and 25% percentiles of the monitoring; n: 12 saplings; (*): first ozone-induced leaf lesions.

ACC rapidly increased with the first appearance of macroscopic lesions (Fig. 4a).

Ethylene emission was analysed from freshly harvested leaves of 12 saplings. The absolute amount of ethylene produced as well as the time course of expression varied greatly between different saplings. Nevertheless, a box blot analysis revealed an apparent correlation between ozone fumigation and ethylene emission (Fig. 4b).

Discussion

Young beech saplings are considered as relatively sensitive to the pollutant ozone, although genetic variabilities are known (Langebartels et al. 1997, Grams et al. 1999, Nunn et al. 2005b, Betz et al. 2009a). The ozone concentration used in this study resulted in an AOT40 value of about 96 $\mu\text{L L}^{-1} \text{h}^{-1}$, which is comparable with previous studies (Nunn et al. 2005a, Nunn et al. 2005b). However it has to be kept in mind that a newly developed ozone flux model might be a better risk assessment approach for European forest trees (Matussek et al. 2007, Paoletti et al. 2008). There is some information indicating that ozone-induced leaf damage in beech saplings is linked to the formation of ethylene (Langebartels et al. 1997, Nunn et al. 2005a). In the present study, we showed that the ozone-induced production of ethylene differed widely from tree to tree, but nevertheless could be correlated with the extent of leaf damage and the appearance of ozone symptoms 42 d after onset of ozone exposure (190 nL L⁻¹ - Fig. 4b). Similar strong variation in ethylene formation and symptom development among individuals of a plant species has been reported in herbaceous plants (Mehlhorn & Wellburn 1987, Tamaoki et al. 2003). However, in contrast to the rapid induction of ethylene production

in herbaceous plants (Nakajima et al. 2001, Moeder et al. 2002), production was remarkably delayed in beech and displayed a markedly longer persistence (Fig. 4b). This is consistent with findings recently published by other investigators (Nunn et al. 2005a).

Previous studies with herbaceous plants have shown that the levels of free and conjugated ACC increase upon ozone exposure (Moeder et al. 2002). Moreover, in 60-year-old beech trees exposed to two times the ambient ozone levels, conjugated ACC was induced only in shade leaves, but not sun leaves (Nunn et al. 2005a). In the present greenhouse study with 4-year-old beech saplings, the maximum levels of free and conjugated ACC, as well as maximal ethylene production, occurred about 9 weeks after the onset of ozone exposure (190 nL L⁻¹; 8 h/d - Fig. 4). This delayed increase in ACC may be explained by the fact that beech is better equipped to detoxify ozone than the herbaceous tomato plant (Tuomainen et al. 1997, Moeder et al. 2002). As the leaves of young beech saplings more closely resemble the sun leaves rather than the shade leaves of a mature tree, this difference in response may reflect an ontogenetic effect. There is also increasing evidence that juvenile trees may differ in physiology and, therefore, in ozone responses compared to mature trees (Matussek & Sandermann 2003, Nunn et al. 2005b).

Previous studies with herbaceous plants have also shown that ozone-induced ethylene emission is regulated by differential transcription of the *ACS* and *ACO* genes (Tuomainen et al. 1997, Moeder et al. 2002). We have analysed the transcript levels of two previously known genes (*ACS1*, *ACO1*), as well as two genes (*ACS2*, *ACO2*) cloned here for the first time. Up-regulation of

ACS2 transcription was delayed compared to that of *ACS1*. However, the relative gene expression was more than 1000-fold and was the highest of all transcripts analysed in this study (Fig. 3). A delayed up-regulation of *ACS2* compared to *ACS1* has also been shown in tomato after a single ozone pulse (Nakajima et al. 2001). Tomato *ACS1*, however, was much more strongly induced compared to *ACS2* (Nakajima et al. 2001). Interestingly, *ACS2* was also up-regulated in sun leaves of adult European beech in an open-air fumigation site in the presence of ozone at two times the ambient level (Jehnes et al. 2007). In a previous study with beech saplings, *ACS2* was up-regulated by a factor of only 150 when exposed to ozone at 200 nL L⁻¹ from September to October (Nunn et al. 2005a). The reason for this difference in induction factor is not obvious. As the present study was carried out from June to September, it may indicate the influence of light conditions, including UV radiation, which have not been considered so far. In 1-year-old clonal birch plants, *ACS1* was induced in all clones irrespective of ozone tolerance or sensitivity, whereas *ACS2* was induced exclusively in sensitive clones (Vahala et al. 2003). Thus, 4-year-old beech saplings can be classified as ozone-sensitive. This idea is further supported by the absence of free SA in sensitive birch clones (Vahala et al. 2003), as well as in beech saplings (Betz et al. 2009b). *ACO1* and *ACO2* were strongly induced at the onset of ozone treatment (150-190 nL L⁻¹), but decreased upon continuous exposure (Fig. 3). Ozone induction of *ACO1* and *ACO2* transcripts has also been reported recently in leaves of 3-year-old beech saplings (Olbrich et al. 2005, Nunn et al. 2005a) and in moderately ozone-tolerant aspen clones in a free-air ozone fu-

migation study (Gupta et al. 2005). Interestingly, *ACO1* and *ACO2* were much more strongly induced in ozone-sensitive than in ozone-resistant birch clones, again favouring the notion that beech saplings are ozone-sensitive. As for leaf injury and ethylene production, the relative gene expression levels among individual saplings showed pronounced variation.

Conclusion

The observed differential induction of *ACS* and *ACO* isoforms, the accumulation of ACC, and the production of ethylene strongly support the idea that ethylene triggers development of the leaf lesions related to ozone-induced cell death. These findings further support the recently postulated idea that ethylene plays an important role in cell death in leaves of both saplings (Vahala et al. 2003) and adult trees (Nunn et al. 2005a).

Acknowledgements

Excellent technical assistance was provided by Lucia Gößl and Rosina Ludwig. We wish to thank the technical staff of EUS for performing the ozone fumigation. This work was supported by the Deutsche Forschungsgemeinschaft (SFB 607) and in part by the European Community (EVOLTREE, 6th Framework Programme and COST E52 Action).

References

- Betz G, Knappe C, Lapierre C, Olbrich M, Welzl G, Langebartels C, Heller W, Sandermann H, Ernst D (2009a). Ozone affects shikimate pathway transcripts and monomeric lignin composition in European beech (*Fagus sylvatica* L.). *European Journal of Forest Research* 128: 109-116. - doi: [10.1007/s10342-008-0216-8](https://doi.org/10.1007/s10342-008-0216-8)
- Betz AG, Gerstner E, Stich S, Winkler B, Welzl G, Kremmer E, Langebartels C, Heller W, Sandermann H, Ernst D (2009b). Ozone affects shikimate pathway genes and secondary metabolites of European beech (*Fagus sylvatica* L.) grown under greenhouse conditions. *Trees - Structure and Function* 23: 539-553.
- Calvo AP, Nicolás C, Lorenzo O, Nicolás G, Rodríguez D (2004). Evidence for positive regulation by gibberellins and ethylene of ACC oxidase expression and activity during transition from dormancy to germination in *Fagus sylvatica* L. seeds. *Journal of Plant Growth Regulation* 23: 44-53. - doi: [10.1007/s00344-004-0074-7](https://doi.org/10.1007/s00344-004-0074-7)
- Corpet F (1988). Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Research* 16: 10881-10890. - doi: [10.1093/nar/16.22.10881](https://doi.org/10.1093/nar/16.22.10881)
- Diara C, Castagna A, Baldan B, Sodi AM, Sahr T, Langebartels C, Sebastiani L, Ranieri A (2005). Differences in the kinetics and scale of signalling molecule production modulate the ozone sensitivity of hybrid poplar clones: the roles of H₂O₂, ethylene and salicylic acid. *New Phytologist* 168: 351-364. - doi: [10.1111/j.1469-8137.2005.01514.x](https://doi.org/10.1111/j.1469-8137.2005.01514.x)
- Grams TEE, Aneeg S, Häberle K-H, Langebartels C, Matyssek R (1999). Interactions of chronic exposure to elevated CO₂ and O₃ levels in the photosynthetic light and dark reactions of European beech (*Fagus sylvatica*). *New Phytologist* 144: 95-107. - doi: [10.1046/j.1469-8137.1999.00486.x](https://doi.org/10.1046/j.1469-8137.1999.00486.x)
- Gupta P, Duplessis S, White H, Karnosky DF, Martin F, Podila GK (2005). Gene expression patterns of trembling aspen trees following long-term exposure to interacting elevated CO₂ and tropospheric O₃. *New Phytologist* 167: 129-142. - doi: [10.1111/j.1469-8137.2005.01422.x](https://doi.org/10.1111/j.1469-8137.2005.01422.x)
- Jehnes S, Betz G, Bahnweg G, Haberer K, Sandermann H, Rennenberg H (2007). Tree internal signalling and defence reactions under ozone exposure in sun and shade leaves of European beech (*Fagus sylvatica* L.) trees. *Plant Biology* 9: 253-264. - doi: [10.1055/s-2006-924650](https://doi.org/10.1055/s-2006-924650)
- Kiefer E, Heller W, Ernst D (2000). A simple and efficient protocol for isolation of functional RNA from plant tissues rich in secondary metabolites. *Plant Molecular Biology Reporter* 18: 33-39. - doi: [10.1007/BF02825291](https://doi.org/10.1007/BF02825291)
- Langebartels C, Kangasjärvi J (2004). Ethylene and jasmonate as regulators of cell death in disease resistance. In: "Molecular ecotoxicology of plants" (Sandermann H ed). Springer, Berlin, Germany, pp. 75-109.
- Langebartels C, Ernst D, Heller W, Lütz C, Payer H-D, Sandermann H (1997). Ozone responses of trees: results from controlled chamber exposures at the GSF phytotron. In: "Forest decline and ozone: a comparison of controlled chamber and field experiments" (Sandermann H, Wellburn AR, Heath RL eds). Springer, Berlin, Germany, pp. 163-200.
- Langebartels C, Schraudner M, Heller W, Ernst D, Sandermann H (2002). Oxidative stress and defence reactions in plants exposed to air pollutants and UV-B radiation In: "Oxidative stress in plants" (Inzé D, Van Montagu M eds). Taylor & Francis, London, UK, pp. 105-135.
- Matyssek R, Sandermann H (2003). Impact of ozone on trees: an ecophysiological perspective. *Progress in Botany* 64: 349-404.
- Matyssek R, Bytnerowicz A, Karlsson PE, Paoletti E, Sanz M, Schaub M, Wieser G (2007). Promoting the O₃ flux concept for European forest trees. *Environmental Pollution* 146: 587-607. - doi: [10.1016/j.envpol.2006.11.011](https://doi.org/10.1016/j.envpol.2006.11.011)
- Mehlhorn H, Wellburn AR (1987). Stress ethylene formation determines plant sensitivity to ozone. *Nature* 327: 417-418. - doi: [10.1038/327417a0](https://doi.org/10.1038/327417a0)
- Moeder W, Barry CS, Tauriainen AA, Betz C, Tuomainen J, Utriainen M, Grierson D, Sandermann H, Langebartels C, Kangasjärvi J (2002). Ethylene synthesis regulated by biphasic induction of 1-aminocyclopropane-1-carboxylic acid synthase and 1-aminocyclopropane-1-carboxylic acid oxidase genes is required for hydrogen peroxide accumulation and cell death in ozone-exposed tomato. *Plant Physiology* 130: 1918-1926. - doi: [10.1104/pp.009712](https://doi.org/10.1104/pp.009712)
- Nakajima N, Matsuyama T, Tamaoki M, Saji H, Aono M, Kubo A, Kondo N (2001). Effects of ozone exposure on the gene expression of ethylene biosynthetic enzymes in tomato leaves. *Plant Physiology and Biochemistry* 39: 993-998. - doi: [10.1016/S0981-9428\(01\)01319-5](https://doi.org/10.1016/S0981-9428(01)01319-5)
- Nunn AJ, Aneeg S, Betz G, Simons S, Kalisch G, Seidlitz HK, Grams TEE, Häberle K-H, Matyssek R, Bahnweg G, Sandermann H, Langebartels C (2005a). Role of ethylene in the regulation of cell death and leaf loss in ozone-exposed European beech. *Plant, Cell and Environment* 28: 886-897. - doi: [10.1111/j.1365-3040.2005.01339.x](https://doi.org/10.1111/j.1365-3040.2005.01339.x)
- Nunn AJ, Kozovits AR, Reiter IM, Heerd C, Leuchner M, Lütz C, Liu X, Löw M, Winkler JB, Grams TEE, Häberle K-H, Werner H, Fabian P, Rennenberg H, Matyssek R (2005b). Comparison of ozone uptake and sensitivity between a phytotron study with young beech and a field experiment with adult beech (*Fagus sylvatica*). *Environmental Pollution* 137: 494-506. - doi: [10.1016/j.envpol.2005.01.036](https://doi.org/10.1016/j.envpol.2005.01.036)
- Olbrich M, Betz G, Gerstner E, Langebartels C, Sandermann H, Ernst D (2005). Transcriptome analysis of ozone-responsive genes in leaves of European beech (*Fagus sylvatica* L.). *Plant Biology* 7: 670-676. - doi: [10.1055/s-2005-873001](https://doi.org/10.1055/s-2005-873001)
- Paoletti E, Ranieri A, Lauteri M (2008). Moving toward effective ozone flux assessment. *Environmental Pollution* 156: 16-19. - doi: [10.1016/j.envpol.2008.02.025](https://doi.org/10.1016/j.envpol.2008.02.025)
- Rao MV, Lee H, Davis KR (2002). Ozone-induced ethylene production is dependent on salicylic acid, and both salicylic acid and ethylene act in concert to regulate ozone-induced cell death. *The Plant Journal* 32: 447-456. - doi: [10.1046/j.1365-3113X.2002.01434.x](https://doi.org/10.1046/j.1365-3113X.2002.01434.x)
- Sandermann H, Ernst D, Heller W, Langebartels C (1998). Ozone: an abiotic elicitor of plant defence reactions. *Trends in Plant Science* 3: 47-50. - doi: [10.1016/S1360-1385\(97\)01162-X](https://doi.org/10.1016/S1360-1385(97)01162-X)
- Schlaghauser CD, Arteca RN, Pell EJ (1997). Sequential expression of two 1-aminocyclopropane-1-carboxylate synthase genes in response to biotic and abiotic stresses in potato (*Solanum tuberosum* L.) leaves. *Plant Molecular Biology* 35: 683-688. - doi: [10.1023/A:1005857717196](https://doi.org/10.1023/A:1005857717196)
- Schütt P, Schuck J, Stimm B (1992). *Lexikon der Forstbotanik*. Ecomed, Landsberg, Germany.
- Tamaoki M, Matsuyama T, Kanna M, Nakajima N, Kubo A, Aono M, Saji H (2003). Differential ozone sensitivity among *Arabidopsis* accessions and its relevance to ethylene synthesis. *Planta* 216: 552-560.
- Tuomainen J, Betz C, Kangasjärvi J, Ernst D, Yin Z-H, Langebartels C, Sandermann H (1997). Ozone induction of ethylene emission in tomato plants: regulation by differential accumulation of transcripts for the biosynthetic enzymes. *The Plant Journal* 12: 1151-1162. - doi: [10.1046/j.1365-3113X.1997.12051151.x](https://doi.org/10.1046/j.1365-3113X.1997.12051151.x)
- Vahala J, Ruonala R, Keinänen M, Tuominen H, Kangasjärvi J (2003). Ethylene insensitivity modulates ozone-induced cell death in birch. *Plant Physiology* 132: 185-195. - doi: [10.1104/pp.102.018887](https://doi.org/10.1104/pp.102.018887)