

Nursery practices increase seedling performance on nutrient-poor soils in *Swietenia humilis*

Erickson Basave Villalobos ⁽¹⁾, Víctor Manuel Cetina Alcalá ⁽¹⁾, Miguel Ángel López López ⁽¹⁾, Arnulfo Aldrete ⁽¹⁾, David H Del Valle Paniagua ⁽²⁾

Swietenia humilis is a valued tree species for its high-quality wood, among other commercial interests. Attempts to plant the species in southwest Mexico have often failed due to poor quality seedlings combined with low soil fertility and dry environments. Nursery top-pruning and fertilization are practices previously reported to improve seedling quality, and facilitate rapid establishment under poor site conditions. In the present study, the effects of three top-pruning intensities (0%, 25%, and 50%), and two fertilizer regimes (traditional and exponential) on several *S. humilis* seedling morphological and physiological indices were tested in the nursery, and a quality test trial on nutrient-poor soils was conducted. Significant interactions between the above two treatments were not detected. Top pruning at 25% and 50% intensity did not improve *S. humilis* seedling quality. However, exponential nutrient supply exhibited favorable effects on seedling growth. Results of the quality test trial revealed exponential fertilization promoted satisfactory seedling performance when low nutrient availability was a limiting factor.

Keywords: Caobilla, Seedling Quality, Top Pruning, Exponential Fertilization

Introduction

Caobilla (*Swietenia humilis* Zucc., Meliaceae) is a small to medium-sized tree (15-20 m tall) native to the Mesoamerican Pacific tropical dry forest zones, and considered a valuable hardwood in Mexican Pacific coastal lowlands for the local forest industry (Gordon et al. 2003). After decades of commercial over-exploitation, natural populations have been substantially reduced to scattered and small isolated populations. The species was therefore listed under the multilateral treaty CITES (Convention on International Trade in Endangered Species of Wild Fauna and Flora), and classified as vulnerable under the IUCN (International Union for Conservation of Nature) Red List.

Currently, the interest in establishing *S. humilis* for environmental and commercial purposes has increased in the dry tropics of Mexico and Central America. However, re-

forestation activities have been limited by low seedling survival and growth rates, largely because nurseries do not produce high-quality stock of container seedlings for out-planting (CONAFOR 2012).

Current practices in many forest nurseries involve container seedlings planted in local forest soil and little or no fertilizer application. Regardless of the production system, seedling germplasm quality along with cultural practices serve a major role in promoting successful morphological and physiological characteristics, resulting in satisfactory plant performance at the plantation site (Mexal et al. 2002, Rodríguez 2008, Zida et al. 2008, Esen et al. 2012, Villar-Salvador et al. 2012).

Drought and high temperatures in the dry tropics can limit nutrient availability, which often severely affects reforestation (Griscom & Ashton 2011). However, success can be

achieved by establishing seedling plantations during favorable seasons, using rapidly acclimating plant material. Villar-Salvador et al. (2012) reported seedlings with high growth rates due to high photosynthetic performance and increased capacity to remobilize nutrients were well-suited to the above conditions; and such attributes were favored in large seedlings under high nutrient concentrations. As a consequence, cultivation practices that enhance these traits might increase *S. humilis* seedling quality.

Top-pruning seedlings in the nursery is an uncommon practice; nevertheless, Cetina et al. (1999, 2001, 2002) provided evidence in *Pinus greggii* that the approach can increase seedling growth and photosynthetic capacity, resulting in favorable effects on seedling field performance. However, Wilson (2000) demonstrated success was dependent on pruning intensity and removal as a consequence of the apical dominance effect (Wilson 2000).

Seedling fertilization is a common nursery practice, which serves to enhance growth and photosynthetic capacity. Reserves can be accumulated through fertilization when nutrient availability exceeds seedling requirements (Chapin et al. 1990, Timmer 1997, Salifu & Timmer 2003), with the benefit of later remobilization when nutrient availability becomes a limiting factor. Different fertilization regimes, including exponential fertilization have been established to promote these conditions (Timmer 1997, Salifu & Timmer 2003, Duan et al. 2013).

Top-pruning and exponential fertilization effects in the nursery have not been documented in *S. humilis*, although improvement in seedling quality has been reported for many other forest species (Cetina et al. 1999, Isaac et al. 2011, Wang et al. 2013). In this study, we examined top-pruning and fertilization effects on *S. humilis* seedlings by respectively evaluating various morphological and physiological indices in the nursery, and a quality test trial on nutrient-poor soils.

Material and methods

The study was conducted under greenhouse conditions in the Nursery of the Forestry Graduate School of *Colegio de Postgraduados* in Montecillo, Texcoco, Mexico (19° 19' N, 98° 53' W; 2250 m a.s.l.). Greenhouse conditions were maintained at an average 30/15 °C (day/night), a 16 h (day/night) photoperiod, and 70-85% relative humidity.

Seeds were collected from healthy *S. humilis* individuals in a tropical dry forest in the Tierra Caliente, Guerrero, Mexico. In August 2012, 1000 seeds were field collected, scarified, and placed in running water for 24 h to sow directly into Jiffy-7 Forestry® compressed peat pellets. One seed per pellet was sown. Germination and emergence was

□ (1) Postgrado Forestal; (2) Colegio de Postgraduados, campus Montecillo, Km. 36.5 Carretera Federal México-Texcoco, 56230, Montecillo (México)

@ Erickson Basave Villalobos (erbavi_88@hotmail.com)

Received: Nov 12, 2013 - Accepted: Jul 23, 2014 - Final Acceptance: Nov 28, 2014

Citation: Basave Villalobos E, Cetina Alcalá VM, López López MA, Aldrete A, Del Valle Paniagua DH, 2014. Nursery practices increase seedling performance on nutrient-poor soils in *Swietenia humilis*. iForest 8: 552-557 [online 2014-12-09] URL: <http://www.sisef.it/forest/contents/?id=ifor1179-007>

Communicated by: Alberto Santini

observed during the last week of August 2012. Seven weeks later on 19 October 2012, 384 seedlings of approximately the same height (9-10 cm) were selected and organized into groups of 16 seedlings, each group corresponding to an experimental unit. Six treatment combinations with four replicates per treatment resulting in a 3x2 factorial arrangement were randomly assigned to each of the 24 experimental units in a completely randomized design.

The two factors tested were top-pruning (TP), with three levels (no-pruning - NP, TP 25%, and TP 50% intensity), and fertilization, with two levels (traditional fertilization - TF, and exponential fertilization - EXF). The six treatment combinations were labeled as T1 (NP + TF), T2 (TP 25% + TF), T3 (TP 50% + TF), T4 (NP + EXF), T5 (TP 25% + EXF), and T6 (TP 50% + EXF). Depending on the effects, NP and TF levels, and their combinations (NP + TF), were considered controls.

Treatments were initiated on 22 October 2012, when nutrient reserves in the seedling endosperm were largely exhausted. Top-pruning (TP) was conducted once, when fertilization was first applied, by removing the stem portion between the second node and the apex (including foliage). Fertilization was further supplied from 22 October to 28 December 2012. The fertilizer amount supplied was designed for a 10-week period, and based on nitrogen (N) concentration estimates in seedling foliage.

TF treatment followed the PRONARE-CO-NAFOR fertilization program for tropical species using Peters® fertilizers (Aldana & Aguilera 2003), which were adjusted to commercial formulations based on seedling developmental stage. Under the TF fertilization regime, seedlings were treated with Peters Professional® Seedling Starter™ (9 N: 45 P₂O₅:15 K₂O) at concentrations of 25 mg N l⁻¹ for a total of six applications, *i.e.*, three weeks, two applications per week. Seedlings were administered eight applications of Peters Professional® General Purpose™ (20 N: 10 P₂O₅: 20 K₂O) at concentrations of 50 mg N l⁻¹ with the same frequency over the following four weeks. During the final three weeks, a total of six treatments, two per

week of Peters Professional® Conifer Finisher™ (4 N: 25 P₂O₅: 35 K₂O) were applied at a 50 mg N l⁻¹ concentration.

The EXF treatment was designed to supply nutrients exponentially, increasing the fertilizer addition levels through time. Calculations were derived from an exponential function described by Timmer (1997 - eqn. 1):

$$N_T = N_S(e^{rt} - N_{(t-1)})$$

where *t* is the number of fertilizer applications during the nursery stage, *N_T* is the fertilizer amount applied on a specific day *t*, *N_(t-1)* is the cumulative fertilizer treatment amount at a specific time (including the previous application), and *N_S* is the initial N content in newly germinated seedlings (7.12 mg N seedling⁻¹). A random sample of 30 newly germinated seedlings was analyzed to determine *N_S*.

The *r* factor is the relative nutrient addition rate to increase *N_S* to the final (*N_T* + *N_S*) level, obtained by the following equation (eqn. 2):

$$r = \frac{\ln(N_T / N_S + 1)(e^t - 1)}{A}$$

where *N_T* is the amount of fertilizer applied on a specific day *t*. Due to limited information on the species nutritional attributes, *N_T* (27.78 mg N seedling⁻¹) was determined from the N amount applied per seedling using the TF scheme. As in TF, fertilizer was applied twice per week, therefore *A* = 20. The doses obtained (mg N seedling⁻¹) for each application date were multiplied by 32. This value was used as a constant considering that for each liter of nutrient solution, 32 seedlings are fertilized. The specific N concentrations used are shown in Fig. 1. For all EXF fertilizer treatments, Peters Professional® General Purpose™ (20 N: 20 P₂O₅: 20 K₂O) was applied to seedlings.

Fertilizer solution pH and electrical conductivity were adjusted following Landis (1989). Fertilizer residue on seedling leaves was removed with water after each application. Additionally, flushing irrigation was conducted weekly. Treatment fertilizer applications at different concentrations, *i.e.*, TF and EXF were the only nutrient sources, as

distilled water was used throughout the experiment.

Three weeks after the final fertilization (21 January 2013), a random sample of 16 seedlings per treatment (four from each replicate) was selected to examine shoot height, root-collar diameter, net photosynthetic rate, and chlorophyll content. Twelve of the above 16 seedlings per treatment were further analyzed for leaf area, shoot and root dry weight, and nitrogen (N), phosphorus (P), and potassium (K) concentrations.

Shoot height (cm) and root-collar diameter (mm) were respectively measured with a graduated ruler and a Mitutoyo digital caliper. The sturdiness index (*SI*) was calculated from the ratio between height and root-collar diameter (Johnson & Cline 1991). Net photosynthetic rate (NPR, μmol m⁻² s⁻¹) was measured between 09:00 and 12:00 a.m. with a CIRAS -2 (PP Systems, Haverhill, MA, USA), and chlorophyll content (CC, SPAD units) was indirectly determined by a Minolta SPAD 502 Plus portable chlorophyll meter (Spectrum Technologies Inc., Plainfield, IL, USA). These last two measures were derived from the same leaves previously selected based on (apparent) similar physiological age. Leaf area (LA, cm²) was determined with an area meter (3100 Biosciences Li-Cor Biosciences, Lincoln, NE, USA).

Shoot and root dry weight were obtained by oven-drying seedlings at 70 °C for 72 h. Dry weight (g) of aboveground components (shoots + leaves), and roots with shoot-heights and root-collar diameters were combined to estimate the Dickson quality index (*DQI*) based on the following equation (Johnson & Cline 1991 - eqn. 3):


$$DQI = \frac{DW_{total}}{(H/D) + (DW_{shoot}/DW_{root})}$$

Finally, N, P, and K concentrations were determined from tissue samples (three per treatment) integrated over the entire seedling by the micro-Kjeldahl method, using photolorimetry by reduction with Molibdo-Vanadate and flame emission spectrometry, respectively. Total dry biomass and nutrient concentrations were used to calculate N, P, and K content.

Seedling quality, *e.g.*, growth, development, and establishment at nutrient-poor sites was examined by conducting a quality test trial in February 2013. A 30-cm deep planting-bed with red volcanic rock substrate (tezontle) was constructed to establish low or no nutrient availability. Ten randomly selected seedlings per treatment (a total of 60 seedlings) were sown in rows on the planting bed following a complete randomized design.

The quality test trial ended in May 2013. During the four month period, seedlings

Fig. 1 - Concentration of N per liter of nutrient solution provided in each application of the exponential fertilization regime.



were periodically watered with distilled water, which was used to prevent any source of external nutrient supply. Seedling height (cm) and stem diameter (mm) were measured at the beginning and end of the test, and monthly relative growth rates were estimated based on the following equation (Pallardy 2008 - eqn. 4):

$$RGR = \frac{\ln X_2 - \ln X_1}{\Delta t}$$

where X_1 and X_2 are the variables measured at the first and second assessment time, respectively, and Δt is the time interval between the two measurements.

Comparisons of the factor effect and its interactions on morphological and physiological quality indices were performed through analysis of variance (ANOVA, $\alpha = 0.05$). Significant differences among treatment means were analyzed using the *post-hoc* Fisher's LSD test ($\alpha = 0.05$). Logarithmic and square root transformations of raw data were conducted when assumptions of normality and homoscedasticity were violated.

Statistical analyses were carried out using the package INFOSTAT[®] (Di Rienzo et al. 2012). Statistical analyses were not conducted on N, P, and K concentration and content. Instead, average values were reported for each composite sample. Graphical vector analysis (Timmer & Stone 1978, López & Alvarado 2010) was performed to interpret nutrient variation among treatments.

Additionally, monthly relative growth rates were analyzed by treatment and not factors, since the quality test was specifically designed to assess seedling performance as a function of nutritional reserves under low soil fertility conditions. Therefore, treatments better reflected the influence of plant nutrient status on seedling growth.

Results

Morphological and physiological quality indices

ANOVA results indicated the main factors, *i.e.*, TP and fertilization regime had significant effects on most morphological and physiological variables analyzed (Tab. 1). However, the interaction between TP and fertilization was not significant for any parameters. In summary, seedlings under the NP and EXF treatments showed the highest values for all variables measured (Tab. 2).

Under the NP treatment, the most divergent morphological and physiological quality index response occurred for shoot and root dry weight, leaf area, and net photosynthetic rate (Tab. 2). NP seedlings accumulated 17% and 47% more shoot biomass than seedlings pruned at 25% and 50%, respectively. Similar results were obtained for root dry weight, where NP control seedlings accumulated 24% and 50% more root biomass compared

Tab. 1 - ANOVA results for top pruning and fertilization regime effects on morphological and physiological variables related to seedling quality in *Swietenia humilis*.

Variables	Factors	df	F-value	P-value
Root-collar diameter	Top pruning	2, 90	9.01	0.0003
	Fertilization regime	1, 90	1.32	0.2542
	Interaction	2, 90	1.47	0.2362
Sturdiness index	Top pruning	2, 90	3.01	0.0542
	Fertilization regime	1, 90	5.25	0.0243
	Interaction	1, 90	1.89	0.1572
Shoot dry weight	Top pruning	2, 66	5.93	0.0001
	Fertilization regime	1, 66	10.31	0.0176
	Interaction	2, 66	0.01	0.99
Root dry weight	Top pruning	2, 66	10.01	0.0002
	Fertilization regime	1, 66	1.28	0.2614
	Interaction	2, 66	0.02	0.9754
Dickson's quality index	Top pruning	2, 66	6.75	0.0022
	Fertilization regime	1, 66	0.56	0.4572
	Interaction	2, 66	0.06	0.9387
Leaf area	Top pruning	2, 66	7.71	0.001
	Fertilization regime	1, 66	33.78	<0.0001
	Interaction	2, 66	0.7	0.5025
Chlorophyll content	Top pruning	2, 90	0.24	0.791
	Fertilization regime	1, 90	26.34	<0.0001
	Interaction	2, 90	0.81	0.4493
Net photosynthesis rate	Top pruning	2, 90	40.88	<0.0001
	Fertilization regime	2, 90	175.3	<0.0001
	Interaction	2, 90	0.59	0.5543

with TP treatments at 25% and 50%, respectively. Additionally, NP seedlings showed the largest leaf area, 40% and 36% greater than 25% and 50% TP seedlings, respectively. Finally, photosynthetic rate of NP seedlings was 49% higher than 25% TP plants, and 32% higher compared with 50% TP plants.

Results of the fertilization effect showed seedlings under the EXF treatment regime produced 17% more biomass and had 50% more leaf area compared with TF treatment seedlings. Similarly, physiological variables, such as chlorophyll content (CC) and net

photosynthetic rate (NPR) were significantly affected by EXF, evidenced by a respective 20% and 60% higher CC and NPR in seedlings compared with those grown under the TF regime (Tab. 2).

N, P, and K concentration and content

Differences in plant nutrient status were detected in response to treatments. Compared with the control treatment (T1), seedlings receiving treatments T4, T5, and T6 exhibited increased N concentration and content, respectively (126% and 157% for T4, 75% and 67% for T5, and 364% and 260%


Tab. 2 - Average values (\pm SE) of morphological and physiological quality indexes assessed in *Swietenia humilis* seedlings in response to top pruning and fertilization treatments. Only means for factors having a significant effect after ANOVA are reported. Means with the same letter within each treatment are not significantly different ($p > 0.05$) after Fischer's LSD test. (NP): No-pruned seedlings; (TP): top-pruned seedlings; (TF): traditional fertilization; (EXF): exponential fertilization; (SI): sturdiness index; (DQI): Dickson quality index; (NPR): net photosynthesis rate; (CC): chlorophyll content.

Variables	Pruning			Fertilization	
	NP	TP 25%	TP 50%	TF	EXF
Diameter (cm)	4.97 \pm 0.11 ^a	4.88 \pm 0.11 ^a	4.37 \pm 0.11 ^b	-	-
SI (cm mm ⁻¹)	-	-	-	2.20 \pm 0.06 ^a	2.39 \pm 0.06 ^b
Shoot dry weight (g)	1.16 \pm 0.06 ^a	0.99 \pm 0.06 ^b	0.79 \pm 0.06 ^c	0.90 \pm 0.05 ^a	1.06 \pm 0.05 ^b
Root dry weight (g)	0.51 \pm 0.03 ^a	0.41 \pm 0.03 ^b	0.34 \pm 0.03 ^b	-	-
DQI	0.36 \pm 0.02 ^a	0.31 \pm 0.02 ^a	0.25 \pm 0.02 ^b	-	-
Leaf area (cm ²)	73.56 \pm 4.07 ^a	52.21 \pm 4.07 ^b	54.07 \pm 4.07 ^b	46.82 \pm 3.33 ^a	73.28 \pm 3.33 ^b
NPR ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	3.54 \pm 0.09 ^a	2.37 \pm 0.09 ^b	2.69 \pm 0.09 ^c	2.18 \pm 0.08 ^a	3.55 \pm 0.08 ^b
CC (SPAD units)	-	-	-	31.83 \pm 0.89 ^a	38.28 \pm 0.89 ^b

Tab. 3 - Concentration and content of N, P and K in response to pruning and fertilization treatments in *Swietenia humilis* seedlings in the nursery.

Treatment combination	Concentration (mg g ⁻¹)			Content (mg plant ⁻¹)		
	N	P	K	N	P	K
NP + TF (T1)	8.1	3	12.2	12.7	4.7	19.13
TP 25% + TF (T2)	8.1	2.4	11.4	10.56	3.13	14.87
TP 50% + TF (T3)	9.1	2.7	11.6	9.49	2.81	12.1
NP + EXF (T4)	18.3	4.6	11.3	32.6	8.19	20.13
TP 25% + EXF (T5)	14.2	3.2	11.5	21.16	4.77	17.14
TP 50% + EXF (T6)	37.6	4.7	18.9	45.68	5.71	22.96

Fig. 2 - Nomograms for response of nitrogen (a), phosphorus (b), and potassium (c), to the application of different top pruning intensities and different fertilization regimes in *Swietenia humilis* seedlings in the nursery. Treatment combinations: (T1) no pruning + TF; (T2) pruning at 25% + TF; (T3) pruning at 50% + TF; (T4) no pruning + EXF; (T5) pruning at 25% + EXF; (T6) pruning at 50% + EXF.



Tab. 4 - Average relative growth rates in *Swietenia humilis* seedlings planted in low or null availability of nutrients. Means with the same letter within columns are not significantly different ($p > 0.05$) after Fischer's LSD test.

Treatment combination	Relative growth rates	
	Height (cm cm ⁻¹ month ⁻¹)	Diameter (mm mm ⁻¹ month ⁻¹)
NP + TF (T1)	0.10 ± 0.01 ^a	0.05 ± 0.01 ^{ab}
TP 25% + TF (T2)	0.09 ± 0.01 ^a	0.04 ± 0.01 ^a
TP 50% + TF (T3)	0.12 ± 0.01 ^{ab}	0.08 ± 0.01 ^{bc}
NP + EXF (T4)	0.15 ± 0.01 ^b	0.11 ± 0.01 ^d
TP 25% + EXF (T5)	0.11 ± 0.01 ^a	0.05 ± 0.01 ^{abc}
TP 50% + EXF (T6)	0.14 ± 0.01 ^b	0.09 ± 0.01 ^{cd}

for T6). P concentration in T4 and T6 was respectively 53% and 74% higher than in T1. Similarly, P content in T4 and T6 treatments were respectively 57% and 21% higher than in T1. Finally, only T6 revealed higher K concentration (> 55%) and content (> 20%) than T1 (Tab. 3).

Quality test trials in low nutrient substrate (Tezontle)

Seedling growth in the quality test trial revealed significant effects for height ($F_{[5, 54]} = 3.4, P < 0.01$) and diameter ($F_{[5, 54]} = 4.91, P < 0.001$), with T4 seedlings showing the highest relative growth rates (Tab. 4).

Discussion

Morphological and physiological quality indices

Our results showed TP at 25% or 50% did not improve the morphological or physiological quality of *S. humilis* seedlings, unless accompanied by EXF fertilization. Indeed, the NP treatment seedlings exhibited the highest value for all quality parameters examined (Tab. 2). The growth responses of TP seedlings, i.e., root-collar diameter, shoot dry weight, leaf area, and *DQI* were likely affected by the stress imposed in shoot biomass loss.

Previous reports demonstrated that after defoliation, plants accumulated nonstructural carbohydrates for regrowth to compensate tissue loss (Kozłowski 1992, von-Fircks & Sennerby-Fores 1998). Chapin et al. (1990) showed the shoot partially and/or temporarily lost the capacity to produce photosynthates, as accumulation was largely dependent on resource allocation to storage. However, allocation of resources to storage directly competes with growth (Berendse et al. 2007). Moreover, growth patterns in different forest species might differ as a consequence of differential biomass allocation to shoots and/or roots (Ericsson 1995). These trade-offs might partly explain reduced growth in the pruned *S. humilis* seedlings observed in this study.

Cetina et al. (1999, 2001) suggested the morpho-physiological responses in *P. greggi* were associated with pruning, and might be beneficial to enhance seedling quality. The same pruning intensity tested in the present study (25% and 50%) was applied to *P. greggi* seedlings, resulting in increased dry biomass accumulation in shoots and roots, as well as in leaf area and the photosynthetic rate (Cetina et al. 1999, 2001). However, our findings suggested TP was not a suitable approach to enhance *S. humilis* seedling quality. First, this lack of congruency might be attributed to among species differences. Cline & Harrington (2007) observed growth and branching patterns among temperate and tropical species differ. Moreover, Chapin et

al. (1990) suggested the benefits obtained by apical dominance removal can vary as a consequence of different species capacities to store the carbohydrates necessary to recover from pruning. Furthermore, seedling age might partially explain these inconsistencies. Seedlings used by Cetina et al. (1999) were older than the seedlings investigated in our work. However, young seedlings have not yet developed an efficient root system. Therefore, seedlings cannot recover from the injuries caused by pruning through resprouting (Kozłowski 1992), unless the resources necessary to support regrowth are provided by fertilization.

Exponential nutrient loading applied in this study improved the morphological and physiological attributes of *S. humilis* seedlings, results inconsistent with pruning. Despite the lack of reports on this particular species, the favorable effects of exponential nutrient supply on growth, and other beneficial characteristics in the nursery have been examined by many authors (Qu et al. 2003, Salifu & Timmer 2003, Salifu & Jacobs 2006, Isaac et al. 2011, Duan et al. 2013, Wang et al. 2013). Timmer (1997) reported this technique has the advantage of supplying the nutrient amounts required at each stage of seedling exponential growth. Seedling responses expressed in morphological and physiological quality indices confirmed these effects (Tab. 2).

Enhanced seedling growth, *i.e.*, leaf area and shoot biomass under the EXF treatment also suggested greater resource exploitation, *i.e.*, light and CO₂, resulting in increased NPR (Tab. 2). Cuesta et al. (2010) reported higher photosynthetic rates in *P. halepensis* seedlings that exhibited greater leaf area. Villar-Salvador et al. (2012) emphasized a large aerial biomass advantage in capturing the resources required to synthesize new tissues, which lead to a positive seedling growth feedback pathway (Kozłowski 1992). However, plant growth and photosynthetic capacity are dependent on plant nutrient status under greenhouse (Villar-Salvador et al. 2012) or natural conditions (Wright et al. 2011). Field evidence demonstrated that both processes were limited by mineral nutrient availability (Pasquini & Santiago 2011). Furthermore, the rate of carbon gain is a function of the photosynthetic rate, respiration, and tissue loss, which typically exhibits interspecific and environmental variability (Berendse et al. 2007, Kitajima 2007, Cuesta et al. 2010).

N, P, and K concentration and content

In the present study, nutrient reserve amounts (N, P, and K) were higher in plants fertilized with the EXF treatment regime (Tab. 3), demonstrating the treatment's effectiveness in promoting the storage of these resources. Kim et al. (1999) reported plant

nutrient reserves declined at the planting site as a consequence of reduced nutrient availability due to interspecific competition (Timmer 1997), or severe soil degradation (Griscom & Ashton 2011). According to Tab. 3, the beneficial effects of exponential fertilization were clearly demonstrated compared with the traditional fertilization regime. Nevertheless, it is worth noting that TP also influenced seedling response to fertilization. Indeed, increased growth and N and P concentration and content were achieved under NP + EXF treatments (T4). However, T6 showed maximum concentration and content of all nutrients (50% TP + EXF), although a lower biomass amount was produced. López & Alvarado (2010) indicated this type of effect resulted from toxicity; however, we determined growth was reduced by pruning, and not by nutrient excess (Fig. 2).

Quality test trial on nutrient-limited soils

Results from the quality test in Tezontle (Tab. 4) confirmed seedling nutrient accumulation due to exponential fertilization facilitated growth on nutrient-limited soils. Heiskanen et al. (2009) obtained similar results in *Picea abies*, where exponential seedling fertilization clearly allowed rapid seedling establishment on poor sites. However, Everett et al. (2007) found a complete lack of congruency in results of a study on morphology and nutritional status of *Pseudotsuga menziesii* var. *glauca* seedlings.

Varied interpretations of species differential responses to exponential fertilization have been proposed. Burgess (1991) attributed the limited response of some fast-growing species to fertilizer addition rates; at early developmental and growth stages, the fertilizer amounts might be insufficient to meet seedling requirements. However, toxic fertilizer addition rates might also generate detrimental effects (Duan et al. 2013).

Based on our short-term trial results, the doses applied (17 to 412 mg N l⁻¹) using a complete formula of soluble fertilizer (20 N: 20 P₂O₅: 20 K₂O) resulted in beneficial effects on accumulation reserves to support growth, when no external nutrient supply was available. However, we cannot exclude toxic effects due to exponential fertilization, particularly when the 50% TP + EXF treatment was administered. Indeed, exponential fertilization assumes a relationship between fertilizer dose and plant size (Timmer 1997), while in the above treatment, plant size, *i.e.*, pruning reduced biomass.

Acknowledgments

We are grateful to Celi G. Calixto Valencia, Noé Bahena Tamayo† and Itzel Dinoran Galindez for their technical support.

References

- Aldana BR, Aguilera RM (2003). Procedimientos y cálculos básicos, útiles en la operación de viveros que producen en contenedor [Procedures and basic calculations useful for forest tree seedlings production systems]. Programa Nacional de Reforestación. Comisión Nacional Forestal, Jalisco, México, pp. 46.
- Berendse F, de Kroon H, Braakhekke WG (2007). Acquisition, use and loss of nutrients. In: "Functional Plant Ecology (2nd edn)" (Pugnaire FI, Valladares F eds). CRC Press, Boca Raton, FL, USA, pp. 259-283.
- Burgess D (1991). Western hemlock and Douglas-fir seedling development with exponential rates of nutrient addition. *Forest Science* 37: 54-57. [online] URL: <http://www.ingentaconnect.com/content/saf/fs/1991/0000037/0000001/art00007>
- Cetina AVM, González HVA, Vargas HJJ (1999). El manejo en vivero de *Pinus greggii* Engelm. y la calidad de planta [Nursery management of *Pinus greggii* Engelm. and plant quality]. *Agrociencia* 33: 423-430.
- Cetina AVM, Ortega DML, González HVA, Vargas HJJ, Colinas LMT, Villegas MA (2001). Fotosíntesis y contenido de carbohidratos de *Pinus greggii* Engelm. en respuesta a la poda y al régimen de riego en vivero [Photosynthesis and carbohydrate content of *Pinus greggii* Engelm. as affected by pruning and irrigation intervals in the nursery]. *Agrociencia* 35: 599-607.
- Cetina AVM, González HVA, Ortega DML, Vargas HJJ, Villegas MA (2002). Supervivencia y crecimiento en campo de *Pinus greggii* Engelm. previamente sometido a podas o sequía en vivero [Plant survival and growth in field conditions of *Pinus greggii* Engelm. previously submitted to pruning or drought stress in nursery.]. *Agrociencia* 36: 233-241.
- Chapin FS, Schulze ED, Mooney HA (1990). The ecology and economics of storage in plants. *Annual Review of Ecology and Systematics* 21: 423-447. - doi: [10.1146/annurev.es.21.110190.002231](https://doi.org/10.1146/annurev.es.21.110190.002231)
- CONAFOR (2012). Evaluación complementaria del PROCOREF ejercicio fiscal 2011 [Complementary evaluation of PROCOREF fiscal year 2011]. Comisión Nacional Forestal, Zapopan, Jalisco, México, pp. 325.
- Cline MG, Harrington CA (2007). Apical dominance and apical control in multiple flushing of temperate woody species. *Canadian Journal of Forest Research* 37: 74-83. - doi: [10.1139/x06-218](https://doi.org/10.1139/x06-218)
- Cuesta B, Villar-Salvador P, Puértolas J, Jacobs DF, Rey B JM (2010). Why do large, nitrogen rich seedlings better resist stressful transplanting conditions? A physiological analysis in two functionally contrasting Mediterranean forest species. *Forest Ecology and Management* 260: 71-78. - doi: [10.1016/j.foreco.2010.04.002](https://doi.org/10.1016/j.foreco.2010.04.002)
- Di Rienzo JA, Casanoves F, Balzarini MG, González L, Tablada M, Robledo CW (2012). InfoStat versión 2012. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. [online] URL: <http://www.infostat.com.ar>

- Duan J, Xu C, Jacobs DF, Ma L, Wei H, Jiang L, Ren J (2013). Exponential nutrient loading shortens the cultural period of *Larix olgensis* seedlings. *Scandinavian Journal of Forest Research* 28(5): 409-418. - doi: [10.1080/02827581.2013.778328](https://doi.org/10.1080/02827581.2013.778328)
- Ericsson T (1995). Growth and shoot: root ratio of seedlings in relation to nutrient availability. *Plant and Soil* 168-169: 205-214. - doi: [10.1007/978-94-011-0455-5_23](https://doi.org/10.1007/978-94-011-0455-5_23)
- Esen D, Yildiz O, Esen U, Edis S, Çetintas C (2012). Effects of cultural treatments, seedling type and morphological characteristics on survival and growth of wild cherry seedlings in Turkey. *iForest* 5: 283-289. - doi: [10.3832/ifor0639-005](https://doi.org/10.3832/ifor0639-005)
- Everett KT, Hawkins BJ, Kiiskila S (2007). Growth and nutrient dynamics of Douglas-fir seedlings raised with exponential or conventional fertilization and planted with or without fertilizer. *Canadian Journal of Forest Research* 37: 2552-2562. - doi: [10.1139/X07-108](https://doi.org/10.1139/X07-108)
- Gordon JE, Hawthorne WD, Sandoval G, Barrance AJ (2003). Trees and farming in the dry zone of southern Honduras. II. The potential for tree diversity conservation. *Agroforestry Systems* 59: 107-117. - doi: [10.1023/A:1026360205113](https://doi.org/10.1023/A:1026360205113)
- Griscom HP, Ashton MS (2011). Restoration of dry tropical forests in Central America: a review of pattern and process. *Forest Ecology and Management* 261: 1564-1579. - doi: [10.1016/j.foreco.2010.08.027](https://doi.org/10.1016/j.foreco.2010.08.027)
- Heiskanen J, Lahti M, Luoranen J, Rikala R (2009). Nutrient loading has a transitory effect on the nitrogen status and growth of outplanted Norway spruce seedlings. *Silva Fennica* 43 (2): 249-260. - doi: [10.14214/sf.210](https://doi.org/10.14214/sf.210)
- Isaac ME, Adjei EO, Issaka RN, Timmer VR (2011). A strategy for tree-perennial crop productivity: nursery phase nutrient additions in cocoa-shade agroforestry systems. *Agroforestry Systems* 81: 147-155. - doi: [10.1007/s10457-010-9365-0](https://doi.org/10.1007/s10457-010-9365-0)
- Johnson JD, Cline ML (1991). Seedling quality of southern pines. In: "Forest Regeneration Manual" (Duryea ML, Dougherty PM eds). Kluwer Academic Publishers, The Netherlands, pp. 143-159. - doi: [10.1007/978-94-011-3800-0_8](https://doi.org/10.1007/978-94-011-3800-0_8)
- Kim YT, Colombo SJ, Hickie DF, Noland TD (1999). Amino acid, carbohydrate, glutathione, mineral nutrient and water potential changes in non-water-stressed *Picea mariana* seedlings after transplanting. *Scandinavian Journal of Forest Research* 14: 416-424. - doi: [10.1080/02827589950154122](https://doi.org/10.1080/02827589950154122)
- Kitajima K (2007). Seed and seedling ecology. In: "Functional Plant Ecology (2nd edn)" (Pugnaire FI, Valladares F eds). CRC Press, Boca Raton, FL, USA, pp. 549-579.
- Kozłowski TT (1992). Carbohydrate sources and sinks in woody plants. *Botanical Review* 58: 108-222. - doi: [10.1007/BF02858600](https://doi.org/10.1007/BF02858600)
- Landis TD (1989). Mineral nutrients and fertilization. In: "The container tree nursery manual" (Landis TD, Tinus RW, McDonald SE, Barnett JP eds). Agriculture Handbook 674, vol. 4, USDA Forest Service, Washington, DC, USA, pp. 1-67. [online] URL: http://admin.rngr.net/publications/ctnm/volume-4/vol_4_chapter_1.pdf
- López LMA, Alvarado LJ (2010). Interpretación de nomogramas de análisis de vectores para diagnóstico nutricional de especies forestales [Interpretation of vector analysis nomograms for nutrient diagnosis of forest tree species]. *Madera y Bosques* 16(1): 99-108. [in Spanish]
- Mexal JG, Cuevas RRA, Negreros-Castillo P, Paraguirre LC (2002). Nursery production practices affect survival and growth of tropical hardwoods in Quintana Roo, México. *Forest Ecology and Management* 168: 125-133. - doi: [10.1016/S0378-1127\(01\)00735-6](https://doi.org/10.1016/S0378-1127(01)00735-6)
- Pallardy SG (2008). Physiology of woody plants (3rd edn). Academic Press, MA, USA, pp. 454.
- Pasquini SC, Santiago LS (2011). Nutrients limit photosynthesis in seedlings of a lowland tropical forest tree species. *Oecologia* 168 (2): 311-319. - doi: [10.1007/s00442-011-2099-5](https://doi.org/10.1007/s00442-011-2099-5)
- Qu L, Quoreshi AM, Koike T (2003). Root growth characteristics, biomass and nutrient dynamics of seedlings of two larch species raised under different fertilization regimes. *Plant and Soil* 255: 293-302. - doi: [10.1023/A:1026159709246](https://doi.org/10.1023/A:1026159709246)
- Rodríguez TDA (2008). Indicadores de calidad de planta forestal [Indicators of tree seedling quality]. Mundi-Prensa, México, pp. 156.
- Salifu KF, Timmer VR (2003). Optimizing nitrogen loading of *Picea mariana* seedlings during nursery culture. *Canadian Journal of Forest Research* 33: 1287-1294. - doi: [10.1139/x03-057](https://doi.org/10.1139/x03-057)
- Salifu KF, Jacobs DF (2006). Characterizing fertility targets and multi-element interactions in nursery culture of *Quercus rubra* seedlings. *Annals of Forest Science* 63: 231-237. - doi: [10.1051/forest:2006001](https://doi.org/10.1051/forest:2006001)
- Timmer VR, Stone EL (1978). Comparative foliar analysis of young balsam fir fertilized with nitrogen, phosphorus, potassium, and lime. *Soil Science Society of America Proceedings* 42: 125-130. - doi: [10.2136/sssaj1978.0361599500420010027x](https://doi.org/10.2136/sssaj1978.0361599500420010027x)
- Timmer VR (1997). Exponential nutrient loading: a new fertilization technique to improve seedling performance on competitive sites. *New Forests* 13: 279-299. - doi: [10.1023/A:1006502830067](https://doi.org/10.1023/A:1006502830067)
- von-Fircks Y, Sennerby-Fores L (1998). Seasonal fluctuations of starch in root and stem tissues of coppiced *Salix viminalis* plants grown under two nitrogen regimes. *Tree Physiology* 18: 243-249. - doi: [10.1093/treephys/18.4.243](https://doi.org/10.1093/treephys/18.4.243)
- Villar-Salvador P, Puértolas J, Cuesta B, Peñuelas JL, Uscola M, Heredia-Guerrero N, Rey BJM (2012). Increase in size and nitrogen concentration enhances seedling survival in Mediterranean plantations. Insights from an ecophysiological conceptual model of plant survival. *New Forests* 43: 755-770. - doi: [10.1007/s11056-012-9328-6](https://doi.org/10.1007/s11056-012-9328-6)
- Wang L, Yan Z, Li J, Wang J, He Q, Su Y, Dong J (2013). Effects of nitrogen exponential fertilization on photosynthetic characteristics of three *Catalpa bungei* clones. *Forest Research* 26 (1): 46-51.
- Wilson BF (2000). Apical control of branch growth and angle in woody plants. *American Journal of Botany* 87 (5): 601-607. - doi: [10.2307/2656846](https://doi.org/10.2307/2656846)
- Wright SJ, Yavitt JB, Wurzburger N, Turner BL, Tanner EVJ, Sayer EJ, Santiago LS, Kaspari M, Hedin LO, Harms KE, Garcia MN, Corre MD (2011). Potassium, phosphorus or nitrogen limit root allocation, tree growth and litter production in a lowland tropical forest. *Ecology* 92 (8): 1616-1625. - doi: [10.1890/10-1558.1](https://doi.org/10.1890/10-1558.1)
- Zida D, Tigabu M, Sawadogo L, Oden PC (2008). Initial seedling morphological characteristics and field performance of two sudanian savanna species in relation to nursery production period and watering regimes. *Forest Ecology and Management* 255: 2151-2162. - doi: [10.1016/j.foreco.2007.12.029](https://doi.org/10.1016/j.foreco.2007.12.029)