

Relationship between tree growth and physical dimensions of *Fagus sylvatica* crowns assessed from terrestrial laser scanning

Dominik Seidel⁽¹⁻²⁾, Peter Schall⁽²⁾,
Malte Gille⁽²⁾, Christian Ammer⁽²⁾

Measurements of physical tree crown dimensions were of subjective character in the past, even though they can be considered important for the management of many silvicultural operations, such as timing of thinning operations. In our study we investigated if and how measures of physical crown dimensions of trees differed when quantified conventionally versus based on 3D-terrestrial laser scanning and how they are related to basal area increment. Some 24 randomly selected predominant or dominant beech trees between 90 and 110 yrs of age and of varying height were used as study trees. We hypothesized that tree crown dimensions obtained from scans are more closely related to tree radial growth than those obtained from conventional field measurements. It was found that from a variety of compared crown size characteristics the scan-based tree attributes mean crown radius, maximum area of the crown and crown projection area were most closely related to individual tree growth. We conclude that the horizontal extension of a tree crown in general is to be considered one of the most important drivers of tree growth. We also conclude that terrestrial laser scanning is a powerful tool to reliably measure physical crown dimensions and TLS-based measurements are more reliable than conventional ones.

Keywords: Tree Geometry, Crown Structure, Crown Surface Area, Basal Area Increment

Introduction

More than 200 years of silviculture provided knowledge on how a forest stand should be managed to produce wood sustainably. This knowledge is based on fundamental relationships, such as between tree crown size and diameter increment, indicating the importance of canopy structure for tree and stand productivity (Dieler & Pretzsch 2013). However, not only biomass production but also other forest ecosystem services and functions, for instance habitat suitability, do very much depend on structural properties of trees and hence on management-related changes of tree and forest structure (Puettmann et al. 2009, Gossner et al. 2014, Winter et al. 2015). Understanding the interplay between forest management and the resulting tree

habitus or growth is crucial for an effective and goal-orientated forest management (Röhrig et al. 2006, Bayer et al. 2013). As tree crown size is closely related to light absorption and productivity (Binkley et al. 2013) reliable estimations of tree crown dimensions are essential for any kind of tree growth modeling. Recent research has revealed that crown size, beside species, tree size and site fertility, is determined by neighborhood density and identity (Pretzsch & Schütze 2009, Dieler & Pretzsch 2013, Metz et al. 2013). For a quantification of the impact of these factors an estimation of the overall tree dimensions, especially of crown size, is required and should be as precise as possible.

Three main characteristics of trees have hindered a detailed acquisition of their

geometry so far: (i) the sheer complexity and overall size (Seidel et al. 2011a); (ii) the dynamic as living organism (Andrejczyk & Brzeziecki 1995); and (iii) the long lifespan (Pretzsch 2010). Therefore, easily accessible attributes of sheer tree size belonged to the most often used data source in forest research of the past. Probably the most important two measures are diameter at breast height (DBH) and total tree height. The combined use of two or more such attributes to predict tree characteristics that are more difficult (or even impossible) to measure is common practice in forest science. Examples are characteristics such as leaf area, crown surface area (Kramer & Akça 1987, Metz et al. 2013) or tree biomass (Hunter et al. 2013).

For a very long time instruments specifically developed to measure a tree's structural attributes were used in forest inventories, such as the Blume-Leiss altimeter for measuring tree height or Wheeler's pentaprism for the determination of upper diameters, to name only two. Several books reviewed the available instruments and the range of possible applications (Van Laar & Akça 2007). Many of these instruments were constantly improved to be more precise, more objective or simply more efficient. The utilization of laser technology (light amplification through stimulated emission of radiation – abbr.: “laser”) is one of these current improvements. Today, laser technology is implemented in range finders that include functions to

□ (1) Department of Forest Ecosystems and Society, College of Forestry, Oregon State University, 321 Richardson Hall, Corvallis, 97331 OR (USA); (2) Department of Silviculture and Forest Ecology of the Temperate Zones, Faculty of Forest Science and Forest Ecology, University of Göttingen, Büsgenweg 1, 37077 Göttingen (Germany)

@ Dominik Seidel (dseidel@gwdg.de)

Received: Jan 17, 2015 - Accepted: May 04, 2015

Citation: Seidel D, Schall P, Gille M, Ammer C (2015). Relationship between tree growth and physical dimensions of *Fagus sylvatica* crowns assessed from terrestrial laser scanning. *iForest* 8: 735-742. - doi: 10.3832/ifor1566-008 [online 2015-06-11]

Communicated by: Chris Eastaugh




Fig. 1 - Exemplary two-dimensional representation of the point cloud of a beech tree (32 m in height).

measure tree height as well as in laser scanners.

Three-dimensional (3D) terrestrial laser scanning is a technology with proven potential for deriving aboveground tree structural attributes. A comprehensive overview of its various applications is given in Maas (2010) and Dassot et al. (2011). Especially, terrestrial laser scanning (TLS) has been demonstrated as a very effective tool in measuring tree/stand properties not directly measurable which are of interest in forest research, such as crown displacement (Seidel et al. 2011b), competition among trees (Metz et al. 2013), leaf area distributions (Béland et al. 2011) or crown volume and surface (Metz et al. 2013). Furthermore, a detailed assessment of branch dimensions became possible by means of 3D-TLS (Bucksch et al. 2010, Dassot et al. 2012).

Here we hypothesize that high resolution spatial information on tree geometry obtained from 3D-TLS also increases the accuracy of measures of tree morphology, which in turn improve models that relate

tree morphology to tree growth. While manifold applications of TLS in forest research have been successfully tested, the use of comprehensive 3D-data from TLS-based measurements of tree crowns is still in an early stage. Tree crowns have mostly been scanned to measure gap fraction or canopy cover (see the review by Dassot et al. 2011). Comparisons with conventional forest inventory measurements focusing on crowns have been rare. Fleck et al. (2007, 2011) showed that conventional eight-point crown projections with fixed angles to capture a tree's horizontal crown extension differed substantially from crown projection areas obtained from TLS measurements. However, no further analysis was made, for example by relating the crown projection areas to tree growth. We conclude that the potential to explore physical tree dimensions by means of TLS is not yet fully exploited and further research in this field could contribute to our understanding of geometry-related processes that take place at all three dimensions of a forest, with growth being an important example.

In this paper we investigated if and how measures of tree size differ when quantified conventionally vs. TLS-based. We hypothesized that tree attributes calculated using measurements of tree crown dimensions obtained from TLS are more closely related to tree growth than those obtained from conventional field measurements. Specifically, we compared TLS-derived values of the mean crown radius (CR_{mean}) as a surrogate for the complex geometric measure of crown surface area (CSA). CSA has successfully been used as an estimator of tree growth performance in the past (Kramer & Akça 1987).

Methods

Study site and study objects

Our study trees were located in the municipality forest of Lübeck, northern Germany, and were distributed over two different unmanaged forested areas named Hevenbruch ($53^{\circ} 38' 41''$ N, $10^{\circ} 32' 53''$ O) and Schattiner Zuschlag ($53^{\circ} 46' 49''$ N, $10^{\circ} 47' 57''$ O). The forests are dominated by oak (*Quercus robur* L. and *Quercus petraea* [Muttuschka] Liebl.) and beech (*Fagus sylvatica* L.), which contribute to about 45% of the total number of stems, according to forest inventory data of 2004. We randomly selected 24 predominant or dominant beech trees between 90 and 110 yrs in age and of varying height (mean: 32.9 m, standard deviation – SD: 5.12 m; measured in 2013) and DBH (mean: 53.48 cm, SD: 13.23 cm; measured in 2013). The stands have not been subject to any tree removals by harvests for at least 20 years and are fully stocked (stocking level: Schattiner Zuschlag = 1.38; Hevenbruch = 1.19).

Conventional forest inventory

DBH of all trees was measured in 2007

and in 2013 using a diameter tape. Tree growth was determined as basal area increment (BAI) between 2007 and 2013. Additionally, in 2013 we determined total tree height (TTH) and crown base height (CBH – lower-most branch of the live crown) using a Vertex (Haglof Inc., Madison, USA) and crown radius using a crown mirror (densiometer). The edge of the crown at its largest and shortest extension and at four cardinal azimuthal directions (North, East, South and West) was determined and the distance to the stem base was obtained from a standard measuring tape. The quadratic mean (CR_{mean}) was then calculated based on the six measurements and used to determine the crown projection area (CPA) based on the formula of the area of the circle ($CPA = \pi \cdot CR_{mean}^2$). CSA was calculated as the sum of the area of the crown exposed to sunlight (CSA_{sun}) and the shaded part of the crown (CSA_{shade}). CSA_{sun} was calculated based on a standard crown shape model using the formula of the surface of a paraboloid, in accordance to Dong & Kramer (1985 – eqn. 1):

$$CSA_{sun} = \frac{\pi CR_{mean}}{6 CL_{sun}^2} [(4 \cdot CL_{sun}^2 + CR_{mean}^2)^{\frac{3}{2}} - CR_{mean}^3]$$

where CL_{sun} is the length of the crown exposed to sunlight. Based on the extensive dataset on beech trees presented by Pretzsch et al. (2002) and Pretzsch (1992, 2001), these authors created a widely used beech crown shape model in which CL_{sun} was set to $0.4 \cdot CL$ (standard crown shape model). We used this model for our analysis. CL was given as difference between TTH and CBH . The area of the shaded surface of the crown (CSA_{shade}) was determined as the lateral surface of a frustum (eqn. 2):

$$CSA_{shade} = (CR_{mean} + CR_{CBH}) \cdot \pi \cdot m$$

with m describing the slant height of the frustum and CR_{CBH} being the diameter of the crown at CBH ($0.33 \cdot CR_{mean}$, according to Pretzsch 2010 - beech standard crown shape model). Consequently, the height of the maximal horizontal crown extension ($H_{maxarea}$) in this standard model is assumed to be located at the base of the sun crown (Metz et al. 2013 – eqn. 3):

$$H_{maxarea} = TTH - (0.4 \cdot CL)$$

The standard crown shape model also assumes that the area of maximum horizontal crown extension is always equal to the crown projection area. This is due to the perfect circularity of the paraboloid (sun-exposed crown) and frustum (shaded crown) that are used in the crown shape model. The interested reader is referred to Metz et al. (2013) for an optical comparison of the simple crown shape model and laser derived (more realistic) tree shapes of beech trees. In reality, it is certainly possi-

ble that a tree does have its maximum horizontal crown extension in a height which is different from the transition between sun-exposed and shaded crown. However, detecting H_{maxarea} is impossible based on conventional field measurements because the vertical dimension is not considered during crown delineations with a crown mirror. Our data enables for a comparison of values obtained from the literature (H_{maxarea}) with those obtained from TLS-data, in which the vertical dimension was taken into account (see below).

Terrestrial laser scanning

Terrestrial laser scans were made in March 2013 (leaf-off conditions) using the Z+F Imager 5006 (Zoller and Fröhlich GmbH, Wangen, Germany) in order to minimize obstruction effects in the crown caused by leaves. The Imager 5006 is a phase-based 3D-terrestrial laser scanner operating with laser light of a wavelength between 650-690 nm. We used a field of view of 310° in vertical and 360° in horizontal direction, scanned with an angular step width of 0.036° (10 000 point per 360° ; point spacing of 6.3 mm at 10 m from the scanner). The maximum range of the instrument (ambiguity interval) was given as 79 m. The scanner was moved to the field sites and mounted on a tripod at breast height (~1.3 m). At each site a first scan (the so called master scan) was made in the vicinity of the tree of interest. Multiple scans were then consecutively made around the tree. Artificial reference targets (checkerboards) distributed in the scanned forest scenes enabled the co-registration of the scans. This kind of scan design with flexible scan position was successfully used in other studies and provided reliable measurements of tree attributes in the past (Seidel et al. 2011b, Metz et al. 2013), even though obstruction effects may still exist in the data to a certain extent.

Data processing

The scan data of each scan session made at a study tree was semi-automatically registered based on the artificial targets, filtered and exported as a single pts file (point cloud data) according to Seidel et al. (2012) using the ZF Laser Control software (Zoller and Fröhlich GmbH, Wangen, Germany). Pts files were used in Cyclone 6 (Leica Geosystems, Heerbrugg, Switzerland) to manually select each tree from the point cloud data, as described by Seidel et al. (2011b). To our knowledge, there is no alternative automated procedure available that performs tree separation for adult trees in dense forest stands with greater accuracy than the manual procedure (Metz et al. 2013). At this stage of the post-processing, all study trees are available with individual three-dimensional descriptions of their habitus based on a point cloud with sub-centimeter resolution. The data was transformed to xyz files (Fig. 1 for visualization) which were then used as

Tab. 1 - Structural attributes used in the present study. [] = Parameters not specifically mentioned in the text but needed to calculate CSA_{TLS} .

Conventional measurement		TLS-based measurement	
Param.	Measurement description	Param.	Measurement description
DBH	Tape; stem circumference at 1.3m above ground	DBH_{TLS}	QR-decomposition procedure to fit a circle to all points at 1.3 m above ground (see Seidel et al. 2011b)
TTH	Vertex; highest point of the tree	TTH_{TLS}	Vertical distance between highest and lowest point of the point cloud
CBH	Vertex; height of the lowermost live-branch	CBH_{TLS}	See Metz et al. (2013)
CR_{mean}	Densimeter; mean of six distance measurements (N, W, S, E, longest, shortest) from stem base to crown edge	CR_{meanTLS}	Mean of the horizontal distances between stem base and crown edge in 360 directions
CPA	$\pi \cdot CR_{\text{mean}}^2$	CPA_{TLS}	Projected area of the convex hull of the crown (on the ground)
CL	$TTH - CBH$	CL_{TLS}	$TTH_{\text{TLS}} - CBH_{\text{TLS}}$
CR_{CBH}	$0.33 \cdot CR_{\text{mean}}$ (according to Pretzsch 2010)	$[CR_{\text{CBHTLS}}]$	$0.33 \cdot CR_{\text{meanTLS}}$ (according to Pretzsch 2010)
CL_{sun}	$0.4 \cdot CL$ (according to Pretzsch 1992)	$[CL_{\text{sun}}]$	$0.4 \cdot CL_{\text{TLS}}$ (according to Pretzsch 1992)
CSA_{sun}	See eqn. 1	$[CSA_{\text{sunTLS}}]$	See eqn. 1 (but with TLS input parameters)
CSA_{shade}	$(CR_{\text{mean}} + CR_{\text{CBH}}) \cdot \pi \cdot m$; with m describing the slant height of the frustum	$[CSA_{\text{shadeTLS}}]$	$(CR_{\text{meanTLS}} + CR_{\text{CBHTLS}}) \cdot \pi \cdot m$; with m describing the slant height of the frustum
CSA	Sum of CSA_{shade} and CSA_{sun}	CSA_{TLS}	Sum of CSA_{shadeTLS} and CSA_{sunTLS}
H_{maxarea}	$TTH - (0.4 \cdot CL)$ (according to Pretzsch 1992, Pretzsch 2010)	$H_{\text{maxareaTLS}}$	Height of the polygon used for Max_{AreaTLS} (see Seidel et al. 2011b)
-	-	CV_{TLS}	See Metz et al. 2013
-	-	Max_{AreaTLS}	Area of the largest crown layer polygon (10-cm layers - see Seidel et al. 2011b)
-	-	$CR_{\text{meanTLSrotated}}$	Mean of 89 rotations around the vertical axis for a virtual CR_{mean} measurement (based on horizontal distance between stem base and crown edge)

inputs for an algorithm implemented in the computational software program Mathematica® (Wolfram Research Inc., Champaign, USA).

Calculation of tree structural properties

Using the above algorithm, several physical attributes were calculated using a fully automatic procedure from the xyz files of all study tree individuals. Diameter at breast height (DBH_{TLS}), total tree height (TTH_{TLS}) and maximum area of the crown as measured in a certain height stratum (Max_{areaTLS}) were calculated as described in Seidel et al. (2011b). In addition, we derived crown base height (CBH_{TLS}), crown volume (CV_{TLS}) and height of the maximum crown extension ($H_{\text{maxareaTLS}}$) as described in Metz et al. (2013). Crown surface area (CSA_{TLS}) was calculated according to eqn. 1 and eqn. 2, but this time based on the TLS derived input variables. Crown length (CL_{TLS}) was calculated in the same way as for the conventional field data ($TTH_{\text{TLS}} - CBH_{\text{TLS}}$). Crown projection area (CPA_{TLS}) was derived from the convex hull of the tree crown by projecting all crown height sections to the ground and calculating their combined

area. Thereby, we used a 10-cm voxel model of the tree crown (only voxels above CBH_{TLS}). Finally, two versions of the mean crown radius were derived from the TLS-data. First, we calculated CR_{meanTLS} based on the mean of 360 virtual measurements of crown radius, one in each azimuth direction in steps of one degree, using the stem base point as starting point and calculating the horizontal distance to the edge of the crown in the voxel model (outermost part of the crown). Secondly, we conducted a simulation of the field measurement scheme with four cardinal directions (North, West, South and East) plus the longest and shortest radius. This measurement scheme was then virtually rotated around the vertical axes in 89 steps of one degree, resulting in 90 readings of CR_{mean} , hereafter referred to as $CR_{\text{meanTLSrotated}}$. All these measurements were based on the trees' voxel models ($10 \times 10 \times 10$ cm) obtained from scanning. Tab. 1 summarizes the structural attributes used in our study.

Acquiring all information on a target tree with the terrestrial laser scanning took about 45 minutes in the field plus additional 60 minutes for the post-processing,




Fig. 2 - (Left): Relationship between the *DBH* of a tree measured by a conventional tape method and *DBH* determined via TLS. (Right): Relationship between the *TTH* obtained from vertex measurements and *TTH* derived from TLS. The 1:1 ratio is represented by the dashed line.




Fig. 3 - (Left): Relationship between the *CBH* of a tree measured with the vertex instrument and *CBH* determined mathematically from TLS-data. (Right): Relationship between the *CL* of a tree derived from the information on *TTH* and *CBH* measured with conventional vertex method and *CL*-values calculated via TLS.




Fig. 4 - (Left): Relationship between $H_{\max\text{area}}$ obtained with literature-based ratios of sun-exposed to shaded portions of the crown and TLS-based measurements. (Right): Relationship between CR_{mean} and CR_{meanTLS} . The 1:1 ratio is represented by the dashed line.

with the manual tree extraction from the point cloud taking most of the time (approx. 20-30 min per tree). The calculation of all tree structural attributes in Mathematica® took no more than 2-3 min per tree.

Statistics

In order to compare the results obtained from the conventional field measurements with TLS-based information, we performed correlation analyses based on linear regressions using the statistical software R (Vers.

3.0.0. - R Development Core Team). We waived using one of the two methods (scan or conventional measurement) as reference. Instead, we decided to focus on the strength of the relationship between measurements and preceding tree growth. The retrospective approach was used to ensure comparability among the approaches as TLS was unavailable in the 2007 inventory. We compared different competing linear models of tree growth, here expressed as 5-yrs increment in the trees' basal area (*BAI* - 2007-2013), based on the Akaike Information Criterion (AICc) according to Burnham & Anderson (2002). In the models tree growth was related to a single tree structural attribute or an interaction of two. In case of the traditional field inventory data, we only used data of 2013 in order to allow comparability with TLS data of the same year.

Results

Comparison of *TTH* and *DBH* values calculated from the TLS-approach with those obtained from traditional field instruments (vertex and tape) yielded high coefficients of determination (Fig. 2). In case of *DBH*, the results showed marginal differences and the linear regression of the measurements obtained from the two approaches was almost identical to the 1:1 line. A mean absolute difference ($|DBH - DBH_{\text{TLS}}|$) of 0.5 cm was found for the TLS measurements.


For the vertex measurements of *TTH* we obtained larger differences to the scanned data, with a mean difference of 2.4 m between the two approaches ($TTH - TTH_{\text{TLS}}$). The mean absolute difference was 2.6 m.

The coefficient of determination for the relationship between vertex-based and TLS-based measurements of *CBH* was 0.7 and the linear regression was statistically significant with $p < 0.001$ (Fig. 3, left panel). The mean absolute difference between measurements of the two approaches ($|CBH - CBH_{\text{TLS}}|$) was 2.8 m. This is not surprising taking into account that different definitions of *CBH* were used, with one being strictly mathematical (CBH_{TLS} as presented in Metz et al. 2013) and the other one (*CBH*) being a field operator's estimate using the Vertex.

CL was derived from *TTH* and *CBH* and hence differences in *CL* were directly dependent on the accordance or discrepancy of these two attributes. A significant linear regression ($p < 0.01$) with a very low coefficient of determination ($R^2 = 0.29$) was found for the relationship between the values obtained from the two different methods. The TLS-based approach resulted in much lower estimates of *CL* when compared to the conventional data, especially for crowns with a large vertical extension (Fig. 3 right). Absolute differences between the two approaches reached up to 10 m.

Interestingly, we found a clear tendency towards lower height of maximum crown

Fig. 5 - Comparison of CR_{meanTLS} , CR_{mean} and $CR_{\text{meanTLsrotated}}$ (rotation around vertical) based on the TLS data.



extension when based on scans ($H_{\text{maxareaTLS}}$) compared to the geometrical crown shape model (H_{maxarea}) (Fig. 4, left panel). Mean absolute difference between the two approaches was 3.0 m and the relationship between both variables was statistically significant ($p < 0.001$) using a linear regression ($R^2 = 0.78$).

A coefficient of determination of 0.6 was found for the relationship between CR_{mean} measured in the field (six azimuthal directions) and the TLS-based measurement (360 azimuthal directions). The scatter plot (Fig. 4, right panel) clearly shows that conventional crown radius measurements resulted in higher values than the TLS-based measures. This was confirmed by the mean difference ($CR_{\text{mean}} - CR_{\text{meanTLS}}$) being 1.4 m.

As displayed in Fig. 5, conventional field based measurements of the mean crown radius (CR_{mean} , indicated by an “X” in the figure) using a densitometer yielded generally higher values than those obtained from scan based simulations of the conventional measurements scheme (Box-Whisker plot of $CR_{\text{meanTLsrotated}}$) or those obtained from 360 measurements in the virtual tree crown (CR_{meanTLS} , indicated by the black dots). Interestingly, CR_{meanTLS} was the more likely to be higher than $CR_{\text{meanTLsrotated}}$ the larger the crown radius was, whereas the opposite was true for trees with small crown radii (Fig. 5).

Values of CPA and CPA_{TLS} were correlated ($R^2 = 0.67$) and yielded a statistically significant linear regression ($p < 0.001$ - Fig. 6, left panel). Crowns of large trees tended to have higher CPA values when calculated conventionally compared to TLS-based estimations that rely on the convex hull of the polygon enclosing the projected area of the crown voxels (precisely, their center coordinates). The mean difference between the two approaches was 9.4 m^2 ($CPA - CPA_{\text{TLS}}$).

CSA values calculated based on TLS data




Fig. 6 - (Left): Relationship between the CPA and a CPA_{TLS} . (Right): Relationship between CSA calculated from conventional field data and determined based on TLS data. The 1:1 ratio is represented by the dashed line.

Tab. 2 - Comparison of models relating the basal area increment (2007-2013) to different tree size attributes (nine measured in 2013 and one in 2007 [initial DBH]). (W_{ic}): AICs weights (Burnham & Anderson 1998); (***) : $p < 0.001$; (**) : $p < 0.01$; (*) : $p < 0.05$.

Tree structural attribute	AICc	ΔAICc	R^2	Level of significance	W_{ic}
CR_{meanTLS}	234.18	0.00	0.44	***	0.32
Max_{areaTLS}	234.35	0.17	0.44	***	0.29
CPA_{TLS}	234.96	0.78	0.42	***	0.21
CSA	237.82	3.64	0.33	**	0.05
CSA_{TLS}	238.94	4.76	0.29	**	0.03
CV_{TLS}	238.98	4.80	0.29	**	0.03
CR_{mean}	239.14	4.96	0.29	**	0.03
CPA	239.75	5.57	0.27	*	0.02
$DBH_{(2007)}$	240.11	5.93	0.25	*	0.02
TTH_{TLS}	244.44	10.26	0.07	n.s.	0.00

and those calculated from traditional field data yielded a statistically significant linear regression ($p < 0.001$) and an intermediate coefficient of determination ($R^2 = 0.48$; Fig. 6 right). The mean absolute difference between crown surface areas calculated from the two approaches was 167.5 m^2 , with the TLS-based values being usually much smaller than the field-based values.

We found that various crown attributes

were closely related to preceding tree diameter growth. Models with interactions and models with more than just one attribute did not result in a stronger relationship than models using only the attribute that showed the strongest relation to growth (results not shown). Closest relation to 5-year basal area increment was given by three simple tree structural attributes obtained from TLS, with the fol-

lowing order in model quality: CR_{meanTLS} , Max_{areaTLS} and CPA_{TLS} (Tab. 2). The conventionally calculated attributes CPA and CR_{mean} were less closely related to tree growth than the respective laser-derived attributes, which is indicated by the differences in AIC (ΔAICc). The strength of the relationship with growth was found to be similar for CSA and CSA_{TLS} (Tab. 2).

Discussion

As tree crown size is closely related to light absorption and hence productivity (Binkley et al. 2013), reliable estimations of tree crown dimensions are essential for any kind of tree growth modeling. Validation of the reliability of structural attributes describing crown dimensions derived from TLS data has been difficult ever since this technology was used in forest research. While attributes such as DBH can be validated based on reliable field data, tree attributes more difficult to measure such as TTH or CR_{mean} may not necessarily be accepted as “more correct” or “closer to the real value” when measured in the field. We decided to avoid this problem using any of the two datasets as “true” reference. Instead, we used information from both approaches to compare how they are related to preceding tree growth.

Tree structural attributes obtained from TLS were more closely related to preceding tree growth than those measured in the field. If spatially detailed information on the crown dimensions is available from TLS, such as CR_{meanTLS} calculated from 360 measurements, estimations of CSA_{TLS} that rely on simple geometric models seem to offer no advantages. Analyzing the same sites considered in our study, Fichtner et al. (2013) showed that tree growth is related to CSA ($R^2 > 0.37-0.63$), though a tree crown model developed only for the sun-exposed crown was mistakenly applied to the whole crown, thereby ignoring the usually assumed differences in shape between the sun-exposed and the shaded part, as presented here and elsewhere (Pretzsch et al. 2002, Pretzsch 2010).

In our study we confirmed that CSA seems a suitable proxy for tree growing performances. However, we demonstrated that crown structural attributes obtained from sophisticated measurements are even more closely related to tree growth. Obtaining high resolution data on crown structural attributes would be very laborious if carried out without the use of TLS. Earlier studies showed that such data can also be obtained by means of TLS for other tree species, including coniferous species (Seidel et al. 2011b, Bayer et al. 2013, Metz et al. 2013). Such data can be used to explain tree growth with higher-than-usual model quality. By conducting 360 measurements of crown radius in the virtual environment of a point cloud, a more precise mean value could be calculated when compared to six field measurements. Fig. 5 showed that field measurements based on

six directions (N, E, S, W, shortest, longest) usually result in higher values of CR_{mean} as compared with TLS estimates. We explain this by the strong influence of the longest crown radius, accounting for the sixth part of the mean value. While the length of the shortest radius can be zero the longest radius can be very large, e.g., due to a strong branch leaning out from the crown center. In the case of 360 measurements based on TLS data, the effect of such a branch on CR_{mean} is much lower. In such a case, one measurement represents only a 360th of the mean, and only a branch with an azimuthal extent of 60° would have the same weight on the mean, which is quite unlikely. However, field-based estimations of the crown radius were generally larger than TLS-based estimations from simulations of the same measurement scheme (see the Box-Whisker plots in Fig. 5). Field measurements should be in the range of the Whiskers if the measurement of the distance between the stem and the edge of a branch in a given direction would yield similar results. This is the case only for two out of 24 trees. We argue that this might be due to difficulties to accurately delineate the end of a branch in the field, especially when trees are very tall. Furthermore, in the case of the TLS-based delineation, small branches and their tiny tips may remain undetected, thus tree crown radii might be underestimated to a certain degree. This effect would be larger in the case of very tall trees, due to the decreasing scan resolution with distance from the scanner. As some of our study trees reached heights greater than 40 m, we expect this error to be present in our dataset, even though we used a 10-cm voxel model.

CSA and CV_{TLS} were the only structural characteristics based on all three dimensions of the tree crown. The first aims at describing the production potential of trees in terms of photosynthesis. Describing the surface area of the tree exposed to sunlight seems reasonable when growth is to be related to crown dimensions. Indeed, the area described by CSA does not represent the crown photosynthetic activity, as neighboring trees cast their shadow on this area and the ratio of leaves adapted to direct sunlight to those adapted to shade is unknown. We hypothesize that CSA values are closely related to growth as they are largely determined by CR_{mean} . Our study showed that attributes describing the horizontal extension of tree crowns were closely related to growth.

In contrast, CV_{TLS} describes the overall three-dimensional extent of the tree crown, with its value being determined by all spatial irregularities of the crown shape both in vertical and horizontal directions, but with no dominant control by the horizontal extent. The three models with the strongest relationship to tree growth were all based on attributes describing the crown width (CR_{meanTLS} , Max_{areaTLS} , CPA_{TLS}). This finding is in line with studies indicating

that trees released from competition tend to increase their crown width and length but not their leaf area density (Forrester et al. 2013).

The horizontal extent of tree crown is often described via the mean crown radius or the crown projection area, which is calculated from the mean radius. In most cases the mean crown radius itself is averaged from a small number of measurements. As mentioned by Fleck et al. (2011) “the most often used resolution [...] is an eight-point crown projection, since four-point projections (into North, East, South, and West direction) are insufficient for most purposes and 16-point approximations require too much time for large-scale investigations”. However, according to Röhle & Huber (1985) eight radii seem to be a minimum for reliable measurements. In our study, the use of six azimuthal directions led to the overestimation of tree crown dimensions (CR_{mean} , CPA) as compared with detailed measurements of crown radius from TLS (360 azimuthal directions), which holds especially for large trees (see Fig. 4, right panel and Fig. 6). Since crown dimensions of such trees are overestimated, their diameter growth may mistakenly be interpreted as lower than expected, which may lead to misinterpretations. In their study at the same sites, Fichtner et al. (2013) concluded that the growth efficiency of beech trees in managed forests, which had been released from competition and which were characterized by large crowns, was lower than that of trees with shallow crowns in an unmanaged stand. Actually, no difference in mean growth efficiency (basal area increment per m² crown projection area) between managed and unmanaged trees exists if the regression equation in Fig. 6 is used for calculating the “true” crown projection areas for the trees investigated in Fichtner et al. (2013). This is in line with the comprehensive study of Pretzsch & Schütze (2009) who found that dominant beech at a given age does hardly change its crown efficiency but immediately starts to occupy new space by enlarging its crown. The high plasticity of beech to effectively explore the canopy space was confirmed by Schröter et al. (2012), who found that the “direction of crown displacement” of a given tree was strongly dependent on the distance from neighbors.

Conclusions

In the present study we compared conventional measures of crown structural attributes with more sophisticated, spatially-explicit measurements based on high resolution point clouds obtained from TLS. Although the accordance between attributes derived from the two approaches was high in general if addressed via correlation analysis, we found large differences for several parameters, especially CR_{mean} and CR_{meanTLS} . A comparison of linear models of tree growth revealed that crown

radius is the attribute with the strongest relationship to tree growth, followed by the two closely related attributes $Max_{areaTLS}$ and CPA_{TLS} . We conclude that the horizontal extension of tree crown has to be considered as one of the most important drivers of tree growth.

From a silvicultural point of view, an early release from competition leads to crown enlargement, resulting not only in high growth responses (Pretzsch 2006) but also in more regular crowns (i.e., circular and centered on the stem - Longuetaud et al. 2013), which in turn reduce the whole stress field in a trunk resulting from tree growth (Jullien et al. 2013).

We demonstrated that terrestrial laser scanning is a powerful tool to measure physical crown dimensions. In this study TLS-based measurements were found more reliable than conventional field methods. Therefore, TLS data should be considered a suitable tool to validate crown radius measurement from airborne laser scanning (Hilker et al. 2010, Hauglin et al. 2014) and it may also be useful for comparisons with high spatial resolution above-canopy imagery. On the long term, airborne approaches for crown radius measurements are likely to be more widely applied to forest areas and they will benefit from detailed ground truth data by means of TLS.

Acknowledgements

We thank the administration of the municipality forest of Lübeck for the permission to scan the trees and for providing us with data on the 2007 forest inventory. We are also grateful to the helpful comments of three anonymous reviewers.

References

- Andrejczyk T, Brzeziecki B (1995). The structure of old-growth *Pinus sylvestris* stands in the Wigry National Park, Northeast Poland. *Vegetatio* 177: 81-95. - doi: [10.1007/BF00033261](https://doi.org/10.1007/BF00033261)
- Bayer D, Seifert S, Pretzsch H (2013). Structural crown properties of Norway spruce (*Picea abies* [L.] Karst.) and European beech (*Fagus sylvatica* [L.]) in mixed versus pure stands revealed by terrestrial laser scanning. *Trees* 27: 1035-1047. - doi: [10.1007/s00468-013-0854-4](https://doi.org/10.1007/s00468-013-0854-4)
- Béland M, Widłowski JL, Fournier RA, Côté JF, Verstraete MM (2011). Estimating leaf area distribution in savanna trees from terrestrial LiDAR measurements. *Agricultural and Forest Meteorology* 151 (9): 1252-1266. - doi: [10.1016/j.agrformet.2011.05.004](https://doi.org/10.1016/j.agrformet.2011.05.004)
- Binkley D, Campoe OC, Gspaltl M, Forrester DI (2013). Light absorption and use efficiency in forests: why patterns differ for trees and stands. *Forest Ecology Management* 288: 5-13. - doi: [10.1016/j.foreco.2011.11.002](https://doi.org/10.1016/j.foreco.2011.11.002)
- Bucksch A, Lindenbergh R, Menenti M (2010). SkelTre - Robust skeleton extraction from imperfect point clouds. *The Visual Computer* 26: 1283-1300. - doi: [10.1007/s00371-010-0520-4](https://doi.org/10.1007/s00371-010-0520-4)
- Burnham KP, Anderson DR (1998). *Model selection and inference: a practical information-theoretic approach*. Springer-Verlag, New York, New York, USA, pp. 353.
- Burnham KP, Anderson DR (2002). *Model selection and multi-model inference: a practical information-theoretic approach*. Springer Science & Business Media, New York, USA, pp. 488.
- Dassot M, Constant T, Fournier M (2011). The use of terrestrial LiDAR technology in forest science: application fields, benefits and challenges. *Annals of Forest Science* 68: 959-974. - doi: [10.1007/s13595-011-0102-2](https://doi.org/10.1007/s13595-011-0102-2)
- Dassot M, Colin A, Santenoise P, Fournier M, Constant T (2012). Terrestrial laser scanning for measuring the solid wood volume, including branches, of adult standing trees in the forest environment. *Computers and Electronics in Agriculture* 89: 86-93. - doi: [10.1016/j.compag.2012.08.005](https://doi.org/10.1016/j.compag.2012.08.005)
- Dieler J, Pretzsch H (2013). Morphological plasticity of European beech (*Fagus sylvatica* L.) in pure and mixed-species stands. *Forest Ecology and Management* 295: 97-108. - doi: [10.1016/j.foreco.2012.12.049](https://doi.org/10.1016/j.foreco.2012.12.049)
- Dong PH, Kramer H (1985). Beziehungen zwischen Zuwachs und Kronenparametern in immissionsgeschädigten Nadelholzbeständen [Relationship between tree growth and crown dimensions in coniferous stands affected by immissions]. *Dt. Verb. Forstl. Forsch. Anst., Kaelberbronn* 5: 1-25. [in German]
- Fichtner A, Sturm K, Rickert C, Oheimb G, Härdtle W (2013). Crown size-growth relationships of European beech (*Fagus sylvatica* L.) are driven by the interplay of disturbance intensity and inter-specific competition. *Forest Ecology and Management* 302: 178-184. - doi: [10.1016/j.foreco.2013.03.027](https://doi.org/10.1016/j.foreco.2013.03.027)
- Fleck S, Obertreiber N, Schmidt I, Brauns M, Jungkunst HF, Leuschner C (2007). Terrestrial LIDAR measurements for analysis canopy structure in an old-growth forest. *IAPRS* 36 (Part 3 / W52): 125-129. [online] URL: http://www.isprs.org/proceedings/XXXVI/3-W52/final_papers/Fleck_2007.pdf
- Fleck S, Mölder I, Jacob M, Gebauer T, Jungkunst HF, Leuschner C (2011). Comparison of conventional eight-point crown projections with LIDAR-based virtual crown projections in a temperate old-growth forest. *Annals of Forest Science* 68: 1173-1185. - doi: [10.1007/s13595-011-0067-1](https://doi.org/10.1007/s13595-011-0067-1)
- Forrester DI, Collopy JJ, Beadle CL, Baker TG (2013). Effect of thinning, pruning and nitrogen fertiliser application on light interception and light-use efficiency in a young *Eucalyptus nitens* plantation. *Forest Ecology and Management* 288: 21-30. - doi: [10.1016/j.foreco.2011.11.024](https://doi.org/10.1016/j.foreco.2011.11.024)
- Gossner MM, Schall P, Ammer C, Ammer U, Engel K, Schubert H, Simon U, Utschick H, Weisser WW (2014). Forest management intensity measures as alternative to stand properties for quantifying effects on biodiversity. *Ecosphere* 5 (9): art113. - doi: [10.1890/ES14-00177.1](https://doi.org/10.1890/ES14-00177.1)
- Hauglin M, Gobakken T, Astrup R, Ene L (2014). Estimating single-tree crown biomass of Norway spruce by Airborne Laser Scanning: a comparison of methods with and without the use of terrestrial laser scanning to obtain the ground reference data. *Forests* 5 (3): 384-403. - doi: [10.3390/f5030384](https://doi.org/10.3390/f5030384)
- Hilker T, Van Leeuwen M, Coops NC, Wulder MA, Newnham GJ, Jupp DL, Culvenor DS (2010). Comparing canopy metrics derived from terrestrial and airborne laser scanning in a Douglas-fir dominated forest stand. *Trees* 24 (5): 819-832. - doi: [10.1007/s00468-010-0452-7](https://doi.org/10.1007/s00468-010-0452-7)
- Hunter MO, Keller M, Vitoria D, Morton DC (2013). Tree height and tropical forest biomass estimation. *Biogeosciences Discussions* 10 (6): 10491-10529. - doi: [10.5194/bgd-10-10491-2013](https://doi.org/10.5194/bgd-10-10491-2013)
- Jullien D, Widmann R, Loup C, Thibaut B (2013). Relationship between tree morphology and growth stress in mature European beech stands. *Annals of Forest Science* 70: 133-142. - doi: [10.1007/s13595-012-0247-7](https://doi.org/10.1007/s13595-012-0247-7)
- Kramer H, Akça A (1987). *Leitfaden für Dendrometrie und Bestandesinventur* [Guidebook for dendrometry and forest inventory]. J. D. Sauerländer's Verlag, Frankfurt, Germany. pp. 358. [in German]
- Longuetaud F, Piboule A, Wernsdörfer H, Collet C (2013). Crown plasticity reduces inter-tree competition in a mixed broadleaved forest. *European Journal of Forest Research* 132 (4): 621-634. - doi: [10.1007/s10342-013-0699-9](https://doi.org/10.1007/s10342-013-0699-9)
- Maas H-G (2010). *Forestry Applications*. In: "Airborne and terrestrial laser scanning" (Vosselman G, Maas H-G eds). Whittles Publishing, Caithness, Scotland, UK, pp. 318.
- Metz J, Seidel D, Schall P, Scheffer D, Schulze ED, Ammer C (2013). Crown modeling by terrestrial laser scanning as an approach to assess the effect of aboveground intra- and interspecific competition on tree growth. *Forest Ecology and Management* 310: 275-288. - doi: [10.1016/j.foreco.2013.08.014](https://doi.org/10.1016/j.foreco.2013.08.014)
- Puettmann KJ, Coates KD, Messier C (2009). *A critique of silviculture: managing for complexity*. Island Press, Washington, USA, pp. 189. [online] URL: <http://journals.cambridge.org/prod/production/action/cjoGetFulltext?fulltextid=6686580>
- Pretzsch H (1992). *Konzeption und Konstruktion von Wuchsmodellen für Rein- und Mischbestände* [Concept and design of growth models for monocultures and mixed stands]. *Forstliche Forschungsberichte München* 115, Forstwissenschaftliche Fakultät der Universität München und der Bayer, Forstliche Versuchs- und Forschungsanstalt, München, Germany. [in German]
- Pretzsch H (2001). *Modellierung des Waldwachstums* [Modelling forest growth]. Blackwell Wissenschaftsverlag GmbH, Berlin/Wien, pp. 357. [in German]
- Pretzsch H, Biber P, Dursky J (2002). The single tree-based stand simulator SILVA: construction, application and evaluation. *Forest Ecology and Management* 162 (1): 3-21. - doi: [10.1016/S0378-1127\(02\)00047-6](https://doi.org/10.1016/S0378-1127(02)00047-6)
- Pretzsch H (2006). Von der Standflächeneffizienz der Bäume zur Dichte-Zuwachs-Beziehung des Bestandes. Beitrag zur Integration von Baum- und Bestandesebene [From growth efficiency of a tree to the density-growth relationship of stands. Contribution to the combined consideration of tree- and stand level]. *Allgemeine Forst- und Jagdzeitung* 177: 188-199. [in German]
- Pretzsch H, Schütze G (2009). Transgressive overyielding in mixed compared with pure stands of Norway spruce and European beech

- in Central Europe: evidence on stand level and explanation on individual tree level. *European Journal of Forest Research* 128: 183-204. - doi: [10.1007/s10342-008-0215-9](https://doi.org/10.1007/s10342-008-0215-9)
- Pretzsch H (2010). *Forest dynamics, growth and yield*. Springer, Berlin, Germany, pp. 664.
- Röhle H, Huber W (1985). Untersuchungen zur Methode der Ablotung von Kronenradien und der Berechnung von Kronengrundflächen [On the method of crown radius delineation and calculation of crown projection area]. *Forstarchiv* 56: 238-243. [in German]
- Röhrig E, Bartsch N, Lüpke B (2006). *Waldbau auf ökologischer Grundlage [Ecological silviculture]* (7th edn). Ulmer, Stuttgart, Germany, pp. 479. [in German]
- Schröter M, Härdtle W, Von Oheimb G (2012). Crown plasticity and neighborhood interactions of European beech (*Fagus sylvatica* L.) in an old-growth forest. *European Journal of Forest Research* 131: 787-798. - doi: [10.1007/s10342-011-0552-y](https://doi.org/10.1007/s10342-011-0552-y)
- Seidel D, Fleck S, Leuschner C, Hammett T (2011a). Review of ground-based methods to measure the distribution of biomass in forest canopies. *Annals of Forest Science* 68 (2): 225-244. - doi: [10.1007/s13595-011-0040-z](https://doi.org/10.1007/s13595-011-0040-z)
- Seidel D, Leuschner C, Müller A, Krause B (2011b). Crown plasticity in mixed forests: quantifying asymmetry as a measure of competition using terrestrial laser scanning. *Forest Ecology and Management* 261 (11): 2123-2132. - doi: [10.1016/j.foreco.2011.03.008](https://doi.org/10.1016/j.foreco.2011.03.008)
- Seidel D, Fleck S, Leuschner C (2012). Analyzing forest canopies with ground-based laser scanning: a comparison with hemispherical photography. *Agricultural and Forest Meteorology* 154-155: 1-8. - doi: [10.1016/j.agrformet.2011.10.006](https://doi.org/10.1016/j.agrformet.2011.10.006)
- Van Laar A, Akça AE (2007). *Forest mensuration* (vol. 13). Springer, Dordrecht, The Netherlands, pp. 399. [online] URL: <http://books.google.com/books?id=ybjdxL2lHOQC>
- Winter M-B, Ammer C, Baier R, Donato DC, Seibold S, Müller J (2015). Multi-taxon alpha diversity following bark beetle disturbance: evaluating multi-decade persistence of a diverse early-seral phase. *Forest Ecology and Management* 338: 32-45. - doi: [10.1016/j.foreco.2014.11.019](https://doi.org/10.1016/j.foreco.2014.11.019)