

## Comparison of drought stress indices in beech forests: a modelling study

Urša Vilhar

Two drought stress indices were applied to managed as well as old-growth beech forests and gaps for the 2001 to 2013 period to aid in the development of an efficient tool for field water supply diagnosis. The relative extractable soil water (REW), which was calculated from the soil water content in the root zone, and the transpiration index (TI), calculated as the ratio between the actual and potential transpiration were used. Both indices were calculated on a daily basis using the water balance model BROOK90, which was fitted and tested using measured data on throughfall and soil water content. A sensitivity analysis apportioned to the input parameters of the drought stress indices was conducted to assess uncertainty. Both drought stress indices showed the greatest drought stress in the years 2009, 2003 and 2011, as also indicated by the Standardized Precipitation Evapotranspiration Index (SPEI) at the nearest meteorological station. However, drought stress intensity and duration differed between the indices and study sites. Greater water supply stress was shown in the forests than the gaps. Furthermore, the agreement among the indices was smaller for gaps compared with forests, which implies that careful index selection is needed when comparing water supply stresses in different stages of forest stand development. Due to the low amount of input data required and the parameters that can be measured with relative ease in the field, REW might be an efficient tool for field water supply diagnosis when analyzing the drought stresses of similar forest types and at unique stages of development. REW satisfactorily indicated drought stress in forests but to a lesser extent in gaps. TI demonstrated more consistent differences in drought stress between forests and gaps and therefore proved to be the appropriate index for a detailed analysis of drought stress variation between different stages of forest stand development. However, due to a greater number of required input data and more demanding parameters, TI appears to be a more complex tool than REW for field water supply diagnosis in forests.

**Keywords:** Relative Extractable Soil Water, Transpiration Index, Standardized Precipitation Evapotranspiration Index, *Fagus sylvatica*, BROOK90 Model, Managed Forest, Old-growth Forest, Canopy Gap

### Introduction

Drought represents a weather-specific phenomenon that affects forests to varying extents and may result in a long-term decrease in forest productivity (Bergès & Balandier 2010, Bertini et al. 2011) or forest decline (Wellpott et al. 2005, Bréda et al. 2006, Briceno-Elizondo et al. 2006, Anderegg et al. 2013b). Over the last 30 years, Europe has been affected by a number of major drought events, most notably in 1976 (Northern and Western Europe), 1989 and 1991 (most of Europe – Mishra & Singh 2010). An extreme drought during the sum-

mer of 2003 led to broad disturbances in European forest ecosystems, which have been reviewed by several authors (Granier et al. 2007, Nikolova et al. 2009, Lebourgeois et al. 2013, Sohn et al. 2013). Repeated droughts are assumed to be responsible for longer fire seasons, increased fire risk (Palahi et al. 2008) and widespread forest mortality (Bréda et al. 2006, Briceno-Elizondo et al. 2006, Anderegg et al. 2013a).

A drought index is usually the primary variable used to assess the effects of drought and to define drought parameters, which include intensity, duration, severity

and spatial extent (Mishra & Singh 2010). Most studies on water supply stress to date have focused on drought stress indices based on climate data (Mishra & Singh 2010, Dogan et al. 2012). Some drought stress indices incorporate topographic and edaphic variability to assess fine-scale soil water availability (Dyer 2009, Piedallu et al. 2013). However, physiological indices, such as those related to soil water deficits (Zahner 1967) or transpiration rates (Zierl 2004, Wellpott et al. 2005), have been shown to be better correlated with radial growth and enable biological interpretations of vegetation responses to climate (Michelot et al. 2012, Tegel et al. 2014).

Currently, European beech (*Fagus sylvatica* L.) represents the most ecologically and economically important forest tree species presently supported by forest management in Slovenia (Kutnar & Kobler 2011). However, the growth and competitive ability of beech may be adversely affected by climate change, particularly drought (Stojanović et al. 2013). Drought stress would be more likely to occur in shal-

□ Slovenian Forestry Institute, Večna pot 2, 1000 Ljubljana (Slovenia)

@ Urša Vilhar ([ursa.vilhar@gozdis.si](mailto:ursa.vilhar@gozdis.si))

Received: Mar 03, 2015 - Accepted: Dec 12, 2015

**Citation:** Vilhar U (2016). Comparison of drought stress indices in beech forests: a modelling study. *iForest* 9: 635-642. - doi: [10.3832/ifor1630-008](https://doi.org/10.3832/ifor1630-008) [online 2016-05-06]

Communicated by: Tamir Klein

low soils such as those in this study. There have been several studies on water supply in canopy gaps (Ritter & Vesterdal 2006, Dalsgaard 2007, Vilhar & Simončič 2012) in which soil water availability influences forest regeneration and establishment (Gray & Spies 1996, Madsen & Hahn 2008, Vilhar et al. 2015). However, drought stress studies are scarce in old-growth forests, and some studies indicate differences in water supply stress between managed and old-growth forests at similar sites (Ritter 2004, Vilhar et al. 2005, Vilhar & Simončič 2012).

The present study compared the performance of two drought stress indices that differed in structure and input data demand in beech forests on shallow soils in Slovenia. The study was carried out in a forest and a gap in an old-growth and a managed forest over a thirteen year period, thereby covering a range of environmental conditions likely to be encountered in Slovenian beech forests. An efficient drought stress index for field water supply diagnosis was sought. We applied (i) the relative extractable soil water (REW), which simulates water deficiency (the availability of soil water in the rooting zone – Bréda et al. 2006, Granier et al. 2007) and (ii) the actual and potential transpiration ratio (*i.e.*, transpiration index, TI – Hammel & Kennel 2001), that considers both atmospheric conditions and the properties and physiological processes of forests and their soils (Zierl 2001, Schwärzel et al. 2009a). Both REW and TI were compared with the Standardized Precipitation Evapotranspiration Index (SPEI – Vicente-Serrano et al. 2010), which is a climatic drought index, using precipitation and reference evapotranspiration at the nearest meteorological station. REW and TI were calculated on a daily basis using measured and modeled data with the water balance model BROOK90 (Federer 2002). We also investigated whether the water supply stress differed between the four sites. This was performed to demonstrate the effect of gap creation as a forest management practice on forest regeneration water supply stress (Vilhar et al. 2015). The comparison of selected drought stress indices should aid in the development of an efficient tool for

field water supply diagnosis (Bergès & Balandier 2010). Such a tool would enable forest managers to develop forest management strategies for adapting to climate change (Kolström et al. 2011).

## Methods

### Site description

The forests under investigation are located in southeastern Slovenia (45° 20' N, 14° 30' E, elevation 860-890 m a.s.l.). They belong to the *Omphalodo-Fagetum* association (Puncer 1980), which is dominated by silver fir (*Abies alba* Mill.) and European beech (*Fagus sylvatica* L.). Norway spruce (*Picea abies* [L.] Karst.), maple (*Acer pseudoplatanus* L.), elm (*Ulmus glabra* Huds.), and lime (*Tiliacordata* Mill.) make up less than 1% of the total stem volume (Vilhar et al. 2015). The bedrock consists of Cretaceous limestone, and the soil depth varies from 10 to 40 cm depending on the highly variable karstic micro-relief. The prevailing soil units were Eutric Cambisols and Rendzic Leptosols (Urbančič et al. 2005). The climate of the region is montane with an annual precipitation of up to 1600 mm. The nearest meteorological station, Kočevje (45° 39' N, 14° 51' E, 467 m a.s.l.), has recorded a long-term (1971-2000) annual average air temperature of 8.0 °C and annual precipitation of 1481 mm (Environmental Agency of the Republic of Slovenia archives). Based on this value and an environmental lapse rate of 6 °C per km of elevation, the long-term annual average temperature at the study area is 5.9°C (Vilhar et al. 2006b).

The specific old-growth and managed forests in this study have similar elevation, aspect and slope. In the old-growth forest, an irregular-shaped gap was formed after windstorms during the winter of 2002-2003 (Tab. 1). In the managed forest, an irregular experimental clear-cut gap was created in the winter of 2000-2001. All of the trees in the experimental gap were harvested and carefully removed by horse skidding.

### Meteorological data and soil hydrological measurements

Meteorological data were collected

above the tree crowns using an automatic weather station (Vantage Pro® wireless, Davis Instruments, Hayward, California, USA). Hourly average values of air temperature and humidity, wind direction and speed, and sums of precipitation were recorded (Vilhar et al. 2006b). Missing air temperature and humidity data were substituted with data from the Kočevje meteorological station, and missing global radiation data were substituted with data from the Iskriba EMEP station (Environmental Agency of the Republic of Slovenia archives) using site-specific regression functions (Vilhar et al. 2006b). In the years 2001 to 2007, monthly or biweekly precipitation (throughfall) was recorded in the forests and gaps at a height of 1.3 m using a series of nine funnel collectors (240 cm<sup>2</sup> each) arranged along a regular grid (spacing: 5×5 m). Incidental precipitation was recorded monthly in the years 2001 to 2007 in an open field close to the study sites using the same type of funnel collectors. For the old-growth gap, which had naturally formed during the winter of 2002-2003, datasets from the years 2003 to 2007 were used.

Using representative soil samples taken from the prevailing soil units, the available water capacity of the mineral soil horizons was calculated for each site from pressure plate measurements of field capacity (moisture content at 0.033 MPa). The permanent wilting point (moisture content at 1.5 MPa) and saturated soil hydraulic conductivity were also determined. The average soil thickness at all sites was 40 cm, and this depth was assumed to be the rooting depth for all sites. The soil water content (SWC) of the 0-40 cm layer was measured monthly at three locations in each plot in 2003 and twice monthly from 2004 to 2007 using time domain reflectometry (TDR – Prenart Equipment, Frederiksberg, Denmark). Double probes were installed vertically, extending through the 40 cm layer, and included the organic layer. Soil-specific calibration curves for vertical TDR probes were used and were obtained using the calibration procedures described in Dirksen (1999). During 2001 and 2002, SWC was determined once per month by taking three replicate volumetric soil samples at

Tab. 1 - General characteristics of the study sites.

Parameter	Managed		Old-growth	
	Forest	Gap	Forest	Gap
Stem volume (m <sup>3</sup> ha <sup>-1</sup> )	255	-	746	-
Basal area (m <sup>2</sup> ha <sup>-1</sup> )	17	-	49	-
Year of creation	-	2000	-	2003
Shape and dimensions	-	Near-circular, 2375 m <sup>2</sup>	-	Irregular, 710 m <sup>2</sup>
Harvesting method	-	Experimental clear-cut	-	Windstorms
Forest floor vegetation - dominant species	<i>Dryopteris filix-mas</i> , <i>Lamium orvala</i> , <i>Polystichum aculeatum</i>		<i>Fagus sylvatica</i>	
Ground vegetation cover (%)	20	6	22	62
Soil depth (cm)	32.2	31.3	32.6	29.9
Stoniness (% vol)	24.4	22.3	29.7	23.1
Soil Texture class	Loam	Loam	Clay loam	Clay loam

depths of 10, 20, and 40 cm in each plot and by measuring the weight loss after oven drying the samples at 105 °C for 24 h. The volumetric moisture contents were converted into depths of water (mm) by multiplying the values by the thickness of the soil layer and correcting for stone content.

#### The BROOK90 simulation

The water balance model BROOK90 Version 4.4g (Federer 1995, Hammel & Kennel 2001, Federer et al. 2003) was applied to forests and gaps to simulate the input parameters needed for the REW and TI calculations. The model calculates daily water fluxes (tree transpiration, evapotranspiration, interception, throughfall, soil evaporation, drainage) and SWC at different depths or for the whole rooting depth. Tree transpiration and soil evaporation are calculated separately using the Shuttleworth-Wallace method (Shuttleworth & Wallace 1985) modified to separate daytime from night time evaporation (Federer 1995). Model parameterisation for the study sites is described in detail by Vilhar & Simončič (2012).

The interception model was fitted by comparing the simulated and the measured monthly throughfall (TF) for the managed and old-growth forest from 2001 to 2003 and tested with the TF from 2004 to 2007. Additional model fitting was performed by comparing the simulated and measured daily SWC of the rooting zone (corrected for stone content) for forests and gaps using datasets from 2001 to 2004 and tested with datasets from 2005 to 2007. For the old-growth gap, which had naturally formed during the winter of 2002-2003, datasets from 2003 to 2007 were used. The extent of TF and SWC fit was evaluated by examining the linear correlation coefficient ( $r$ ), which describes the degree of correspondence between the measured and simulated values; the index of agreement ( $D$  – Thompson 1999), which describes relative error; and the root mean square error (RMSE), which expresses the error between the measured and simulated values.

#### Drought stress indices

##### Relative extractable soil water (REW)

The relative extractable soil water (REW) indicates the availability of soil water in the rooting zone. As a simple drought stress index, the REW may be computed from soil water content in the root zone at any given time, as follows (Bréda et al. 2006, Granier et al. 2007 – eqn. 1):

$$REW = \frac{SWC_{day} - SWC_{min}}{SWC_{max} - SWC_{min}}$$

where  $SWC_{day}$  is the daily soil water content (mm), whereas  $SWC_{min}$  and  $SWC_{max}$  are the minimum and maximum soil water content (mm) for the entire rooting zone during

the period of interest. The REW is between 1.0 (maximum soil water content) and 0 (minimum soil water content). In the present study, the daily REW values for forests and gaps were calculated from the simulated daily SWC during the growing season (i.e., May to October) from 2001 to 2013 (with the exception of the natural gap from 2003 to 2013), using the BROOK90 model. For the REW calculation, the daily  $SWC_{min}$  and  $SWC_{max}$  for the entire modeling period were used. Because a site-specific critical value of the matrix potential was not available for the assessment of the soil water deficits, water supply stress was assumed to occur when the REW dropped below the threshold of 0.4 ( $REW_c$ ), thereby inducing stomatal regulation in forest trees (Granier et al. 1999). Although Lagergren & Lindroth (2002) report on slightly lower REW threshold values for pine and spruce trees in Sweden, the threshold of  $REW < 0.4$  has been widely used in different forest ecosystems (Bréda et al. 2006, Granier et al. 2007, Michelot et al. 2012, Zhou et al. 2013). Additionally, the duration of the water supply stress, i.e., the percentage of growing season days with an REW below 0.4 ( $REW_c$ ) for forests and gaps was calculated to quantify the duration of the water supply stress (Bréda et al. 2006).

##### Transpiration index (TI)

The transpiration index (TI) is a daily ratio between simulated actual and potential transpiration. TI is a more complex drought stress indicator than REW because measurements of transpiration rates in forest ecosystems using the sap flow measurements of single trees and scaling up to forest stands (as in TI) are costly and time consuming (Granier & Loustau 1994, Cermák et al. 2004). Therefore, the mathematical modeling of transpiration rates has become an alternative approach in daily determinations of the seasonal variation of transpiration in forest trees (Granier et al. 2000, Matejka et al. 2007). In the present study, daily TI values for forests and gaps were calculated from simulated actual and potential transpiration during the growing season using the BROOK90 model from 2001 to 2013, with the exception of the natural gap from 2003 to 2013. Potential transpiration in the BROOK90 model is defined as the theoretical transpiration rate that would occur if the soil-plant system was able to satisfy the atmospheric demand and the stomatal regulation was unaffected by drought stress (Federer 1995). Actual transpiration, by contrast, is defined as the actual transpiration rate with stomatal opening adapted to the current drought stress conditions. A TI equal to 1 indicates that the water supply of the forest is optimal, whereas a TI value of less than 1 indicates temporary water deficiency (Wellpott et al. 2005). Hammel & Kennel (2001) defined sites as having frequent water deficiency when the 25<sup>th</sup> percentile of their daily TI values was below 0.95. Schwärzel

et al. (2009a) reported that transpiration in beech stands was significantly reduced when the daily actual transpiration was < 70% of the daily potential transpiration ( $TI < 0.7$ ). Therefore, the percentage of days with a TI below the critical threshold of 0.7 ( $TI_c$ ) for each growing season was calculated to quantify the duration of water supply stress.

##### Standardized Precipitation Evapotranspiration Index (SPEI)

The Standardized Precipitation Evapotranspiration Index (SPEI – Vicente-Serrano et al. 2010) is a climatic drought index that indicates general water supply for a study area. The SPEI may be calculated using monthly (or weekly) precipitation and grass reference evapotranspiration at the nearest meteorological station as follows (eqn. 2):

$$SPEI = P_i - Ref ETP_i$$


where  $P_i$  is monthly (or weekly) precipitation and  $Ref ETP_i$  is grass reference evapotranspiration, which is calculated using the FAO-56 Penman-Monteith equation (Allen et al. 1998, Vicente-Serrano et al. 2010). The SPEI varies between maximum precipitation (when  $Ref ETP_i$  is zero) and reaches negative values when  $Ref ETP_i$  exceeds the precipitation amount, indicating a negative climatic water balance on a monthly basis. In the present study, the monthly SPEI was calculated from  $P$  and  $Ref ETP$  data at the Kočevje meteorological station (Environmental Agency of the Republic of Slovenia archives) and compared with monthly REW and TI for forests and gaps. Additionally, the percentage of days with a negative climatic water balance ( $SPEI < 0.0$ ) for each growing season from 2001 to 2013 was calculated to quantify the duration of the negative climatic water balance in the study area.

##### Statistical analyses

Due to a non-normal distribution, the Spearman's rank correlation coefficient ( $R$ ) was used to evaluate the relationship between the daily REW and TI in forests and gaps. Differences between REW and TI at each site were assessed using the Kruskal-Wallis test. Probability values of  $p < 0.05$  (\*),  $p < 0.01$  (\*\*) and  $p < 0.001$  (\*\*\*) were considered significant. The data analysis was performed using the software package Graphpad® (Graphpad 2014).

##### Results

The results of this study clearly demonstrate significantly greater simulated daily SWC in gaps compared with forests during the growing seasons 2001-2013 ( $H = 5615$ ,  $df = 3$ ,  $N = 9200$ ,  $p < 0.001$  – Fig. 1a). Furthermore, the simulated daily actual transpiration significantly differed between all sites over the course of the study period ( $H = -486.6$ ,  $df = 3$ ,  $N = 9200$ ,  $p < 0.001$ ) with higher average seasonal values in the



forests than the corresponding gaps (Fig. 1b).

#### BROOK90 model fitting and testing

Tab. S1 in Supplementary Material summarises the input parameters used in the simulations. Monthly TF and daily SWC were accurately simulated in the BROOK90 model in respect to both time and magnitude for all sites. The average D was 0.881 and the average RMSE was 11.8 for the SWC model fitting; the average D was 0.754 and the average RMSE was 16.1 for model testing (Tab. S2 and Fig. S1 in Supplementary Material). For TF, the average D for model fitting was 0.823 and average RMSE was 49.4; for model testing, the average D was 0.752 and average RMSE was 78.1 (Tab. S3 and Fig. S2 in Supplementary Material).

#### Drought stress indices

The daily average air temperature in the period from 1971 to 2000 at the Kočevje meteorological station was 8.0 °C with a minimum in January (-1.3 °C) and maximum in July (17.5 °C). The daily average air temperature for the study period from 2001 to 2013 at the Kočevje meteorological station was 9.2 °C, with the highest daily air temperature reported in 2002 ( $9.7 \pm 7.4$  °C) and 2007 ( $9.7 \pm 7.3$  °C) and the lowest in 2005 ( $8.1 \pm 8.5$  °C). The average annual precipita-

tion from 2001 to 2013 was 1408 mm, which was 95% of the long-term (1971-2000) average annual precipitation for the area (1481 mm). The years 2011 and 2003 were substantially drier than normal. There was 1025 mm of precipitation in 2011 and 1152 mm in 2003, accounting for 69% and 78% of the long-term average. When accounting only for growing season, however, precipitation was lowest in 2009 (516 mm), followed by the 2003 (626 mm) and 2013 (634 mm), corresponding to 64%, 77% and 78% of the long-term (1971-2000) average seasonal precipitation for the area (813 mm). Furthermore, the annual SPEI was lowest in 2011 with 267 mm, followed by 358 mm in 2003, corresponding to 37% and 50%, respectively, of the long-term annual SPEI (717 mm from 1971 to 2000). When accounting for only growing seasons, however, a SPEI was lowest in 2009 (-21.5 mm), followed by the 2003 growing season with a SPEI of 2.3 mm. Furthermore, the percentage of days with SPEI < 0.0 was highest in the 2011 growing season (81%), followed by that in the 2001, 2003 and 2009 growing seasons (79%).

The resulting time-courses of the daily REW and TI from 2001 to 2013 followed the opposite pattern of precipitation in the open area and indicated the highest water deficiency in June, July and August at all sites and sufficient water supply in the


autumn months. The drought stress indices showed the greatest drought stress in the growing seasons of 2003, 2009 and 2011; however, the drought stress intensity and duration differed between indices and study sites. According to REW, drought stress was most intense in 2009 at all sites with the exception of the old-growth site with a maximal number of days with  $REW < REW_c$  in 2006 (Fig. 2). Longer drought spells in both forests were also indicated in 2011 and 2008, when the number of days with  $REW < REW_c$  was more than 88% of the days in the growing season. TI indicated the greatest drought stress in the managed forest in 2003 and the managed gap in the 2006 growing season. In the old-growth forest, the greatest drought stress according to TI was in 2011; however, the maximal number of days with  $TI < TI_c$  occurred in 2003 (59% of days in the growing season). For the old-growth gap, however, the greatest drought stress according to TI was in 2013 but the maximal number of days with  $TI < TI_c$  occurred in 2012 (39% of days in the growing season).

Both REW and TI indicated significantly greater drought stress in forests than gaps ( $REW H = 4283.8$ ,  $TI H = 327.5$ ,  $df = 3$ ,  $N = 9200$ ,  $p < 0.001$  – Fig. 3). Furthermore, slightly greater drought stress was shown for the old-growth forest by both indices compared with managed forest. Nevertheless, REW indicated greater and TI minor drought stress in the managed compared with old-growth gap. As a consequence, the number of days with  $TI < TI_c$  indicated a similar difference in drought stress duration between forests and gaps (25% in managed and 23% in old-growth sites), whereas the number of days with  $REW < REW_c$  indicated larger differences in the old-growth (70%) compared with managed sites (55%).

Daily values of both indices were in good agreement between the forests ( $REW R = 0.995$ ,  $TI R = 0.949$ ;  $p < 0.05$ ) as well as between the gaps ( $REW R = 0.898$ ,  $TI R = 0.882$ ;  $p < 0.05$ ). However, there was slightly lower agreement in daily REW or TI between forests and gaps (daily REW between the managed forest and gap:  $R = 0.638$ , daily TI  $R = 0.789$ ; daily REW between the old-growth forest and gap:  $R = 0.679$ , daily TI  $R = 0.657$ ;  $p < 0.05$ ).

When comparing the drought stress intensity at all sites, we found significant differences between the daily REW and TI ( $H = 733.2$ ,  $df = 1$ ,  $N = 5486$ ,  $p < 0.001$ ), as well as between the days below  $REW_c$  and those below  $TI_c$  ( $H = 2039.9$ ,  $df = 1$ ,  $N = 9200$ ,  $p < 0.001$ ). Nevertheless, the agreement between the days below  $REW_c$  and those below  $TI_c$  was slightly higher ( $R = 0.591$ ) than the agreement between the REW and TI ( $R = 0.567$ ,  $p < 0.05$ ), respectively. There was a better agreement between daily REW and TI for the old-growth ( $R = 0.831$ ) and managed forests ( $R = 0.815$ ,  $p < 0.05$ ) than between REW and TI for the gaps ( $R = 0.574$  for the managed gap,  $R = 0.503$  for the old-growth

**Fig. 2** - The number of days with a Standardized Precipitation Evapotranspiration Index below 0.0 (SPEI < 0.0) at the Kočevje meteorological station. (a) The number of days with a relative extractable water content below 0.4 (REW < REW<sub>c</sub>, %) and (b) the number of days with a transpiration index below 0.7 (TI < TI<sub>c</sub>, %) for the managed and old-growth forests and gaps for the 2001 to 2013 growing seasons (i.e., May - October), with the exception of the natural gap from 2003 to 2013.




gap;  $p < 0.05$ ). The agreement between the days below REW<sub>c</sub> and those below TI<sub>c</sub> was better in the managed compared with old-growth sites ( $R = 0.749$  for the managed gap,  $R = 0.509$  for the managed forest,  $R = 0.480$  for the old-growth gap and  $R = 0.461$  for the old-growth forest;  $p < 0.05$ ).

A comparison of the monthly SPEI at all sites with the REW and TI showed statistically significant differences ( $H = 85.0$ ,  $df = 2$ ,  $N = 978$ ,  $p < 0.001$ ). However, the agreement between SPEI and TI was slightly higher ( $R = 0.461$ ) than between the SPEI and REW ( $R = 0.402$ ,  $p < 0.001$ ). Additionally, the difference between the SPEI < 0.0 and the number of days with REW < REW<sub>c</sub> or TI < TI<sub>c</sub> was statistically significant ( $H = 270.8$ ,  $df = 2$ ,  $N = 978$ ,  $p < 0.001$ ). The agreement was higher between the number of days with SPEI < 0.0 and days with TI < TI<sub>c</sub> ( $R = 0.513$ ) than between the number of days with SPEI < 0.0 and the days with REW < REW<sub>c</sub> ( $R = 0.416$ ,  $p < 0.001$  - Fig. 4). Regarding the sites, the highest agreement between SPEI and REW was found in the managed forest ( $R = 0.582$ ) and gap ( $R = 0.546$ ), followed by the old-growth forest ( $R = 0.491$ ) and gap ( $R = 0.483$ ,  $p < 0.001$ ). However, the agreement between the number of days with SPEI < 0.0 and days with REW < REW<sub>c</sub> was highest in the old-growth forest ( $R = 0.607$ ), followed by the managed forest ( $R = 0.580$ ), managed gap ( $R = 0.578$ ) and old-growth gap ( $R = 0.505$ ,


$p < 0.001$ ). The highest agreement between SPEI and TI was found in the old-growth forest ( $R = 0.675$ ) followed by the managed forest ( $R = 0.631$ ,  $p < 0.001$ ),

whereas the agreement was not significant for the gaps. The agreement between the number of days with SPEI < 0.0 and the days with TI < TI<sub>c</sub> was highest in the man-



**Fig. 3** - Daily mean relative extractable soil water (REW) and transpiration index (TI) in the managed and old-growth forests and gaps from 2001 to 2013 growing seasons (i.e., May - October), with the exception of the natural gap from 2003 to 2013. REW and TI equaling 1 indicate no drought stress.

**Fig. 4** - Comparison of monthly percentage of days with a Standardized Precipitation Evapotranspiration Index below 0.0 (SPEI < 0.0, %) at the Kočevje meteorological station and a) the number of days with the relative extractable water content below 0.4 (REW < REW<sub>c</sub>, %) and b) the number of days with the transpiration index below 0.7 (TI < TI<sub>c</sub>, %) in the managed and old-growth forests and gaps from 2001 to 2013 growing seasons (i.e., May - October), with the exception of the natural gap from 2003 to 2013. The line is the regression line for all sites (n = 300).



aged ( $R = 0.715$ ) and old-growth forests ( $R = 0.700$ ), followed by the old-growth ( $R = 0.479$ ) and managed gap ( $R = 0.424$ ,  $p < 0.001$ ).

## Discussion

Both drought stress indices showed the greatest drought stress in the years 2009, 2003 and 2011, as also indicated by the SPEI at the nearest meteorological station. However, drought stress intensity and duration differed between indices and study sites. Both TI and REW showed greater drought stress in forests than gaps in managed as well as old-growth forests. Higher Spearman rank coefficients indicated a better agreement between daily REW and TI for the forests than for the gaps, implying that indices should be carefully selected when comparing water supply stress between different stages of forest stand development.

According to REW, the greatest and longest drought stress in all sites occurred in 2009, followed by the 2001 growing season. TI indicated the greatest and longest drought stress in 2011, followed by the 2003 and 2009 growing seasons. TI was more strongly correlated with the SPEI than REW but only for the forests. The SPEI and TI were not significantly correlated for the gaps.

Both indices (TI and REW) showed a greater water supply stress for the forests than for the gaps, which is in accordance with previous studies (Vilhar et al. 2005, Vilhar & Simončič 2012). Due to lower interception and lower transpiration (Cognard-Plancq et al. 2001, Zirlewagen & Von Wilpert 2001), plants in the gaps benefit from higher precipitation input and increased soil water storage compared with the sur-

rounding forest (Gray et al. 2002, Ritter & Vesterdal 2006, Dalsgaard 2007). However, REW indicated larger differences in drought stress duration between forests and their corresponding gaps (55-70%); similar differences were found via TI (23-25%). The REW calculation is based on the daily, minimum and maximum SWC for the measurement period, which can be derived from modeled data or from continuous SWC profile measurements (Schwärzel et al. 2009b). Such measurements are currently cost-effective (Ferlan & Simončič 2012) and can be numerous installed in larger forest areas. However, a sensitivity analysis revealed that REW is sensitive to both minimum and maximum SWC for the measurement period. Furthermore, REW satisfactorily indicated drought stress in forests but to lesser extent in gaps. Due to the low amount of required input data as well as parameters that can be measured in the field with relative ease, REW may be an efficient tool for field water supply diagnosis (Bergès & Balandier 2010) when analyzing the drought stress of similar forest types and at unique stages of development.

TI demonstrated more consistent differences between forests and gaps in regard to drought stress in different stages of forest stand development. TI is a more complex drought stress indicator than REW. Measurements of actual and potential transpiration rates in trees are costly and time consuming (Granier & Loustau 1994, Cermák et al. 2004). Potential transpiration rates in forest trees are also difficult to measure; however, some studies have reported on the transpiration rates of trees under non-limiting soil water conditions (Cermák et al. 1982, 1993, Cienciala et al.

1994, Granier et al. 2000). Transpiration rates of forests are therefore often simulated using complex water balance models (Granier et al. 2000, Vilhar et al. 2006a, Matejka et al. 2007) with greater input data requirements (Hammel & Kennel 2001). The actual and potential transpiration in the BROOK90 model can be controlled using a number of parameters related to the tree stand, including tree height, leaf water potential, soil matrix potential, inner plant resistance, and others. However, the better performance of TI is associated with a substantial increase in the need for input data and parameters compared with REW. Daily meteorological data as well as parameters describing the site, the canopy, the soil and the water flow through the soil are required. Many of these parameters are rarely measured in the field and are often obtained from the literature. Thus, TI may provide more accurate estimates of drought stress in different forest types and stages of forest stand development; however, due to the greater number of required input data and parameters, TI also appears to be a less efficient tool for field water supply diagnosis in forests.

Further research on site-specific REW and TI thresholds is needed, as well as the integration of drought stress indicators with forest reactions, e.g., crown defoliation (Zierl 2004) or increment losses (Michelot et al. 2012, Tegel et al. 2014) for beech forests in Slovenia.

## Conclusions

Both the relative extractable soil water (REW) and the transpiration index (TI) indicated the greatest drought stress in the years 2009, 2003 and 2011, as also shown by the SPEI at the nearest meteorological station.

Both indices (TI and REW) showed greater water supply stress for the forests than for the gaps in managed as well as old-growth forests. However, REW indicated larger differences in drought stress duration between forests and their corresponding gaps (55-70%), whereas for TI differences between forests and corresponding gaps were smaller (23-25%).

REW may be an efficient tool for field water supply diagnosis when analyzing the drought stress of similar forest types and at unique stages of development due to the low amount of required input data.

TI proved to be an appropriate index for the detailed analysis of drought stress variation in different forest types and stages of forest stand development. However, due to more complex measurements or the mathematical modeling of transpiration rates in forest ecosystems, TI also appears to be a more complex tool for field water supply diagnosis in forests compared to REW.

## Acknowledgements

The study was part of the European 5FW project NAT-MAN (QLRT1-CT99-1349); the

ManFor C.BD project “Managing forests for multiple purposes: carbon, biodiversity and socio-economic wellbeing” (LIFE09 ENV/IT/000078); EUFORINNO “European Forest Research and Innovation” (Reg. Pot No. 315982); several projects within the Programme group “Forest biology, ecology and technology”; and the Programme group Forest biology, ecology and technology (0404-501), which is financed by the Ministry of Education, Science and Sport, Republic of Slovenia.

## References

- Allen RG, Pereira LS, Raes D, Smith M (1998). Crop evapotranspiration. Guidelines for computing crop water requirements. FAO irrigation and drainage paper no. 65, FAO, Rome, Italy, pp. 300. [online] URL: [http://appgeodb.nancy.inra.fr/biljou/pdf/Allen\\_FAO1998.pdf](http://appgeodb.nancy.inra.fr/biljou/pdf/Allen_FAO1998.pdf)
- Anderegg LDL, Anderegg WRL, Abatzoglou J, Hausladen AM, Berry JA (2013a). Drought characteristics role in widespread aspen forest mortality across Colorado, USA. *Global Change Biology* 19: 1526-1537. - doi: [10.1111/gcb.12146](https://doi.org/10.1111/gcb.12146)
- Anderegg LDL, Anderegg WRL, Berry JA (2013b). Not all droughts are created equal: translating meteorological drought into woody plant mortality. *Tree Physiology* 33: 701-712. - doi: [10.1093/treephys/tp044](https://doi.org/10.1093/treephys/tp044)
- Bergès L, Balandier P (2010). Revisiting the use of soil water budget assessment to predict site productivity of sessile oak (*Quercus petraea* Liebl.) in the perspective of climate change. *European Journal of Forest Research* 129: 199-208. - doi: [10.1007/s10342-009-0315-1](https://doi.org/10.1007/s10342-009-0315-1)
- Bertini G, Amoriello T, Fabbio G, Piovosi M (2011). Forest growth and climate change: evidences from the ICP-Forests intensive monitoring in Italy. *iForest* 4: 262-267. - doi: [10.3832/ifor0596-004](https://doi.org/10.3832/ifor0596-004)
- Bréda N, Huc R, Dreyer E, Granier A (2006). Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaptation processes and long-term consequences. *Annals of Forest Science* 63: 625-644. - doi: [10.1051/forest:2006042](https://doi.org/10.1051/forest:2006042)
- Briceno-Elizondo E, Garcia-Gonzalo J, Peltola H, Matala J, Kellomaki S (2006). Sensitivity of growth of Scots pine, Norway spruce and silver birch to climate change and forest management in boreal conditions. *Forest Ecology and Management* 232: 152-167. - doi: [10.1016/j.foreco.2006.05.062](https://doi.org/10.1016/j.foreco.2006.05.062)
- Cermák J, Kucera J, Nadezhkina N (2004). Sap flow measurements with some thermodynamic methods, flow integration within trees and scaling up from sample trees to entire forest stands. *Trees* 18 (5): 529-546. - doi: [10.1007/s00468-004-0339-6](https://doi.org/10.1007/s00468-004-0339-6)
- Cermák J, Matyssek R, Kucera J (1993). Rapid response of large, drought-stressed beech trees to irrigation. *Tree Physiology* 12: 281-290. - doi: [10.1093/treephys/12.3.281](https://doi.org/10.1093/treephys/12.3.281)
- Cermák J, Ulehla J, Kučera J, Penka M (1982). Sap flow rate and transpiration dynamics in the full-grown oak (*Quercus robur* L.) in floodplain forest exposed to seasonal floods as related to potential evapotranspiration and tree dimensions. *Biologia Plantarum (Praha)* 24: 446-460. - doi: [10.1007/BF02880444](https://doi.org/10.1007/BF02880444)
- Cienciala E, Eckersten H, Lindroth A, Hällgren J-E (1994). Simulated and measured water uptake by *Picea abies* under non-limiting soil water conditions. *Agricultural and Forest Meteorology* 71: 147-164. - doi: [10.1016/0168-1923\(94\)90105-8](https://doi.org/10.1016/0168-1923(94)90105-8)
- Cognard-Plancq A-L, Marc V, Didon-Lescot J-F, Norman M (2001). The role of forest cover on streamflow down sub-Mediterranean mountain watersheds: a modelling approach. *Journal of Hydrology* 254: 229-243. - doi: [10.1016/S0022-1694\(01\)00494-2](https://doi.org/10.1016/S0022-1694(01)00494-2)
- Dalsgaard L (2007). Above and belowground gaps-the effects of small canopy opening on throughfall, soil moisture and tree transpiration in Suserup Skov, Denmark. *Ecological Bulletins* 52: 81-102.
- Dirksen C (1999). Soil physics measurements. Catena Verl., Reiskirchen, Germany, pp. 154. [online] URL: <http://www.cabdirect.org/abstracts/20013083835.html>
- Dogan S, Berktaş A, Singh VP (2012). Comparison of multi-monthly rainfall-based drought severity indices, with application to semi-arid Konya closed basin, Turkey. *Journal of Hydrology* 470-471: 255-268. - doi: [10.1016/j.jhydrol.2012.09.003](https://doi.org/10.1016/j.jhydrol.2012.09.003)
- Dyer JM (2009). Assessing topographic patterns in moisture use and stress using a water balance approach. *Landscape Ecology* 24: 391-403. - doi: [10.1007/s10980-008-9316-6](https://doi.org/10.1007/s10980-008-9316-6)
- Federer CA (1995). BROOK90 manual: a simulation model for evaporation, soil water and streamflow, version 3.1. USDA Forest Service, Durham, NH, USA, pp. 40.
- Federer CA (2002). BROOK90: a simulation model for evaporation, soil water, and streamflow. Version 4.4g. Computer freeware, C. Anthony Federer, Falmouth ME 04105, USA. [online] URL: <http://www.ecoshift.net/brook/brook90.htm>. BROOK90
- Federer CA, Vorosmarty C, Fekete B (2003). Sensitivity of annual evaporation to soil and root properties in two models of contrasting complexity. *Journal of Hydrometeorology* 4: 1276-1290. - doi: [10.1175/1525-7541\(2003\)004<1276:SOAETS>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<1276:SOAETS>2.0.CO;2)
- Ferlan M, Simončič P (2012). Robust and cost-effective system for measuring and logging of data on soil water content and soil temperature profile. *Agricultural Sciences* 3: 865-870. - doi: [10.4236/as.2012.36105](https://doi.org/10.4236/as.2012.36105)
- Granier A, Biron P, Lemoine D (2000). Water balance, transpiration and canopy conductance in two beech stands. *Agricultural and Forest Meteorology* 100: 291-308. - doi: [10.1016/S0168-1923\(99\)00151-3](https://doi.org/10.1016/S0168-1923(99)00151-3)
- Granier A, Breda N, Biron P, Villette S (1999). A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands. *Ecological Modelling* 116: 269-283. - doi: [10.1016/S0304-3800\(98\)00205-1](https://doi.org/10.1016/S0304-3800(98)00205-1)
- Granier A, Loustau D (1994). Measuring and modelling the transpiration of a maritime pine canopy from sap-flow data. *Agricultural and Forest Meteorology* 71: 61-81. - doi: [10.1016/0168-1923\(94\)90100-7](https://doi.org/10.1016/0168-1923(94)90100-7)
- Granier A, Reichstein M, Breda N, Janssens IA, Falge E, Ciais P, Gruenwald T, Aubinet M, Berbigier P, Bernhofer C, Buchmann N, Facini O, Grassi G, Heinesch B, Ilvesniemi H, Keronen P, Knohl A, Koestner B, Lagergren F, Lindroth A, Longdoz B, Loustau D, Mateus J, Montagnani L, Nys C, Moors E, Papale D, Peiffer M, Pilegaard K, Pita G, Pumpanen J, Rambal S, Rebmann C, Rodrigues A, Seufert G, Tenhunen J, Vesala T, Wang Q (2007). Evidence for soil water control on carbon and water dynamics in European forests during the extremely dry year: 2003. *Agricultural and Forest Meteorology* 143: 123-145. - doi: [10.1016/j.agrformet.2006.12.004](https://doi.org/10.1016/j.agrformet.2006.12.004)
- Graphpad (2014). GraphPad Prism. Software MacKiev, La Jolla, CA, USA. [online] URL: <http://www.graphpad.com>
- Gray A, Spies T (1996). Gap size, within-gap position and canopy structure effects on conifer seedling establishment. *Journal of Ecology* 84: 635-645. - doi: [10.2307/2261327](https://doi.org/10.2307/2261327)
- Gray A, Spies T, Easter M (2002). Microclimatic and soil moisture responses to gap formation in coastal Douglas-fir forests. *Canadian Journal of Forest Research* 32: 332-343. - doi: [10.1139/x01-200](https://doi.org/10.1139/x01-200)
- Hammel K, Kennel M (2001). Charakterisierung und Analyse der Wasserverfügbarkeit und des Wasserhaushalts von Waldstandorten in Bayern mit dem Simulationsmodell BROOK90 [Characterization and analysis of water availability and the water balance of forest sites in Bavaria with the simulation model BROOK90]. *Forstliche Forschungsberichte München* 185: 135. [in German]
- Kolström M, Lindner M, Vilén T, Maroschek M, Seidl R, Lexer MJ, Netherer S, Kremer A, Delzon S, Barbati A, Marchetti M, Corona P (2011). Reviewing the science and implementation of climate change adaptation measures in European forestry. *Forests* 4: 961-982. - doi: [10.3390/f2040961](https://doi.org/10.3390/f2040961)
- Kutnar L, Kobler A (2011). Prediction of forest vegetation shift due to different climate-change scenarios in Slovenia. *Sumarski list* 135: 113-126.
- Lagergren F, Lindroth A (2002). Transpiration response to soil moisture in pine and spruce trees in Sweden. *Agricultural and Forest Meteorology* 112: 67-85. - doi: [10.1016/S0168-1923\(02\)00060-6](https://doi.org/10.1016/S0168-1923(02)00060-6)
- Lebourgeois F, Gomez N, Pinto P, Mérian P (2013). Mixed stands reduce *Abies alba* tree-ring sensitivity to summer drought in the Vosges mountains, western Europe. *Forest Ecology and Management* 303: 61-71. - doi: [10.1016/j.foreco.2013.04.003](https://doi.org/10.1016/j.foreco.2013.04.003)
- Madsen P, Hahn K (2008). Natural regeneration in a beech-dominated forest managed by close-to-nature principles - a gap cutting based experiment. *Canadian Journal of Forest Research* 38: 1716-1729. - doi: [10.1139/X08-026](https://doi.org/10.1139/X08-026)
- Matejka F, Srelcová K, Hortalová T, Gomoryová E (2007). Modelling transpiration and soil water potential in a spruce primeval forest during dry period. In: Proceedings of the International Scientific Conference on “Bioclimatology and natural hazards”. Polana nad Detvou (Slovakia) 17-20 Sep 2007. Česká bioklimatologická společnost, Praha, Czech Republic, pp. 1-7.
- Michelot A, Bréda N, Damesin C, Dufrêne E (2012). Differing growth responses to climatic variations and soil water deficits of *Fagus sylvatica*, *Quercus petraea* and *Pinus sylvestris* in a temperate forest. *Forest Ecology and Management* 265: 161-171. - doi: [10.1016/j.foreco.2011](https://doi.org/10.1016/j.foreco.2011)

- 10.024
- Mishra AK, Singh VP (2010). A review of drought concepts. *Journal of Hydrology* 391: 202-216. - doi: [10.1016/j.jhydrol.2010.07.012](https://doi.org/10.1016/j.jhydrol.2010.07.012)
- Nikolova PS, Raspe S, Andersen CP, Mainiero R, Blaschke H, Matyssek R, Häberle K-H (2009). Effects of the extreme drought in 2003 on soil respiration in a mixed forest. *European Journal of Forest Research* 128: 87-98. - doi: [10.1007/s10342-008-0218-6](https://doi.org/10.1007/s10342-008-0218-6)
- Palahi M, Mavsar R, Gracia C, Birot Y (2008). Mediterranean forests under focus. *International Forestry Review* 10: 676-688. - doi: [10.1505/ifor.10.4.676](https://doi.org/10.1505/ifor.10.4.676)
- Piedallu C, Gégout J-C, Perez V, Lebourgeois F (2013). Soil water balance performs better than climatic water variables in tree species distribution modelling. *Global Ecology and Biogeography* 22: 470-482. - doi: [10.1111/geb.12012](https://doi.org/10.1111/geb.12012)
- Puncer I (1980). Dinarski jelovo-bukovi gozdovi na Kočevskem [Dinaric silver fir - beech forests in Kočevje region]. *Razprave* 22: 161. [in Czech]
- Ritter E (2004). The effect of gap formation on soil temperature, soil water, and processes in the nitrogen cycle in temperate beech forests of different management intensities. PhD Thesis, The Royal Veterinary and Agricultural University, Copenhagen, Denmark, pp. 232.
- Ritter E, Vesterdal L (2006). Gap formation in Danish beech (*Fagus sylvatica*) forests of low management intensity: soil moisture and nitrate in soil solution. *European Journal of Forest Research* 125: 139-150. - doi: [10.1007/s10342-005-0077-3](https://doi.org/10.1007/s10342-005-0077-3)
- Schwärzel K, Feger K-H, Häntzschel J, Menzer A, Spank U, Clausnitzer F, Köstner B, Bernhofer C (2009a). A novel approach in model-based mapping of soil water conditions at forest sites. *Forest Ecology and Management* 258: 2163-2174. - doi: [10.1016/j.foreco.2009.03.033](https://doi.org/10.1016/j.foreco.2009.03.033)
- Schwärzel K, Menzer A, Clausnitzer F, Spank U, Haentzschel J, Gruenwald T, Koestner B, Bernhofer C, Feger K-H (2009b). Soil water content measurements deliver reliable estimates of water fluxes: A comparative study in a beech and a spruce stand in the Tharandt forest (Saxony, Germany). *Agricultural and Forest Meteorology* 149: 1994-2006. - doi: [10.1016/j.agrformet.2009.07.006](https://doi.org/10.1016/j.agrformet.2009.07.006)
- Shuttleworth WJ, Wallace JS (1985). Evaporation from sparse crops - an energy combination theory. *Quarterly Journal of the Royal Meteorological Society* 111 (469): 839-855. - doi: [10.1002/qj.49711146910](https://doi.org/10.1002/qj.49711146910)
- Sohn JA, Gebhardt T, Ammer C, Bauhus J, Häberle K-H, Matyssek R, Grams TEE (2013). Mitigation of drought by thinning: Short-term and long-term effects on growth and physiological performance of Norway spruce (*Picea abies*). *Forest Ecology and Management* 308: 188-197. - doi: [10.1016/j.foreco.2013.07.048](https://doi.org/10.1016/j.foreco.2013.07.048)
- Stojanović DB, Kržič A, Matović B, Orlović S, Duputic A, Djurdjević V, Galić Z, Stojnić S (2013). Prediction of the European beech (*Fagus sylvatica* L.) xeric limit using a regional climate model: an example from southeast Europe. *Agricultural and Forest Meteorology* 176: 94-103. - doi: [10.1016/j.agrformet.2013.03.009](https://doi.org/10.1016/j.agrformet.2013.03.009)
- Tegel W, Seim A, Hakelberg D, Hoffmann S, Panev M, Westphal T, Büntgen U (2014). A recent growth increase of European beech (*Fagus sylvatica* L.) at its Mediterranean distribution limit contradicts drought stress. *European Journal of Forest Research* 133: 61-71. - doi: [10.1007/s10342-013-0737-7](https://doi.org/10.1007/s10342-013-0737-7)
- Thompson SA (1999). Hydrology for water management. Balkema, Rotterdam, The Netherlands, pp. 476. [online] URL: [http://129.118.24.171/research/txdot\\_0-6070/Literature\\_Archive/thompson1999/thompson1999.pdf](http://129.118.24.171/research/txdot_0-6070/Literature_Archive/thompson1999/thompson1999.pdf)
- Urbančič M, Simončič P, Cater M (2005). Impacts of gaps on humus forms in dinaric silver fir-beech (*Omphalodo-Fagetum*) and soil solution quality. *Mitteilungen Österreichisches Bodenkundliche Gesellschaft* 72: 179-187.
- Vicente-Serrano SM, Beguería S, López-Moreno JI (2010). A multi-scalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index-SPEI. *Journal of Climate* 23: 1696-1718. - doi: [10.1175/2009JCLI2909.1](https://doi.org/10.1175/2009JCLI2909.1)
- Vilhar U, Nadezhkina N, Cermak J, Gasperek J, Urbančič M, Simončič P (2006a). Meritve in modeliranje transpiracije podsajene bukve v smrekovem sestoji na Pohorju [Measuring and modeling of the transpiration of underplanted beech in spruce stand on Pohorje]. In: "Splošne ekološke in gozdnogojitvene osnove za podsadnjo bukve (*Fagus sylvatica* L.) v antropogenih smrekovih sestojih" (Simončič P, Cater M eds). *Silva Slovenica, Gozdarski Inštitut Slovenije, Ljubljana*, vol. 129, pp. 86-103. [in Slovenian]
- Vilhar U, Roenberger D, Simončič P, Diaci J (2015). Variation in irradiance, soil features and regeneration patterns in experimental forest canopy gaps. *Annals of Forest Science* 72: 253-266. - doi: [10.1007/s13595-014-0424-y](https://doi.org/10.1007/s13595-014-0424-y)
- Vilhar U, Simončič P (2012). Water status and drought stress after gap formation in managed and semi-natural silver fir-beech forests. *European Journal of Forest Research* 131: 1381-1397. - doi: [10.1007/s10342-012-0605-x](https://doi.org/10.1007/s10342-012-0605-x)
- Vilhar U, Simončič P, Kajfe-Bogataj L, Katzensteiner K, Diaci J (2006b). Mikroklimatske razmere v vrzelih in sestojih dinarskega jelovobukovega gozda [Microclimate conditions in gaps and mature stands of Dinaric silver fir-beech forests]. *Zbornik gozdarstva in lesarstva* 81: 21-36. [in Slovenian]
- Vilhar U, Starr M, Urbančič M, Smolej I, Simončič P (2005). Gap evapotranspiration and drainage fluxes in a managed and a virgin dinaric silver fir-beech forest in Slovenia: a modelling study. *European Journal of Forest Research* 124: 165-175. - doi: [10.1007/s10342-005-0067-5](https://doi.org/10.1007/s10342-005-0067-5)
- Wellpott A, Imbery F, Schindler D, Mayer H (2005). Simulation of drought for a Scots pine forest (*Pinus sylvestris* L.) in the southern upper Rhine plain. *Meteorologische Zeitschrift* 14: 143-150. - doi: [10.1127/0941-2948/2005/0015](https://doi.org/10.1127/0941-2948/2005/0015)
- Zahner R (1967). Refinement in empirical functions for realistic soil-moisture regimes under forest cover. *Forest Hydrology*. In: *Proceedings of the "National Science Foundation Advanced Science Seminar"*. Pergamon Press, Oxford, Pennsylvania State University, PA, USA, pp. 261-273.
- Zhou J, Zhang Z, Sun G, Fang X, Zha T, McNulty S, Chen J, Jin Y, Noormets A (2013). Response of ecosystem carbon fluxes to drought events in a poplar plantation in Northern China. *Forest Ecology and Management* 300: 33-42. - doi: [10.1016/j.foreco.2013.01.007](https://doi.org/10.1016/j.foreco.2013.01.007)
- Zierl B (2001). A water balance model to simulate drought in forested ecosystems and its applications to the entire forested area in Switzerland. *Journal of Hydrology* 242: 115-136. - doi: [10.1016/S0022-1694\(00\)00387-5](https://doi.org/10.1016/S0022-1694(00)00387-5)
- Zierl B (2004). A simulation study to analyse the relations between crown condition and drought in Switzerland. *Forest Ecology and Management* 188: 25-38. - doi: [10.1016/j.foreco.2003.07.019](https://doi.org/10.1016/j.foreco.2003.07.019)
- Zirlewagen D, Von Wilpert K (2001). Modeling water and ion fluxes in a highly structured, mixed-species stand. *Forest Ecology and Management* 143: 27-37. - doi: [10.1016/S0378-1127\(00\)00522-3](https://doi.org/10.1016/S0378-1127(00)00522-3)

## Supplementary Material

**Tab. S1** - Values for selected parameters used in the BROOK90 simulation.

**Tab. S2** - Linear regression ( $y = a + bx$ ), correlation coefficients ( $r$ ), indices of agreement ( $D$ ), root mean square error (RMSE) and number of measurements ( $n$ ) between the BROOK90 simulated ( $y$ ) and measured ( $x$ ) values for the soil water contents (SWC, mm) of the rooting depth (0 to 40 cm).

**Tab. S3** - Linear regression ( $y = a + bx$ ), correlation coefficients ( $r$ ), indices of agreement ( $D$ ), root mean square error (RMSE) and number of measurements ( $n$ ) between the BROOK90 simulated ( $y$ ) and the measured ( $x$ ) throughfall (TF, mm).

**Fig. S1** - (a) BROOK90 model fitting and b) BROOK90 model testing: simulated and measured values for soil water contents (SWC, mm) of the rooting depth (0 to 40 cm) for the managed forest, gap, and the old-growth forest during 2001-2007 and the old-growth gap during 2003-2007.

**Fig. S2** - Model fitting and testing: simulated and measured values for throughfall TF (mm) for the managed and the old-growth forest during 2001-2007.

**Link:** [Vilhar\\_1630@suppl001.pdf](mailto:Vilhar_1630@suppl001.pdf)