

Influence of soil and topography on defoliation intensity during an extended outbreak of the common pine sawfly (*Diprion pini* L.)

Maiju Kosunen,
Tuula Kantola,
Mike Starr,
Minna Blomqvist,
Mervi Talvitie,
Päivi Lyytikäinen-Saarenmaa

Insect herbivore disturbances are likely to intensify as a consequence of climate change. In Finland, outbreaks of the common pine sawfly (*Diprion pini* L.), which feeds on Scots pine (*Pinus sylvestris* L.) needles, and resulting damage to forests have already increased. Although drivers of sawfly outbreak dynamics have been investigated, the effects of topography and soil fertility have not been fully elucidated. We studied the effect of elevation, slope and soil properties (carbon and nitrogen contents, C/N ratio, pH, texture and horizon thicknesses) on the defoliation intensity of 28 plots (227–531 m²), located in a 34.5 km² forested area in eastern Finland suffering from an extended outbreak of *D. pini*. Plot elevation and slope (relative relief 35 m, maximum elevation 200 m a.s.l.) were derived from a digital elevation model and the soil properties from samples of the humus layer (Of+Oh), (Ah+)E and B horizons of podzol profiles. Defoliation was greater on the more fertile and flatter sites than on less fertile and steeper sites, but independent of elevation. The soil property most strongly correlated to plot mean defoliation was the C/N ratio of the humus layer (Spearman's $\rho = -0.68$). However, logistic modelling showed that the thickness of the (Ah+)E-horizon had the highest classification accuracy in predicting the probability of a plot having moderate to severe (>20%) defoliation. Our study showed that forest damage caused by *D. pini* was related to topography and soil fertility. Taking these factors into account could help in understanding the population dynamics of *D. pini*, in modeling of insect outbreaks and in forest management planning.

Keywords: C/N Balance, Defoliation, Pine Sawfly, Soil, Topography

Introduction

Changes in climate have already been shown to affect the outbreak severity (Haynes et al. 2014) and shift in the range of defoliating insects (Battisti et al. 2006), and thereby alter the functioning of forest ecosystems. In addition to climate, defoliator populations are regulated by a range of biotic and abiotic factors, including natural enemies (Kollberg et al. 2014), topography (Kharuk et al. 2007) and soil (Mayfield et al. 2007). Their impacts and interactions with each other are, however, only partly known.

Forest defoliator damage and performance have been shown to be related to elevation, slope and aspect. Elevation not

only acts as a physical boundary limiting the distribution of defoliators and host trees but it also affects local climatic conditions (temperature, precipitation, wind speed and radiation) and influences insect physiology and performance (Hodkinson 2005). Natural enemy abundance and host-species nutritional value and distribution can also vary with elevation and have an impact on defoliator tree damage and insect performance (Niemelä et al. 1987, McMillin et al. 1996, Hengxiao et al. 1999, Kharuk et al. 2007). Steep, especially south to west-facing slopes, are generally drier, warmer and have more nutrient deficient conditions than other slopes, and can result in more stressed host trees and possi-

bly more favorable local climates and habitats for defoliators (Morse & Kulman 1986, Kharuk et al. 2007). However, damage by defoliators has also been found to be higher on flat areas compared to steeper slopes (Kharuk et al. 2009) and sapsucking insects have been shown to favor north-northwest-facing slopes (Kantola et al. 2014).

Soil fertility affects the nutritional value of the foliage and production of defense compounds by host trees and therefore has an important indirect effect on defoliator performance and related tree damage. Several studies have suggested that trees growing on poorer sites are more prone to defoliator-caused damage compared to trees growing on more fertile sites (Larsson & Tenow 1984, Nevalainen et al. 2015). Greater defoliator performance and higher damage intensity have been associated with coarser soil texture (Cobb et al. 1997, Mayfield et al. 2007) and thin soil A-horizons (Hood et al. 1988), both indicating poor soil fertility (Brady & Weil 2014). Fertilization experiments have shown that an increase in soil fertility has a positive effect on defoliator performance (Mopper & Whitham 1992). However, other fertilization experiments have shown a negative effect (Larsson & Tenow 1984) or no effect (Björkman et al. 1991) on defoliator performance. Tree foliage nitrogen (N) and car-

□ Department of Forest Sciences, University of Helsinki, P.O. Box 27 (Latokartanonkaari 7), FI-00014 Helsinki (Finland)

@ Maiju Kosunen (maiju.kosunen@helsinki.fi)

Received: Mar 26, 2016 - Accepted: Oct 18, 2016

Citation: Kosunen M, Kantola T, Starr M, Blomqvist M, Talvitie M, Lyytikäinen-Saarenmaa P (2017). Influence of soil and topography on defoliation intensity during an extended outbreak of the common pine sawfly (*Diprion pini* L.). *iForest* 10: 164-171. - doi: [10.3832/ifor2069-009](https://doi.org/10.3832/ifor2069-009) [online 2016-11-19]

Communicated by: Massimo Faccoli

bon (C) contents relate to both their nutritional value (Lyytikäinen 1994, Giertych et al. 2007), which can benefit insect performance, and to the production of defense compounds such as C-based monoterpenes (Barre et al. 2003) and phenolics (Giertych et al. 2007) and N-based alkaloids (Barbosa & Krischik 1987), which can repel insects. For example, Scots pine (*Pinus sylvestris* L.) needle N contents increase with soil N contents (Björkman et al. 1991, Raitio 1999), which have also been shown to affect the contents of secondary defense compounds (Björkman et al. 1991, Kainulainen et al. 1996). In addition, predators and parasitoids of defoliators have been shown to be affected by site fertility (Hanski & Parviainen 1985, Herz & Heitland 2005).

The common pine sawfly (*Diprion pini* L., Hymenoptera: Diprionidae) is distributed throughout Europe and northern Africa (Geri 1988). Until the 1990's the species caused only small-scale defoliation of *P. sylvestris* in Finland (De Somviele et al. 2007). In 1997, *D. pini* populations reached outbreak densities in western Finland and the outbreak started to spread eastwards (De Somviele et al. 2007). By 2001, this largest insect-driven forest damage in the Finnish forest health records, had spread over an area of ca. 500 000 ha (Lyytikäinen-Saarenmaa & Tomppo 2002). Consumption of all of the needle year-classes by *D. pini* reduces the photosynthesizing leaf area, leading to decreased tree growth (Långström et al. 2001, Lyytikäinen-Saarenmaa & Tomppo 2002) and mortality if the needle consumption continues over many consecutive years (Långström et al. 2001). In Finland the economic losses of reduced tree growth and mortality related to *D. pini* has been estimated to be 93-530 US\$ ha⁻¹ depending on defoliation intensity (Lyytikäinen-Saarenmaa & Tomppo 2002). Overall, sawfly outbreaks seem to be more frequent and severe on monoculture, even-aged and plantation stands, compared to forests with several tree species, age classes and layers (Geri 1988, McMillin & Wagner 1993). In Scandinavia, the performance of pine sawflies and the frequency, intensity and spatial extent of damage has been related to temperature (Kollberg et al. 2015) and host plant quality, especially needle carbohydrate and terpene concentrations (Larsson et al. 1986, Lyytikäinen 1994, Kollberg et al. 2015). In addition, site and stand characteristics, such as site type and stand age (Larsson & Tenow 1984, De Somviele et al. 2007, Nevalainen et al. 2015), and natural enemies (Hanski & Parviainen 1985, Kollberg et al. 2014) have been observed to modify sawfly performance and affect the level of tree damage. However, the effects of topography and soil properties on sawfly outbreaks have received little attention. A better understanding of the relationship between pine sawfly outbreak intensity and these abiotic factors would improve our ability to iden-

tify sites prone to pine sawfly outbreaks.

In this study we investigated the spatial variation in the severity of defoliation by *D. pini* in managed *P. sylvestris* stands in eastern Finland. The overall aim was to determine the extent to which the level of defoliation was related to topographical features (elevation and slope) and various chemical and physical properties of the surface organic layer and underlying mineral soil. Our specific objectives were: (1) to determine the variation in *P. sylvestris* defoliation by *D. pini*; (2) to determine a number of topographic parameters and soil physical and chemical properties; and (3) to investigate the relationships between these abiotic factors and defoliation. According to our knowledge, this is the first study investigating the effects of topographic characteristics and soil properties on *D. pini* related defoliation of *P. sylvestris* in Fennoscandian boreal forest ecosystems.

Material and methods

Research area

The study was carried out in Palokangas, a forested area of 34.5 km² located in the municipality of Ilomantsi (62° 52' N, 30° 56' E), eastern Finland. *D. pini* has caused severe damage to the managed *P. sylvestris* dominated forests in the area since 1999. Although having a typical eruptive population dynamic, the local *D. pini* population has in some parts of the area adopted a chronic outbreak pattern with high population levels. At its most extensive range (beginning of 2000's), the outbreak area covered ca. 10 000 ha (Kantola et al. 2010, Talvitie et al. 2011). Changes in the *D. pini* population and resulting tree damage intensity within the study area have been monitored annually since 2000 (De Somviele et al. 2007).

The soils in the study area are mainly podzols and developed in moraine and glaciofluvial deposits. The surface organic layer is a mor type. According to the Cajanderian site type classification (Mikola 1982), which describes site productivity and soil fertility, the area is dominated by rather poor *Vaccinium* type (VT) and poor *Calluna* type (CT) site types with the ground vegetation consisting mainly of lingonberry (*Vaccinium vitis-idaea* L.), heather (*Calluna vulgaris* L. Hull), blueberry (*Vaccinium myrtillus* L.), mosses, and occasional lichens. The mean annual temperature and precipitation for the study area in the period 1981-2010 were 2-3 °C and 650-700 mm, respectively (Pirinen et al. 2012).

Sample plots and stand inventory

The study was carried out on 28 circular sampling plots located throughout the study area. The radius of the plots varied between 8.5 and 13 m, resulting in an average of 24 trees on each plot. As part of a larger project (Kantola et al. 2010, 2013, Talvitie et al. 2011), eleven of the plots were

established in 2002 and 17 in 2007. The locations of the plots were chosen subjectively to ensure that the range in defoliation intensity levels within the study area was covered. The plots were established in even-aged *P. sylvestris* dominated stands, with occasional Norway spruce (*Picea abies* L.), silver birch (*Betula pendula* Roth), aspen (*Populus tremula* L.) and juniper (*Juniperus communis* L.) also present. Most (26) of the plots were located on poor CT site types and two plots on somewhat more fertile VT site types. Tree and plot-wise characteristics were inventoried in May and June 2010. The coordinates of the plot centers were determined using a Trimble Pro XH-GPS® device (Sunnyvale, CA, USA). All the trees on each plot were located by measuring distance and azimuth from the plot center and classified into hierarchy classes: dominant, co-dominant, or suppressed. The diameter at breast height (dbh) of all trees having a dbh > 5.6 cm was measured and tree height of every third tree and that of the median tree in each hierarchy class measured (7-8 sampling trees per plot).

The level of defoliation of all the *P. sylvestris* trees was visually estimated into percentage defoliation classes according to Eichhorn et al. (2010), except we used 10% defoliation classes rather than 5% classes. Accordingly, the defoliation intensity of each tree was assessed by comparing the foliage density of the upper two thirds of the tree crown (i.e., that part of the crown not influenced by shading from other trees) to an imaginary healthy tree growing on the same site type and at the same canopy cover layer. Trees with 100% defoliation level corresponded to dead trees and trees with 0% defoliation had no symptoms of insect defoliation. The plot-wise mean defoliation intensity was calculated as the average defoliation level of the dominant and co-dominant trees on the plot. Suppressed trees were excluded because their defoliation may be due to shading and competition for nutrients. For logistic reasons the defoliation assessment of 17 of the plots was carried out in autumn 2009 and that of the remaining 11 plots in spring 2010. However, as the *D. pini* larvae feed in July-September, the defoliation situation of autumn 2009 corresponds to the situation in spring 2010. The complete history of tree defoliation is unknown as annual surveys of the tree defoliation on the study plots have only been carried out since plot establishment. The outbreak in the area started in 1999 and a peak in defoliation was reached in 2005, after which some of the plots have recovered to a healthier state. However, the pattern of defoliation among the plots has not greatly changed, although the overall level of defoliation has decreased after 2005.

Using the plot-wise mean defoliation levels, each plot was classified as having either "mild" (< 20% defoliation intensity) or "moderate to severe" defoliation (> 20%

defoliation intensity). The 20% defoliation intensity value was used because defoliation exceeding 20% in the Nordic countries is generally considered harmful for tree growth (Strand 1997, Lyytikäinen-Saarenmaa 1999, Lyytikäinen-Saarenmaa & Tomppo 2002). Of the plots, 21 (75%) were classified as having *mild* defoliation and 7 (25%) as having *moderate to severe* defoliation in 2010.

Topography

Elevation and slope were derived from a high pulse density airborne scanning LiDAR (Light Detection and Ranging) data set. The LiDAR point cloud was acquired from the Palokangas area in July 2008 using an ALS50-II SNO58[®] laser scanner (Leica Geosystem AG, Heerbrugg, Switzerland) as described by Kantola et al. (2010). With the standard TerraScan approach (Axelsson 2000), the data set was classified into ground or non-ground points and a high resolution (1 m) digital elevation model (DEM) was generated using classified ground points.

Elevation above the sea level (m a.s.l.) and slope were computed from the DEM for each plot using ArcMap[®] (ArcGis v. 9.3, ESRI, Redlands, CA). Plot mean values for elevation (m a.s.l.) were calculated from the point data. The slope layer was delineated according to the perimeter of each plot and the mean values of the pixels within the plot were used. Topographical data for two of the western-most sample plots could not be used due to a lack of information on the elevation on the plots.

Soil sampling and laboratory analysis

In June 2010, soil cores (n = 140) were collected from the center of each plot and at 5.3 m distance from the center in each cardinal direction in order to get a set of samples representative of each plot. A steel auger (diameter 58 mm) was used to take intact soil cores to a depth of ca. 0.5 m, unless limited by the bedrock. Each soil core was extracted from the auger, the surface litter layer (Ol) removed, and the thicknesses of the humus layer (Of+Oh), (Ah+)E, B and C-horizons measured to a precision of 0.1 cm. The C-horizon was missing from some cores as a consequence of the thin soil. Each soil core was then cut into the above horizons and each placed in a separate plastic bag and kept at 5 °C until analyzed in the laboratory. The plot-wise mean thickness of each horizon was calculated from the five samples from each plot with the exception of three plots where horizon thickness could not be measured due to the unclear interpretation of the horizons and their boundaries. As the (Ah+)E and C-horizon samples were not always sufficient for physical and chemical analysis, only the humus layer and B-horizon samples of each core were analyzed.

The soil samples were dried at 50 °C for 24 hours after which they were weighed. The humus layer samples were milled,

passed through a 2 mm sieve and composited by plot. For the B-horizon samples, a 25 ml subsample from each dried sample was taken and composited by plot. The composited samples were passed through a 2 mm sieve and the weight of >2 mm and <2 mm fractions recorded. The <2 mm fraction was retained for analysis.

The particle size distribution of the <2 mm B-horizon soil was determined using laser fractionation (Coulter LS230[®], Beckman Coulter Inc., Brea, CA, USA). The following texture classification was used: coarse sand (0.6-2 mm), medium sand (0.2-0.6 mm), fine sand (0.06-0.2 mm), coarse silt (0.02-0.06 mm), medium silt (0.006-0.02 mm), fine silt (0.002-0.006 mm), and clay (< 0.002 mm). Total organic C and total N contents were determined from the milled humus layer samples and <2 mm B-horizon samples using a VarioMax CN device (Elementar Analysensysteme GmbH, Hanau, Germany). The soil C/N ratio was calculated from the C and the N contents. Soil pH was determined from a suspension of the milled humus layer samples and 0.1 M calcium chloride solution (1:2), using a glass electrode.

Statistical analysis

Spearman's rank correlation coefficients were used to describe the relationships between plot topographical features, soil properties and plot-wise mean defoliation intensity. The non-parametric Mann-Whitney U-test was used to test if there were significant differences in each predictor variable (topography and soil variables) between the *mild* defoliation (< 20% defoliation) and *moderate to severe* defoliation (> 20% defoliation) classes. Logistic regression was used to investigate the power of the topographical and soil variables to predict the probability of a plot belonging to the *moderate to severe* defoliation class. Predictor variables that were strongly correlated with each other were not added to the same model. This ensured the models were simple (only up to three predictors) and avoided possible over-fitting. The logistic regression was performed for several

different combinations of predictor variables. Evaluation of the models was based on predictor variable p-values, overall classification accuracies, and Cohen's Kappa-values (Landis & Koch 1977). The R-statistical computing environment (R Core Team 2013) and the "irr" package (Gamer et al. 2012) were used to make the analysis.

Results

Characteristics of the study trees

Plot mean dbh and mean height were similar in the *mild* defoliation and *moderate to severe* defoliation classes (20.5 vs. 19.2 m, and 20.6 vs. 18.9 m, respectively – Tab. 1). The mean number of stems per hectare and mean basal area were higher in the *mild* defoliation class (565 stems ha⁻¹, 18 m² ha⁻¹) than in the *moderate to severe* defoliation class (446 stems ha⁻¹, 15 m² ha⁻¹). Mean defoliation in the *mild* defoliation class was 10% and 54% in the *moderate to severe* defoliation class (Tab. 1).

Topographical characteristics and soil properties

The study area was generally rather flat (Fig. 1). The highest elevations were found along the central southwest-northeast axis running through the study area and sloping towards the northwest and southeast. Plot-mean elevations varied between 165 and 200 m a.s.l. and plot-mean slope between 1 and 14° (Tab. 2).

The thickness of the humus layer varied between 2.9 and 6.4 cm and the thickness of (Ah+)E-horizon between 2.1 and 7.4 cm (Tab. 2). The soils were rather acidic, with a mean pH of 4.03. Humus layer C and N contents varied between 19.5 and 45.7% and between 0.50 and 1.43%, respectively, resulting in C/N ratios varying between 27 and 56 (Tab. 2). The C and N contents of the B-horizon were an order of magnitude lower than humus layer values, and C/N ratios varied between 16 and 30. Soil texture varied considerably among the plots. The plot mean proportion of coarse silt and finer (< 0.06 mm) varied between 10.7 and 44.2% and the proportions of medium silt

Tab. 1 - Forest characteristics (mean, median, minimum, maximum, and standard deviation - STD) of the study plots by defoliation class *mild* (<20% defoliation) and *moderate to severe* (>20% defoliation). (dbh): diameter at breast height; (h): tree height; (N): number of stems per hectare; (BA): basal area.

Defoliation	Statistics	dbh (cm)	h (m)	N (stems ha ⁻¹)	BA (m ² ha ⁻¹)	Defoliation (%)
<i>Mild</i> (n=21)	Mean	20.5	19.2	565	18	10
	Median	20.5	19.3	526	19	10
	Min	16.5	16.4	320	10	1
	Max	24.0	21.6	1057	29	19
	STD	2.1	1.4	184	5	6
<i>Moderate to severe</i> (n=7)	Mean	20.6	18.9	446	15	54
	Median	20.8	18.3	464	16	51
	Min	16.4	15.4	283	9	26
	Max	24.1	23.1	575	21	77
	STD	2.5	2.4	90	4	22




Fig. 1 - Digital elevation model of the study area and location of the plots (black dots).

Tab. 2 - Topographic characteristics and soil properties of the study plots (mean \pm standard deviation, minimum and maximum; n=26 for topographical characteristics, n=25 for soil horizon thickness, n=28 for other soil properties).

Variable	Mean \pm STD	Min	Max
Elevation (m a.s.l.)	192 \pm 10	165	200
Slope ($^{\circ}$)	4 \pm 3	1	14
Humus layer thickness (cm)	4.5 \pm 0.9	2.9	6.4
(Ah+)E-horizon thickness (cm)	4.0 \pm 1.4	2.1	7.4
Humus layer pH	4.03 \pm 0.29	3.54	4.78
Humus layer C (%)	34.6 \pm 6.2	19.5	45.7
Humus layer N (%)	1.01 \pm 0.22	0.50	1.43
Humus layer C/N ratio	35 \pm 5	27	56
B-horizon C (%)	1.7 \pm 0.6	0.7	3.4
B-horizon N (%)	0.08 \pm 0.03	0.04	0.15
B-horizon C/N ratio	21 \pm 4	16	30
B-horizon medium sand & finer (< 0.6 mm) (%)	86.8 \pm 7.6	61.3	97.4
B-horizon fine sand & finer (< 0.2 mm) (%)	54.1 \pm 10.1	34	77.5
B-horizon coarse silt & finer (< 0.06 mm) (%)	23.5 \pm 8.6	10.7	44.2
B-horizon medium silt & finer (< 0.02 mm) (%)	8.4 \pm 2.6	4.6	13.6
B-horizon fine silt & finer (< 0.006 mm) (%)	3.2 \pm 0.9	1.9	4.8
B-horizon clay (< 0.002 mm) (%)	1.4 \pm 0.4	0.8	2.4

and finer (< 0.02 mm) between 4.6 and 13.6 %. The proportion of clay (< 0.002 mm) was rather low on all of the plots (Tab. 2).

The thickness of (Ah+)E-horizon had a significant negative correlation with slope, but otherwise the soil properties were not significantly correlated with the topographical variables (Tab. 3). Several of the variables describing site fertility, such as the C/N ratio and proportion of fine soil particles (< 0.02 mm), were significantly correlated with each other (Tab. 3).

Relationships between topography, soil and defoliation

Slope had a significant negative correlation with defoliation but there was no significant difference in slope between the mild and moderate to severe defoliation classes (Tab. 4, Fig. 2). The C/N ratio had a significant negative correlation with plot mean defoliation but it was clearly

Tab. 3 - Spearman's correlation coefficients (ρ) between plot topographical characteristics and soil properties (n=26 for topographical characteristics, n=25 for soil horizon thickness, n=28 for other soil properties). See Tab. 2 for explanation of variable abbreviations. (*): $P < 0.05$; (**): $P < 0.01$; (***): $P < 0.001$.

Parameters	Elevation (m a.s.l.)	Slope ($^{\circ}$)	Humus layer (cm)	(Ah+)E-hor. (cm)	Humus layer pH	Humus layer C (%)	Humus layer N (%)	Humus layer C/N	B-hor. C (%)	B-hor. N (%)	B-hor. C/N	B-hor. < 0.02 mm (%)
Elevation	1	0.06	-0.35	-0.37	0.02	-0.04	-0.14	0.08	0.33	0.30	0.11	-0.02
Slope	-	1	0.05	-0.49*	-0.11	-0.09	-0.15	0.28	-0.06	-0.18	0.06	-0.09
Humus layer	-	-	1	0.36	-0.29	0.45*	0.28	0.30	0.07	0.08	-0.07	0.25
(Ah+)E-hor.	-	-	-	1	0.11	0.35	0.46*	-0.23	-0.51**	-0.22	-0.41*	0.31
Humus layer pH	-	-	-	-	1	-0.08	0.13	-0.41*	-0.13	-0.12	0.08	0.07
Humus layer C	-	-	-	-	-	1	0.85***	0.00	0.18	0.35	-0.29	0.33
Humus layer N	-	-	-	-	-	-	1	-0.45*	0.20	0.52**	-0.49**	0.55**
Humus layer C/N	-	-	-	-	-	-	-	1	-0.13	-0.45*	0.38*	-0.43*
B-hor. C	-	-	-	-	-	-	-	-	1	0.80***	0.16	0.25
B-hor. N	-	-	-	-	-	-	-	-	-	1	-0.38*	0.53**
B-hor. C/N	-	-	-	-	-	-	-	-	-	-	1	-0.54**
B-hor. < 0.02 mm	-	-	-	-	-	-	-	-	-	-	-	1

stronger for the humus layer. The thickness of (Ah+)E-horizon, the N content of humus layer and the proportion of medium silt and finer (< 0.02 mm) in B-horizon were all significantly and positively correlated with plot mean defoliation (Tab. 4, Fig. 2). Concerning differences between the *mild* and *moderate* to *severe* defoliation classes, the thickness of (Ah+)E-horizon, the N content and C/N ratio of the humus layer, and the C/N ratio and proportion of medium silt and finer (< 0.02 mm) in the B-horizon all showed a significant difference (Tab. 4).

Logistic regression models

The best model in predicting the probability of a plot having *moderate* to *severe* defoliation (>20%) included the thickness of the (Ah+)E-horizon as a single predictor and had a classification accuracy of 88 %

Tab. 4 - Spearman's correlation coefficients (ρ) between various plot topographical characteristics and soil properties and mean defoliation (%), and Mann-Whitney U test values and significance of difference between plots classified as having *mild* (<20 % foliage loss) and plots having *moderate* to *severe* (>20 % foliage loss) defoliation (n=26 for topographical characteristics, n = 25 for soil horizon thickness, n = 28 for other soil properties). See Tab. 2 for explanation of variable abbreviations. (*): P < 0.05; (**): P < 0.01; (***) : P < 0.001.

Variable	Spearman's ρ	Mann-Whitney U test
Elevation (m a.s.l.)	-0.26	42.0
Slope (°)	-0.52**	33.0
Humus layer (cm)	0.10	39.0
(Ah+)E-hor. (cm)	0.62***	19.0*
Humus layer pH	0.05	43.5
Humus layer C (%)	0.14	52.0
Humus layer N (%)	0.45*	36.5*
Humus layer C/N	-0.68***	35.0*
B-hor. C (%)	-0.05	70.0
B-hor. N (%)	0.25	49.0
B-hor. C/N	-0.39*	35.5*
B-hor. <0.02 mm (%)	0.44*	36.0*




Fig. 2 - Scatter plots of plot-wise mean defoliation against significantly (p<0.05) correlated (Spearman ρ , Tab. 4) topographical characteristics and soil properties. See Tab. 2 for explanation of variable abbreviations. Lines are best fitting (highest R²) regression models.

Tab. 5 - Logistic regression models for predicting probability of a plot having *moderate* to *severe* defoliation (>20 % foliage loss) ($\log[p/(1-p)] = \beta_0 + \beta_1x_1 + \dots + \beta_kx_k$, where p is the probability (0, 1), β_0 the intercept, and β_1, \dots, β_k the regression coefficients for predictor variables x_1, \dots, x_k). (SE): standard error; (P-value): significance of predicting variable being associated with the probability of plot having *moderate* to *severe* defoliation; (Accuracy): classification accuracy (%); (Kappa): Kappa statistic (0.41-0.60 = moderate agreement, 0.61-0.80 = substantial agreement). See Tab. 2 for explanation of variable abbreviations. (*): P < 0.05; (**): P < 0.01; (***) : P < 0.001.

Model	Variable	Coefficient	SE	P-value	Accuracy	Kappa
Model 1(n=25)	Intercept	-6.779	2.507	0.007**	-	-
	(Ah+)E-hor. (cm)	1.277	0.531	0.016*	88	0.65
Model 2 (n=28)	Intercept	55.662	24.648	0.024*	-	-
	Humus layer C/N	-0.721	0.346	0.037*	-	-
	Humus layer pH	-7.910	3.617	0.029*	86	0.58
Model 3 (n=26)	Intercept	11.055	5.155	0.032*	-	-
	B-hor. C/N	-0.437	0.213	0.040*	-	-
	Slope (°)	-0.960	0.536	0.073	85	0.57
Model 4 (n=28)	Intercept	9.732	10.014	0.331	-	-
	Humus layer N (%)	6.773	3.128	0.030*	-	-
	Humus layer pH	-4.429	2.694	0.100	82	0.50

and a substantial Kappa-value of 0.65 (Model 1 – Tab. 5). The models using the C/N ratio and pH of the humus layer (Model 2), C/N ratio of the B-horizon and slope (Model 3), and N content and pH of humus layer (Model 4) all had moderate Kappa-values (Tab. 5).

Discussion

Topography effects

We did not find a significant impact of elevation on *D. pini* defoliation in our study. Other studies concerning insect outbreaks and elevation imply that patterns of insect performance along elevational gradients differ between insect species. Only a few studies regarding pine sawfly performance in relation to elevation have been conducted. However, Niemelä et al. (1987) observed the damage caused by the European pine sawfly (*Neodiprion sertifer* Geoffr.) on *P. sylvestris* in northern Finland to be more severe at upper summits compared to the lower ones and was due to differences in foliar nutrient contents. Similarly, larval and cocoon masses of a pine sawfly, *Neodiprion xiangyunicus* (Xiao & Huang), feeding on Yunnan pine (*Pinus yunnanensis* Franch.) were found to increase with elevation in south-western China (Hengxiao et al. 1999). McMillin et al. (1996) observed that pine defoliation by the sawfly *Neodiprion autumnalis* (Smith) was present only at elevations of 2410–2440 m a.s.l. and by the sawfly *Neodiprion xiangyunicus* (Xiao and Hung) at 1850–2050 m a.s.l. in southern USA and southern China. Kantola et al. (2014) observed higher hemlock woolly adelgid (*Adelges tsugae* Annand) induced tree mortality on higher elevations within a southern Appalachian forest landscape in western North Carolina, USA. In contrast, Kharuk et al. (2007) found that the highest conifer mortality caused by the Siberian silkmoth (*Dendrolimus superans sibiricus* Tschetw.) in central Siberia was at lower (200 m) elevations than at higher elevations (300 m). While *N. sertifer* has been found to adapt to environmental conditions at various elevations and latitudes in Europe (Pschorn-Walcher 1991), the low range in elevation among the plots most probably accounts for the lack of an elevation effect in our study.

Defoliator susceptible sites have been found on both steep and flat slopes. Morse & Kulman (1986) found steeper and south-facing slopes to have a higher probability for the white spruce (*Picea glauca* [Moench] Voss) defoliation by the yellow-headed spruce sawfly (*Pikonema alaskensis* Rohver) in Minnesota, USA, and Kharuk et al. (2007) found the most intensive Siberian silkmoth damage associated with slopes of 5–20° compared to slopes of less than 5°. In a later study, Kharuk et al. (2009) found that while the initial outbreak of Siberian silkmoth in the Eastern Sayan Mountains occurred on slopes of less than 5°, it later spread to less favorable sites on

steeper slopes. While we found no significant difference in slope between the *mild* and *moderate to severe* defoliation classes, plot mean *D. pini* defoliation was negatively correlated to slope. However, the *D. pini* outbreak has been on-going for more than 10 years; if it were to follow the pattern observed by Kharuk et al. (2009), one would have expected the intensity of damage to have shifted to steeper slopes. However, the higher defoliation on flatter slopes we found could be related to a more favorable forest structure or microclimate for *D. pini* during critical parts of its life cycle, such as oviposition early in the season or larval feeding in early fall (Jactel et al. 2009).

Soil fertility effects

Although having a rather limited range in fertility, we found that plot-mean defoliation levels were significantly greater on the more fertile plots (i.e., humus layers having higher N contents and lower C/N ratios and B-horizons having higher contents of medium silt and finer). The results from other studies concerned with the effects of soil fertility on insect damage to trees and insect performance have been partly contradictory to what we found. Larsson & Tenow (1984) found an outbreak of *N. sertifer* in southern Sweden to be concentrated to unfertile soils and Nevalainen et al. (2015) showed that *N. sertifer* and *D. pini* damage in Finland was more common on sub-xeric heath forests and poorer sites (i.e., sites with relatively poor soil fertility) than on more fertile sites. Studies by Hood et al. (1988), Cobb et al. (1997) and Mayfield et al. (2007) also found that soil properties indicating lower fertility were associated with greater tree damage. Mayfield et al. (2007) found a negative correlation between pine false webworm sawfly (*Acantholyda erythrocephala* L.) population densities and A-horizon silt content and Cobb et al. (1997) showed that tree mortality caused by the pinyon tip moth (*Dioryctria albivittella* Hulst) was more severe on cinder (volcanic) soils with lower silt and clay contents, compared to sandy soils with higher silt and clay contents. Hood et al. (1988) observed that a thin A-horizon and lower calcium (Ca) contents in soil led to higher rates of pine tip moth species (*Rhyacionia* spp.) infestation.

The N contents of the humus layer and B-horizon in our study were within the range (0.3–1.5% for humus layer and 0.01–0.14% for 0–30 cm mineral soil layer, n=47) reported for similar site types in southern Finland (Tamminen 1991). Those studies that found the incidence of insect outbreak and degree of tree damage decreased with site fertility were for sites covering a greater range or having a higher level of site fertility, as compared with our study. The apparent contradictory relationship between insect outbreak and damage and site fertility between these studies and ours may thus be because the relationship is curvilinear,

with one side of the relationship showing a positive relationship and the other side showing a negative relationship, or because the relationship within groups of sites having similar levels of fertility differs from that across all the groups. In sites of poor fertility, the N nutritional value of the trees may be below the optimal for *D. pini*. However, the outbreak of *D. pini* in our study had been chronic for several years and therefore might exhibit a special kind of behavior.

Using fertilization experiments to mimic the effects of soil fertility on defoliator-caused damage and defoliator performance have given contradictory results. Mopper & Whitham (1992) found that cocoons of pinyon sawfly (*Neodiprion edulicollis* Ross) had higher masses in sites with NPK-fertilized and watered pinyon pine (*Pinus edulis* Englm.) compared to those having no additions. In contrast, Larsson & Tenow (1984) observed that ammonium fertilized plots suffered from milder damage and had a lower number of *N. sertifer* cocoons than non-fertilized plots, while Björkman et al. (1991) did not find ammonium-nitrate fertilization to have any effect on the performance of *N. sertifer* larvae. Differences in the level of tree damage and insect performance in relation to soil fertility are probably related to several factors, including differences in host species chemistry and insect ecology, outbreak patterns, scale of the study and range in site fertility and/or predation and parasitism.

N, soluble carbohydrates glucose and fructose, and phosphorus are important in the diet of pine sawflies and therefore also important determinants of pine sawfly performance (Lyytikäinen 1994, Giertych et al. 2007). C-based defensive compounds, such as resin acids (Larsson et al. 1986), phenolics (Giertych et al. 2007) and 3-carene monoterpene (Barre et al. 2003) have been found to have negative effects on pine sawfly performance. Bryant et al. (1983) suggested that plants growing on nutrient-poor sites have a surplus of C from which they can allocate a part to C-based defense compounds. Bryant et al. (1983) also proposed that increased N concentrations in soil would enhance the allocation of C to tree growth and decrease the amount of C-based defense compounds and carbohydrates and increase the contents of N and N-based defense compounds. The N content of the humus layer in Finnish forests has been shown to significantly correlate with the N content of *P. sylvestris* current year needles (Raitio 1999). Increased soil N content via fertilization has also been found to increase *P. sylvestris* needle N and resin acid concentrations (Björkman et al. 1991) and the number of resin glands (Kainulainen et al. 1996), but to decrease contents of monoterpenes (Kainulainen et al. 1996). *P. sylvestris* foliar contents of N and P decrease (Helmisaari 1992) and contents of phenolic compounds increase (Giertych et al. 2007) with needle age. Current and

previous year needles, both of which are consumed by *D. pini*, may also respond differently to N additions. For example, Kainulainen et al. (1996) found that while phenolic contents in current-year *P. sylvestris* needles decreased after ammonium-nitrate fertilization, there was no effect on the previous year needles. In our study, the trees on plots with lower soil C/N ratios possibly were able to allocate more C to growth and less to defense, and had needles with higher amounts of soluble N and lower amounts of phenolics and monoterpenes, resulting in more attractive nutrition for *D. pini*. The *D. pini* population in our study area has also been at gradation and post-gradation level since 1999. In such a chronic outbreak situation, continuous tree defoliation may increase the N content and decrease C-based defensive chemical content of the host-plant material due to changed C allocation patterns, making them more favorable for defoliators (McMillin & Wagner 1997). This partly self-maintaining mechanism might have affected the outbreak pattern of *D. pini* and the relationship between defoliation and soil N in our study area.

Since insect herbivores can influence soil characteristics, e.g., by nutrient input into the soil via frass (Frost & Hunter 2007, Kaukonen et al. 2013), the higher N content of humus layer on our study plots with higher mean defoliation intensity could have resulted from *D. pini* rather than vice versa. However, we observed that N content of the humus layer had a significant positive correlation with the proportion of fine soil particles of the B-horizon. Soil N contents often increase with increasing proportion of fine soil particles (Brady & Weil 2014) and the particle size distribution of mineral soil is not affected by *D. pini* defoliation. Therefore N contents of the humus layer in our study indicate the long-term nutrient status of the site rather than the consequences of *D. pini* defoliation.

Conclusions

Although the fertility was generally poor and topographical variation low across our study plots, we found *D. pini* defoliation of *P. sylvestris* trees to be greater on the more fertile and flatter sites than on the less fertile and steeper sites, but to be independent of elevation. Plot mean defoliation was most strongly correlated with the C/N ratio of the humus layer. However, the thickness of (Ah+E) horizon was the most successful soil property determining the probability of having moderate to severe defoliation (>20% foliage loss). The positive relationship between *D. pini* defoliation and site fertility we found contrasts with the findings of other studies in which insect performance and damage has been found to be negatively correlated to site fertility. While this may be related to differences in the scale of site fertility, the outbreak of *D. pini* in our study had been chronic for several years and therefore

might exhibit a special kind of behavior.

The effect of defoliators on C and nutrient cycling in forest ecosystems is not well known however, and the interaction between soil, topography and defoliators, especially in relation to climate change, needs further study. In particular, the relationship between soil properties, needle nutrient and defense chemical contents, and defoliator performance needs to be clarified. We have shown that soil and terrain conditions contribute to pine sawfly population dynamics and forest damage. Ways should be sort to take these factors into account in forest management planning, selection of silvicultural practices and in the modeling of insect outbreaks in forests.

Acknowledgements

We wish to thank Antero Pasanen and Jari Tahvanainen from Tornator Ltd., who enabled this study in Palokangas area in Ilomantsi. Bert De Somviele and Anna-Maija Kokkonen who kindly assisted in establishing the sampling plots in 2002, and Saara Ilvesniemi in 2007. We would also like to thank Marjut Wallner for help in the laboratory. In addition, we wish to thank the three anonymous reviewers of this paper for valuable comments. This study was made possible by grants from the Finnish Cultural Foundation – South Karelia Regional Fund, *Societas pro Fauna et Flora Fennica*, Niemi Foundation, Maj and Tor Nessling Foundation, and the Finnish Academy project “Centre of Excellence in Laser Scanning Research” (CoE-LaSR, decision number 272195).

References

- Axelsson P (2000). DEM generation from laser scanner data using adaptive TIN models. In: Proceedings of the “XIX ISPRS Congress, Commission I-VII” (Fritsch D, Molenaar M eds). Amsterdam (The Netherlands), 16-23 July 2000. pp. 110-117. [online] URL: http://www.isprs.org/proceedings/XXXIII/congress/part4/111_XXXIII-part4.pdf
- Barbosa P, Krischik VA (1987). Influence of alkaloids on feeding preference of Eastern deciduous forest trees by the gypsy moth *Lymantria dispar*. *The American Naturalist* 130: 53-69. - doi: [10.1086/284697](https://doi.org/10.1086/284697)
- Barre F, Goussard F, Geri C (2003). Variation in the suitability of *Pinus sylvestris* to feeding by two defoliators, *Diprion pini* (Hym., Diprionidae) and *Graellsia isabellae galliaegloria* (Lep., Attacidae). *Journal of Applied Entomology* 127: 249-257. - doi: [10.1046/j.1439-0418.2003.00655.x](https://doi.org/10.1046/j.1439-0418.2003.00655.x)
- Battisti A, Stastny M, Buffo E, Larsson S (2006). A rapid altitudinal range expansion in the pine processionary moth produced by the 2003 climatic anomaly. *Global Change Biology* 12: 662-671. - doi: [10.1111/j.1365-2486.2006.01124.x](https://doi.org/10.1111/j.1365-2486.2006.01124.x)
- Björkman C, Larsson S, Gref R (1991). Effects of nitrogen fertilization on pine needle chemistry and sawfly performance. *Oecologia* 86: 202-209. - doi: [10.1007/BF00317532](https://doi.org/10.1007/BF00317532)
- Brady NC, Weil RR (2014). Elements of the nature and properties of soils. Prentice Hall,

New Jersey, USA, pp. 1046.

- Bryant JP, Chapin FS, Klein DR (1983). Carbon/nutrient balance of boreal plants in relation to vertebrate herbivory. *Oikos* 40: 357-368. - doi: [10.2307/3544308](https://doi.org/10.2307/3544308)
- Cobb NS, Mopper S, Gehring CA, Caouette M, Christensen KM, Whitham TG (1997). Increased moth herbivory associated with environmental stress of pinyon pine at local and regional levels. *Oecologia* 109: 389-397. - doi: [10.1007/s004420050098](https://doi.org/10.1007/s004420050098)
- Eichhorn J, Roskams P, Ferretti M, Mues V, Szepesi A, Durrant D (2010). Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests Part IV. Visual Assessment of Crown Condition and Damaging Agents. UNECE ICP Forests Programme Coordinating Centre, Hamburg, Germany, pp. 49. [online] URL: http://www.icp-forests.org/pdf/FINAL_Crown.pdf
- Frost CJ, Hunter MD (2007). Recycling of nitrogen in herbivore feces: plant recovery, herbivore assimilation, soil retention, and leaching losses. *Oecologia* 151: 42-53. - doi: [10.1007/s00442-006-0579-9](https://doi.org/10.1007/s00442-006-0579-9)
- Gamer M, Lemon J, Fellows I, Singh P (2012). irr: Various coefficients of interrater reliability and agreement. R package version 0.84. [online] URL: <http://CRAN.R-project.org/package=irr>
- Geri C (1988). The pine sawfly in central France. In: “Dynamics of forest insect populations” (Berryman AA ed). Plenum Press, New York, USA, pp. 377-405. - doi: [10.1007/978-1-4899-0789-9_19](https://doi.org/10.1007/978-1-4899-0789-9_19)
- Giertych MJ, Karolewski P, Grzebyta J, Oleksyn J (2007). Feeding behavior and performance of *Neodiprion sertifer* larvae reared on *Pinus sylvestris* needles. *Forest Ecology and Management* 242: 700-707. - doi: [10.1016/j.foreco.2007.02.005](https://doi.org/10.1016/j.foreco.2007.02.005)
- Hanski I, Parviainen P (1985). Cocoon predation by small mammals, and pine sawfly population dynamics. *Oikos* 45: 125-136. - doi: [10.2307/3565230](https://doi.org/10.2307/3565230)
- Haynes KJ, Allstadt AJ, Klimetzek D (2014). Forest defoliator outbreaks under climate change: effects on the frequency and severity of outbreaks of five pine insect pests. *Global Change Biology* 20: 2004-2018. - doi: [10.1111/gcb.12506](https://doi.org/10.1111/gcb.12506)
- Helmisaari H-S (1992). Spatial and age-related variation in nutrient concentrations of *Pinus sylvestris* needles. *Silva Fennica* 26: 145-153. - doi: [10.14214/sf.a15643](https://doi.org/10.14214/sf.a15643)
- Hengxiao G, McMillin JD, Wagner MR, Zhou J, Zhou Z, Xu X (1999). Altitudinal variation in foliar chemistry and anatomy of yunnan pine, *Pinus yunnanensis*, and pine sawfly (Hym., Diprionidae) performance. *Journal of Applied Entomology* 123: 465-471. - doi: [10.1046/j.1439-0418.1999.00395.x](https://doi.org/10.1046/j.1439-0418.1999.00395.x)
- Herz A, Heitland W (2005). Species diversity and niche separation of cocoon parasitoids in different forest types with endemic populations of their host, the common pine sawfly *Diprion pini* (Hymenoptera: Diprionidae). *European Journal of Entomology* 102: 217-224. - doi: [10.14411/eje.2005.034](https://doi.org/10.14411/eje.2005.034)
- Hodkinson ID (2005). Terrestrial insects along elevation gradients: species and community responses to altitude. *Biological Reviews* 80:

- 489-513. - doi: [10.1017/S1464793105006767](https://doi.org/10.1017/S1464793105006767)
- Hood WM, Hedden RL, Berisford CW (1988). Hazard rating forest sites for pine tip moth, *Rhyacionia* spp., in the Upper Piedmont Plateau. *Forest Science* 34: 1083-1093. [online] URL: <http://www.ingentaconnect.com/content/saf/ffs/1988/00000034/00000004/art00021>
- Jactel H, Nicoll BC, Branco M, Gonzalez-Olabarria JR, Grodzki W, Långström B, Moreira F, Netherer S, Orazio C, Piou D, Santos H, Schelhaas MJ, Tojic K, Vodde F (2009). The influences of forest stand management on biotic and abiotic risks of damage. *Annals of Forest Science* 66: 701-719. - doi: [10.1051/forest/2009054](https://doi.org/10.1051/forest/2009054)
- Kainulainen P, Holopainen J, Palomäki V, Holopainen T (1996). Effects of nitrogen fertilization on secondary chemistry and ectomycorrhizal state of Scots pine seedlings and on growth of grey pine aphid. *Journal of Chemical Ecology* 22: 617-636. - doi: [10.1007/BF02033574](https://doi.org/10.1007/BF02033574)
- Kantola T, Vastaranta M, Yu X, Lyytikäinen-Saarenmaa P, Holopainen M, Talvitie M, Kaasalainen S, Solberg S, Hyyppä J (2010). Classification of defoliated trees using tree-level airborne laser scanning data combined with aerial images. *Remote Sensing* 2: 2665-2679. - doi: [10.3390/rs2122665](https://doi.org/10.3390/rs2122665)
- Kantola T, Vastaranta M, Lyytikäinen-Saarenmaa P, Holopainen M, Kankare V, Talvitie M, Hyyppä J (2013). Classification of needle loss of individual Scots pine trees by means of airborne laser scanning. *Forests* 4: 386-403. - doi: [10.3390/f4020386](https://doi.org/10.3390/f4020386)
- Kantola T, Lyytikäinen-Saarenmaa P, Coulson RN, Strauch S, Tchakerian MD, Holopainen M, Saarenmaa H, Streett DA (2014). Spatial distribution of hemlock woolly adelgid induced hemlock mortality in the southern Appalachians. *Open Journal of Forestry* 4: 492-506. - doi: [10.4236/ojf.2014.45053](https://doi.org/10.4236/ojf.2014.45053)
- Kaukonen M, Ruotsalainen AL, Wäli PR, Männistö MK, Setälä H, Saravesi K, Huusko K, Markkola A (2013). Moth herbivory enhances resource turnover in subarctic mountain birch forests? *Ecology* 94: 267-272. - doi: [10.1890/12-0917.1](https://doi.org/10.1890/12-0917.1)
- Kharuk VI, Ranson KJ, Fedotova EV (2007). Spatial pattern of Siberian silkmoth outbreak and taiga mortality. *Scandinavian Journal of Forest Research* 22: 531-536. - doi: [10.1080/02827580701763656](https://doi.org/10.1080/02827580701763656)
- Kharuk VI, Ranson KJ, Im ST (2009). Siberian silkmoth outbreak pattern analysis based on SPOT VEGETATION data. *International Journal of Remote Sensing* 30: 2377-2388. - doi: [10.1080/01431160802549419](https://doi.org/10.1080/01431160802549419)
- Kollberg I, Bylund H, Huitu O, Björkman C (2014). Regulation of forest defoliating insects through small mammal predation: reconsidering the mechanisms. *Oecologia* 176: 975-983. - doi: [10.1007/s00442-014-3080-x](https://doi.org/10.1007/s00442-014-3080-x)
- Kollberg I, Bylund H, Jonsson T, Schmidt A, Gershenson J, Björkman C (2015). Temperature affects insect outbreak risk through tritrophic interactions mediated by plant secondary compounds. *Ecosphere* 6: 1-17. - doi: [10.1890/ES15-00021.1](https://doi.org/10.1890/ES15-00021.1)
- Landis JR, Koch GG (1977). The measurement of observer agreement for categorical data. *Biometrics* 33: 159-174. - doi: [10.2307/2529310](https://doi.org/10.2307/2529310)
- Larsson S, Björkman C, Gref R (1986). Responses of *Neodiprion sertifer* (Hym., Diprionidae) larvae to variation in needle resin acid concentration in Scots pine. *Oecologia* 70: 77-84. - doi: [10.1007/BF00377113](https://doi.org/10.1007/BF00377113)
- Larsson S, Tenow O (1984). Areal distribution of a *Neodiprion sertifer* (Hym., Diprionidae) outbreak on Scots pine as related to stand condition. *Holarctic Ecology* 7: 81-90. [online] URL: <http://www.jstor.org/stable/3682253>
- Lyytikäinen P (1994). Effects of natural and artificial defoliations on sawfly performance and foliar chemistry of Scots pine saplings. *Annales Zoologici Fennici* 31: 307-318. [online] URL: <http://www.jstor.org/stable/23735535>
- Lyytikäinen-Saarenmaa P (1999). Growth responses of Scots pine (Pinaceae) to artificial and sawfly (Hymenoptera: Diprionidae) defoliation. *The Canadian Entomologist* 131: 455-463. - doi: [10.4039/Ent131455-4](https://doi.org/10.4039/Ent131455-4)
- Lyytikäinen-Saarenmaa P, Tomppo E (2002). Impact of sawfly defoliation on growth of Scots pine *Pinus sylvestris* (Pinaceae) and associated economic losses. *Bulletin of Entomological Research* 92: 137-140. - doi: [10.1079/BER2002154](https://doi.org/10.1079/BER2002154)
- Långström B, Annala E, Hellqvist C, Varama M, Niemelä P (2001). Tree mortality, needle biomass recovery and growth losses in Scots pine following defoliation by *Diprion pini* (L.) and subsequent attack by *Tomicus piniperda* (L.). *Scandinavian Journal of Forest Research* 16: 342-353. - doi: [10.1080/02827580118325](https://doi.org/10.1080/02827580118325)
- Mayfield AE, Allen DC, Briggs RD (2007). Site and stand conditions associated with pine false webworm populations and damage in mature eastern white pine plantations. *Northern Journal of Applied Forestry* 24: 168-176. [online] URL: <http://www.ingentaconnect.com/content/saf/njaf/2007/00000024/00000003/art00003>
- McMillin JD, Hengxiao G, Wagner MR, Long X (1996). Spatial distribution patterns of pine sawflies (Hymenoptera: Diprionidae) in Arizona, US and Sichuan, PR of China. *Forest Ecology and Management* 86: 151-161. - doi: [10.1016/S0378-1127\(96\)03793-0](https://doi.org/10.1016/S0378-1127(96)03793-0)
- McMillin JD, Wagner MR (1993). Influence of stand characteristics and site quality on sawfly population dynamics. In: "Sawfly life history adaptation to woody plants" (Wagner MR, Raffa KF eds). Academic Press, London, UK, pp. 333-361.
- McMillin JD, Wagner MR (1997). Chronic defoliation impacts pine sawfly (Hymenoptera: Diprionidae) performance and host plant quality. *Oikos* 79: 357-362. - doi: [10.2307/3546019](https://doi.org/10.2307/3546019)
- Mikola P (1982). Application of vegetation science to forestry in Finland. In: "Handbook of vegetation Science, Part 12" (Jahn G ed). Dr W. Junk Publishers, The Hague/Boston/London, pp. 199-224.
- Mopper S, Whitham TG (1992). The plant sex paradox: effects on pinyon sawfly sex ratios and fecundity. *Ecology* 73: 515-525. - doi: [10.2307/1940757](https://doi.org/10.2307/1940757)
- Morse BW, Kulman HM (1986). A method of hazard-rating white spruce plantations for yellow-headed spruce sawfly defoliation. *Northern Journal of Applied Forestry* 3: 104-105. [online] URL: <http://www.ingentaconnect.com/content/saf/njaf/1986/00000003/00000003/art00009>
- Nevalainen S, Sirkkiä S, Peltoniemi M, Neuvonen S (2015). Vulnerability to pine sawfly damage decreases with site fertility but the opposite is true with *Scleroderris* canker damage; results from Finnish ICP Forests and NFI data. *Annals of Forest Science* 72: 909-917. - doi: [10.1007/s13595-014-0435-8](https://doi.org/10.1007/s13595-014-0435-8)
- Niemelä P, Rousi M, Saarenmaa H (1987). Topographical delimitation of *Neodiprion sertifer* (Hym., Diprionidae) outbreaks on Scots pine in relation to needle quality. *Journal of Applied Entomology* 103: 84-91. - doi: [10.1111/j.1439-0418.1987.tb00962.x](https://doi.org/10.1111/j.1439-0418.1987.tb00962.x)
- Pirinen P, Simola H, Aalto J, Kaukoranta J-P, Karlsson P, Ruuhela R (2012). Climatological statistics of Finland 1981-2010. Reports 2012-1, Finnish Meteorological Institute, Helsinki, Finland, pp 96. [online] URL: <http://hdl.handle.net/10138/35880>
- Pschorn-Walcher H (1991). Development and diapause of different European provenances of the pine sawfly *Neodiprion sertifer* (Geoff.) (Hym., Diprionidae) under identical outdoor conditions. *Journal Applied Entomology* 112: 382-388. - doi: [10.1111/j.1439-0418.1991.tb01071.x](https://doi.org/10.1111/j.1439-0418.1991.tb01071.x)
- R Core Team (2013). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [online] URL: <http://www.R-project.org/>
- Raitio H (1999). Needle chemistry. In: "Forest condition monitoring in Finland, National Report 1998" (Raitio H, Kilponen T eds). The Finnish Forest Research Institute, Research Papers 743: 51-69. [online] URL: <http://urn.fi/URN:ISBN:951-40-1692-0>
- De Somville B, Lyytikäinen-Saarenmaa P, Niemelä P (2007). Stand edge effects on distribution and condition of diprionid sawflies. *Agricultural and Forest Entomology* 9: 17-30. - doi: [10.1111/j.1461-9563.2006.00313.x](https://doi.org/10.1111/j.1461-9563.2006.00313.x)
- Strand L (1997). Monitoring the environmental quality of Nordic forests. *NORD 1997: 14*, Nordic Council of Ministers, Copenhagen, Denmark, pp. 77. [online] URL: <http://books.google.com/books?id=zLD0ZDHAgUC>
- Talvitie M, Kantola T, Holopainen M, Lyytikäinen-Saarenmaa P (2011). Adaptive cluster sampling in inventorying forest damage by the common pine sawfly (*Diprion pini*). *Journal of Forest Planning* 16: 1-7. [online] URL: <http://ci.nii.ac.jp/naid/110009357764/en>
- Tamminen P (1991). Expression of soil nutrient status and regional variation in soil fertility of forested sites in southern Finland. *Folia Forestalia* 777. The Finnish Forest Research Institute, Helsinki, Finland, pp. 40. [in Finnish with English summary]