

Comparison of soil CO₂ emissions between short-rotation coppice poplar stands and arable lands

Chiara Ferré,
Roberto Comolli

Bioenergy crops are considered to have the potential for climate mitigation and socio-economic benefits owing to their capacity to sequester C and partially replace the consumption of fossil fuels. In this study, the effects on soil respiration of a recent conversion of arable land to high (H) and very high (VH) density short-rotation coppices (SRCs), as well as of agronomic treatments (fertilization with or without irrigation) and lane/row location, were investigated in an alluvial area in Italy. A survey of soil variability was carried out by collecting soil cores (0-60 cm depth) at 67 points to characterize surface and subsurface spatial distributions of pH, organic carbon, nitrogen and carbonates and identify comparable points for monitoring soil respiration. Soil CO₂ emissions were monitored over the period April 2010-November 2011 at 27 locations covering the whole study site. The influence of land use (H-SRC, VH-SRC, corn and alfalfa) or treatments on soil respiration was evaluated considering both factors as fixed effects in a linear mixed model. Our results showed that (i) the high variability of soil properties even at small spatial scale has to be considered when selecting points for monitoring soil respiration in the field; (ii) the cumulative soil respiration over the study period at the VH-SRC was lower ($1299 \pm 30 \text{ g C m}^{-2}$) than in croplands ($1600 \pm 145 \text{ g C m}^{-2}$) and higher along the rows than in the lanes; (iii) no significant differences in soil respiration were found between the H-SRC and corn field; (iv) two years after VH-SRC establishment, agronomic treatments did not appear to influence soil respiration; (v) land-use change affected the vertical soil organic carbon distribution and soil surface temperature, as reflected in soil respiration differences.

Keywords: Land-use Change, Soil Spatial Variability, Soil Respiration, Short-rotation Coppice, Cropland, Mixed Model

Introduction

The identification of alternatives to fossil fuels has become a worldwide priority. Various biofuels based on food crops such as corn and soybean, or produced from lignocellulosic compounds of non-food crops, such as grasses and woody crops, play an ecologically and economically fundamental role as renewable energy supplies. In Italy, energy crops have mainly been developed under the regional financial support of EC Regulation 1257/1999. In particular, a short-rotation coppice (SRC) culture, which is a high-density plantation of fast-growing trees (such as poplars or willows) with cutting periods from 2 to 8 years, is considered to have a high potential for climate

mitigation and socio-economic benefits owing to its capacity to sequester C and partially replace the consumption of fossil fuels (Smith 2004).

Bioenergy cultivation can be considered carbon neutral since as much carbon is absorbed during the growth of the SRC plantation as it is released upon conversion to energy. Nevertheless, bioenergy cultivation may represent a CO₂ source in relation to its associated land-use changes, acting as a CO₂ source mainly during the first years of establishment, as a result of the cultivation practices that accelerate the short-term decomposition of organic matter in the soil (Grigal & Berguson 1998, Arevalo et al. 2011).

Many studies have been carried out on poplar short-rotation coppice biomass production and its quality, energy balance and management intensity (Kauter et al. 2003, Nassi et al. 2010), by comparing the advantages of poplar SRC with other bioenergy crops (Karp & Shield 2008) or by evaluating the conversion of agricultural lands from food production to poplar SRC (Boehmel et al. 2008, Gasol et al. 2010). However, there are few studies on SOC dynamics and CO₂ fluxes under SRC (Vande Walle et al. 2007, Verlinden et al. 2013b, Zenone et al. 2015).

To assess the impact of land-use changes on soil properties and CO₂ fluxes, space-for-time substitution studies on a paired-plot, for which the assumption of an identical starting point has to be verified, are mainly used (Poeplau & Don 2013, Ferré et al. 2014). Since soil spatial variability is reflected in the differences in soil respiration rates (Ferré et al. 2015), it follows that an appropriate sampling design should vary according to site conditions and depending on study objectives.

This study is part of the SUSFACE project (<http://www.gruppo-panacea.it/biomasse/progetti/susface>), which aims to investigate the conversion of agricultural lands from food production (sugar beet) to SRC

□ Department of Earth and Environmental Sciences, Milano Bicocca University, Milan (Italy)

@ Chiara Ferré (chiara.ferre@unimib.it)

Received: Sep 02, 2017 - Accepted: Dec 14, 2017

Citation: Ferré C, Comolli R (2018). Comparison of soil CO₂ emissions between short-rotation coppice poplar stands and arable lands. *iForest* 11: 199-205. - doi: [10.3832/ifor2621-010](https://doi.org/10.3832/ifor2621-010) [online 2018-03-01]

Communicated by: Werther Guidi Nissim

poplar plantations for bioenergy. In the context of the SUSCACE project, the objectives of this study were to: (i) identify comparable soil respiration monitoring points based on a soil spatial variability survey; (ii) assess the effects of recent conversion of arable land to SRC on soil respiration; and (iii) investigate of the influence of agronomic treatments and of lane/row position on soil CO₂ efflux in a 2-year-old very high density SRC.

Materials and methods

Study site

The experimental site (Fig. 1) is located close to the Po River within the CREA (Italian Council for Agricultural Research and Economics) experimental farm at Casale Monferrato in Piedmont, Italy (45° 08' 07.77" N; 08° 30' 42.16" E). The climate is temperate continental, as monitored by the long-term meteorological station at the experimental site, with a yearly average rainfall of 715 mm and mean air temperature of 12.5 °C.

Soils did not exhibit a high pedogenetic degree since they developed on recent alluvial deposits. According to the WRB classification (IUSS Working Group 2015), they are Calcaric Regosols (Epiloamic, Endoarenic) and sometimes Endoskeletal (see Fig. S1 in Supplementary Materials). The soil surface texture (USDA class) is mostly sandy loam, while the soil texture becomes sandy once deeper in the soil. In agronomical terms (Peverill & La Sparrow Reuter 1999), soils are low in SOM, moderately calcareous and slightly alkaline.

The study area (1.52 ha) had been managed in a uniform manner for at least 60 years before the land was converted. In the autumn of 2006, the total area of former cropland was alfalfa cultivated (*Medicago sativa*). Beginning in the spring of 2009, the area was subdivided into 4 fields (each field was approximately 0.38 ha):

one alfalfa, one corn, and two poplar SRC (*Populus × canadensis* Mönch) fields. One of the two poplar SRC fields was a high density trial (H-SRC – 1,111 trees ha⁻¹; 3 × 3 m spacing), and one was a very high density trial (VH-SRC – 8,333 trees ha⁻¹; 3 × 0.4 m spacing), with cutting periods of 5 and 2 years, respectively.

For the VH-SRC, four different cultivation treatments were tested: control (CTR), fertilization (F), irrigation (I), and both fertilization and irrigation (FI). The SRC models and treatments were tested using a split-plot design with three replications (each plot covered approximately 50 m²).

The entire area of the experiment, except for a section that remained alfalfa cultivated, was ploughed to a depth of 35 cm and harrowed before field subdivision and planting or seeding. In the VH-SRC trial, post-planting residual herbicides (pendimethalin and metolachlor) were applied for 30-40 days after planting to prevent germination of annual weeds. During the planting year, disc harrowing was carried out in the lanes twice in the VH-SRC and only once in the H-SRC. During the cultivation period, the harrowing was carried out once per year for both the stands.

Every year, three chemical pesticide treatments against insects (*Chrysomela populi*, *Cryptorhynchus lapathi* and *Hyphantria cunea*) were needed for both poplar trials; fertilization was carried out once a year in the early spring using 60 kg of nitrogen per hectare. A drip irrigation system was utilized, and during the summer, a quantity of water corresponding to the estimated evapotranspiration (by a Class A pan) was applied three times a week. Alfalfa and corn were sprinkler irrigated twice every summer (350 m³ ha⁻¹). Only corn was ploughed and fertilized (210 kg ha⁻¹ of nitrogen) once every spring.

Multiple harvestings per year (4-5 times) were applied for alfalfa and once per year for corn, while in the VH-SRC plantation,

harvesting was carried out at the end of the first year.

Survey of soil spatial variability

A survey of soil spatial variability was carried out to characterize the surface and subsurface spatial distribution of soil organic carbon (SOC), total nitrogen (N_{tot}), pH in water and KCl (pH_w and pH_{KCl}), and total carbonates (CaCO₃) at the time of land-use conversion.

The thickness of soil layers was inferred from the average limits between the horizons, according to description of pedological pits. Soil samples were collected in 67 randomly selected georeferenced points from 4 layers: 0-15 cm (average harrowing thickness, corresponding to the Ap1 horizon), 15-35 cm (35 cm is the depth of recent ploughing), 35-60 cm (60 cm is the depth of the old deep ploughing for poplar planting), and 60-100 cm (corresponding to the C horizon, never affected by ploughing). The sampling was performed during the spring of 2010, using a gouge auger (30 mm diameter – Eijkelkamp, Giesbeek, The Netherlands).

Soil samples were air dried, sieved (2 mm mesh) and analysed to determine the SOC and N_{tot} content (Flash EA 1112 NCSOIL, Thermo Fisher Scientific elemental analyzer, Pittsburgh, USA), pH_{KCl} and pH_w (soil to KCl 1 M solution/water ratio of 1:2.5) and CaCO₃ (Dietrich-Frühling calcimeter, in the 0-15 cm layer only).

Along with the characterization of the initial soil conditions to monitor soil changes over time, the spatial variability study provided a guide for the selection of points for soil respiration (R_s) monitoring and comparison. Data related to the soil surface layer, from which the major part of a gas flux usually originates, were processed by geostatistical ordinary kriging techniques to map the spatial distribution of soil properties (Goovaerts 1997, Wackernagel 2003, Webster & Oliver 2007). The goodness-of-fit of interpolated values to the observed values at each sampling point was evaluated with cross-validation by calculating: (1) the mean error (ME), which proves the unbiasedness of an estimator if its value is close to 0; and (2) the variance of the root mean square standardized error (RMSSE). If the estimation is accurate, the variance of the RMSSE should vary within the 1 ± 3·√(2/n) interval, where n is the number of samples (Chilès & Delfiner 1997).

For each variable, an exploratory data analysis, including the calculation of descriptive statistics, identification of data distribution types and detection of outliers were performed.

All geostatistical analyses were performed using the ISATIS software package, release 13.01 (Géovariances 2013).

Soil respiration

For the comparison of soil respiration (R_s) between land uses, soil CO₂ flux was measured every 15-30 days (depending on the




Fig. 1 - Study area and field subdivision. (Alfalfa): crop-field cultivated with alfalfa; (Corn): crop-field cultivated with corn; (H-SRC): high-density short rotation coppice (1,111 trees ha⁻¹); (VH-SRC): very high-density short rotation coppice (8,333 trees ha⁻¹).

season) over a period of 19 months (April 2010 to October 2011) at 15 locations (see below). Measures were taken using plastic collars with a diameter of 10 cm and height of 6 cm. At the beginning of the experiment, the collars were inserted into the soil down to 2.5 cm from the soil surface and remained in place throughout the duration of the experiment. Collars were momentarily removed only during harrowing and repositioned immediately after.

The 15 monitoring sites were distributed throughout the experimental fields, with three locations for each land-use type, except for VH-SRC where each collar in the lane was coupled to a nearby collar in the row. The monitoring points for each land-use type were selected based on the spatial distribution maps of soil properties that were used to identify soils with comparable characteristics, so that the effects of land-use changes could be monitored. Since the soil texture was relatively uniform at the study site, the choice of monitoring sites was based on the spatial distribution of the SOC stock and rock fragment content.

The comparison of R_s between land uses was conducted for the same type of management (fertilization and irrigation), and therefore, only the FI thesis was considered for the VH-SRC.

The assessment of the influence of agronomic treatments on R_s was carried out on the same dates as the land-use comparison was conducted. In the VH-SRC, the soil CO_2 fluxes of the three experimental treatments (CTR, F, FI) were compared using the 6 collars already installed for land-use comparison. Moreover, 12 additional collars equally distributed between rows and lanes were placed at the CTR and F treatments, in order to deeper investigate the differences between rows and lanes in VH-SRC.

CO_2 soil fluxes at monitoring sites were measured using an EGM-3 portable infra-

red gas analyser (PP-Systems, Hitchin, UK) equipped with an SRC-1 chamber based on the principle of closed dynamic cylindrical systems with a diameter of 10 cm and volume of 1170 cm³. A small fan in the chamber ensured air mixing. For each monitoring day, a single measurement was performed at each sampling point. To reduce the effects of diurnal variation, measurements were taken during the central hours of the day within a time span of four hours. Simultaneously with R_s , soil water content and soil temperature (T_s) were measured at all points within 10 cm from the collar. Soil water content was measured with a portable TDR system (IMKO Micromodultechnik, GmbH, Ettlingen, Germany) using a 10 cm long trifurcated probe and the results were converted to water-filled pore space (WFPS, %). T_s was measured with the built-in temperature probe of the CO_2 analyser at a depth of 5 cm. Soil temperature was also measured at the same depth as T_s and was recorded continuously at the meteorological station of the experimental site.

The R_s and T_s data from each collar were fitted to exponential models (data not shown); consequently, cumulative respiration rates (R_{cum} , g C m⁻²) over the period of the experiment (1st of April 2010 – 31st of October 2011) were computed by integrating (hourly base) the R_s estimated as a function of the soil temperature. The average R_{cum} rates were computed for each land-use type. For the VH-SRC, the average R_{cum} rates were calculated, weighing the results obtained in the rows and lane.

At the end of the experiment (November 2011), the soil at all the gas monitoring points was sampled (0-15 cm) to determine the root density (RD) and SOC content. For successful separation of the roots from the soil, the soil samples were dispersed in a sodium hexametaphosphate solution. Roots were then washed with water and divided into three diameter classes: (i) fine (f, <2 mm); (ii) medium (m, 2-5 mm); (iii)

coarse (c, >5 mm). The roots were oven-dried at 105 °C for 48 h to determine their dry mass.

Soil bulk density (BD) at each monitoring point was determined for the 0-15 cm layers with the cylindrical core method (core diameter = 5.4 cm) on undisturbed core samples, considering the volume of the stones when present. Through the BD and stone volume, the SOC content on an area basis was computed for the mineral 0-15 cm layers in correspondence with the monitoring plots.

Statistical analyses

To compare soil properties between the monitoring locations and test for differences in R_{cum} and pedoclimate parameters between land-use and management types, the mixed effect model procedure was performed (Bolker et al. 2009) and autocorrelation among the model residuals was tested (Searle et al. 2009).

After verifying the absence of a residual spatial correlation (data not shown), the evaluation of each variable response in relation to land use or a treatment was carried out considering land use or treatment as a fixed effect in the linear mixed model. The differences between the mean values were obtained through contrast analysis. The restricted maximum likelihood (REML) procedure was used to estimate the variance components (Littell et al. 2006).


All statistical analyses were performed using the SAS[®] software (release 9.4, SAS Institute Inc., Cary, NC, USA).

Results

Spatial distribution of soil properties

The vertical distribution of the SOC content and pH_{KCl} is displayed in Fig. 2. The average SOC content (\pm standard deviation, SD) was 0.92 \pm 0.20% and 0.74 \pm 0.12% at the 0-15 cm and 0-35 cm depths, respectively, with differences in the vertical distri-

Fig. 2 - Vertical distribution of SOC (%) – left panel) and pH_{KCl} (right panel) mean values in the soil cores obtained from the survey of soil spatial variability at the four experimental fields. (VH-SRC): very high-density short rotation coppice; (H-SRC): high-density short rotation coppice; (SD): standard deviation.






Fig. 3 - Maps obtained by spatial interpolation (kriging) of soil parameters (0-15 cm layer). Black crosses represent the soil sampling points of the spatial survey.

bution between the investigated fields. The alfalfa field showed a significantly ($p < 0.05$) higher SOC value ($1.11 \pm 0.18\%$) in the 0-15 cm layer than in the 15-35 cm layer ($0.70 \pm 0.09\%$), while the remaining fields exhibited a more uniform vertical distribution in the ploughed layer (0-35 cm). Since old, deep ploughing had been made within 60 cm of depth, the SOC content was very low under this limit. The vertical trend of N was comparable with that of SOC (data not shown).

The pH_w average values increased with depth in all the fields (from 7.7 ± 0.1 to 8.5 ± 0.1); the vertical trend of pH_w was comparable with that of pH_{KCl} , except for the deepest layer where the values of pH_{KCl} were lower (corn and alfalfa) or nearly equal (SRCs) to those of the third layer. The average surface $CaCO_3$ content was $8.5 \pm 0.9\%$.

An isotropic variogram model was fitted to the experimental variograms of the surface (0-15 cm) SOC content and stock, pH_{KCl}

and coarse rock fragments. The variogram consisted of two basic structures, a nugget effect and a spherical model with 82, 78, 32 and 118 m ranges, respectively, for the four aforementioned variables. The variogram fitting was satisfactory because the mean error was close to 0 and the RMDDE was within the $1 \pm 3 \cdot \sqrt{(2/n)}$ interval.

The experimental variogram of $CaCO_3$ content showed a pure nugget effect, which indicated a complete lack of spatial correlation, at least at the sampling scale.

The spatial distribution maps of the SOC content and stock, as well as the pH_{KCl} and coarse rock fragments of the surface layer, are shown in Fig. 3. A clearly increasing gradient in rock fragment percentage was observed in the NW-SE direction, mainly in the eastern part of the study area where the SRC plantations are located. The SOC content in the SRC fields showed a similar spatial distribution trend, while for carbon stock such gradient was attenuated.

The alfalfa field showed a higher SOC sur-

face content and stock than the other fields. As for pH_{KCl} , a substantial homogeneity of values was observed throughout the study site.

Based on the obtained maps, the points for R_s monitoring were identified, taking care of selecting points as similar as possible in terms of properties affecting soil CO_2 emissions. The average values of BD, SOC stock and RDf (fine root density) of the 0-15 cm layer for each land-use and management type are shown in Tab. 1. Preliminarily, R_s monitoring points were selected in areas with no rock fragments; however, significant differences in BD were found, with the lowest values at the alfalfa field and the highest values at the VH-SRC field. No significant differences were found in SOC stock and RDf among the selected monitoring points.

Soil respiration

During the monitored period, the measured R_s rates ranged from 0.07 to $8.2 \mu mol CO_2 m^{-2} s^{-1}$ in the corn field, from 0.11 to $6.8 \mu mol CO_2 m^{-2} s^{-1}$ in the alfalfa field, from 0.19 to $5.2 \mu mol CO_2 m^{-2} s^{-1}$ in the VH-SRC (FI) and from 0.34 to $6.2 \mu mol CO_2 m^{-2} s^{-1}$ in the H-SRC (Fig. 4a). The different agronomic treatments in the VH-SRC showed similar soil CO_2 emissions (Fig. 4b) with average values (\pm standard deviation) of 3.3 ± 1.4 , 2.8 ± 1.5 and $3.2 \pm 1.2 \mu mol CO_2 m^{-2} s^{-1}$ in the FI, F and CTR treatments, respectively. R_s generally showed a typical seasonal pattern following soil temperature (T_s), with the highest soil CO_2 fluxes during the summer months and the lowest during the winter months.

The comparison of average R_{cum} , T_s and WFPS (water-filled pore space) values for

Tab. 1 - Average values (\pm standard deviation) of soil bulk density (BD), soil organic content (SOC) stock and fine root density (RDf) at monitoring points for each land use (VH-SRC, H-SRC, corn, alfalfa – see text) and agronomic treatment (CTR: control; F: fertilization; FI: fertilization and irrigation). Different letters denote significant ($p < 0.05$) differences between land uses. Number of monitoring points: 3 couples (one in the lane and one in the row) of collars for the VH-SRCs and 3 collars for all the other land uses. Average values at the VH-SRC were obtained by weighing the results in rows and lanes.

Variable	VH-SRC			H-SRC	Corn	Alfalfa
	CTR	F	FI			
BD	1.6 ± 0.1^a	1.5 ± 0.1^{ab}	1.6 ± 0.1^a	1.4 ± 0.1^b	1.4 ± 0.1^b	1.3 ± 0.1^c
RDf	0.6 ± 0.3^a	0.4 ± 0.1^a	0.9 ± 0.5^a	0.6 ± 0.3^a	0.1 ± 0.4^a	0.5 ± 0.1^a
SOC stock	1.7 ± 0.1^a	1.6 ± 0.2^a	1.5 ± 0.1^a	1.5 ± 0.1^a	1.6 ± 0.1^a	1.8 ± 0.2^a





Fig. 4 - Comparison of the temporal trends in soil CO₂ emissions ($\mu\text{mol m}^{-2} \text{s}^{-1}$) between land uses (a) and treatments (b).

Fig. 5 - Comparison of soil parameters among land use types (a) and agronomic treatments (b). (R_{cum}): soil cumulative respiration rate (g C m^{-2}); (T_s): soil temperature ($^{\circ}\text{C}$); (WFPS): water filled pore space (%). Treatments: (CTR) control; (F) fertilization and irrigation. Different letters denote significant differences ($p < 0.05$) between means. Number of sampling points: four for each VH-SRC treatment and three for H-SRC, corn and alfalfa.



the analysed land uses are shown in Fig. 5a. The corn field, which was more exposed to solar radiation, showed an average T_s significantly higher than that measured in the other plots, followed by the H-SRC and alfalfa fields, whose T_s were comparable to

each other and significantly different from the VH-SRC field. The WFPS in the H-SRC was the lowest.

The R_{cum} in the alfalfa field ($1689 \pm 100 \text{ g C m}^{-2}$) was significantly higher ($p < 0.05$) compared with that of the other land uses. The

H-SRC and corn field showed similar CO₂ fluxes (approximately 1500 g C m^{-2}), which were significantly higher than that measured at VH-SRC plantation ($1299 \pm 30 \text{ g C m}^{-2}$).

The agronomic treatments tested inside the VH-SRC plantation did not result in differences in the R_{cum} or in differences in the T_s and WFPS (Fig. 5b).

In the VH-SRC stand, the rows and lanes were compared (Tab. 2). The cumulative CO₂ emissions in the rows were significantly higher (p value < 0.05) than in the lanes. The SOC was higher ($p < 0.10$) in the rows, and the BD was higher ($p < 0.10$) in the lanes.

Discussion

As previously reported by many studies, soil heterogeneity is elevated even at small spatial scales. Therefore, the adoption of

Tab. 2 - Comparison of soil parameter values (average \pm standard deviation) between rows and lanes at the VH-SRC. Number of sampling points: nine for rows and nine for lanes. (1): 0-15 cm; (2): 0-5 cm; (3): 0-10 cm.

Label	Parameter	Units	Rows	Lanes	p-value
R_{cum}	Cumulative soil respiration	g C m^{-2}	1621 ± 200	1274 ± 60	0.002
SOC ⁽¹⁾	Soil organic C	g kg^{-1}	7.7 ± 0.8	6.8 ± 0.4	0.095
RDf ⁽¹⁾	Fine root density	g dm^{-3}	0.78 ± 0.09	0.62 ± 0.08	0.656
BD ⁽¹⁾	Bulk density	g cm^{-3}	1.5 ± 0.1	1.6 ± 0.1	0.092
T_s ⁽²⁾	Soil temperature	$^{\circ}\text{C}$	18.02 ± 0.76	18.49 ± 0.98	0.406
WFPS ⁽³⁾	Water filled pore space	%	46.1 ± 3.5	43.5 ± 2.8	0.294

an adequate sampling strategy is crucial for monitoring organic carbon changes and respiration rates in soils (Saiz et al. 2006, Rodeghiero & Cescatti 2008). For our purposes, the preliminary analysis of soil spatial variability made it possible to identify areas with soil characteristics as much similar as possible, so that soil CO₂ fluxes could be correctly monitored and the effects of land-use changes appropriately investigated. Indeed, the spatial survey revealed a degree of soil variability that could have invalidated any conclusions related to the effects of the land-use changes and management if the plot choice had been random. For example, rock fragment content in the study area showed a spatial variability not connected to the land-use change but to the alluvial origin of the soil. Rock fragments were absent in about half of the study area but strongly influenced the remaining study area. In fact, rock fragment content significantly affected the soil bulk density and SOC stock (Schrumpf et al. 2011, Zabowski et al. 2011), and therefore, the knowledge of its spatial distribution was critical for the identification of comparable R_s in the monitored area.

The observed SOC enrichment in the alfalfa field was probably the consequence of the previous four years of alfalfa cultivation in the same field. In forage production systems, perennial grasses such as alfalfa increase SOM by increasing root biomass and stimulating microbial growth in the rhizosphere (Paustian et al. 1997, Sánchez-Moreno 2002). In such systems, where aboveground biomass is removed through mowing, roots and rhizodeposit-C are the primary sources of SOC (Van Der Krift & Berendse 2002). After land use conversion (2009), the portion of the study area kept with alfalfa cultivation was not ploughed, so that it could be part of the 5-year rotation period. This could have favored the accumulation of SOC in the soil surface layers (0-15 cm), in contrast with deeper layers (0-35 cm). At the end of the rotation period, alfalfa will be replaced by corn; pre-sowing soil preparations include ploughing, which result in a sudden change in the vertical SOC distribution by mixing the upper enriched layer with the deeper depleted one (up to 35 cm). This will likely promote the decomposition of SOC because of the increase in soil aeration.

Due to the heterogeneity of the above-mentioned soil properties, we chose to locate the points for monitoring soil respiration in the zones with no rock fragments and similar in SOC stocks. During the monitoring period (April 1st 2010 – October 31st 2011), the calculated SRC cumulative respiration values averaged 1351 ± 56 and 1459 ± 11 g C m⁻² for the VH and H stands, respectively. Reporting such values on an annual scale, the observed fluxes were higher than those measured by Verlinden et al. (2013a, 2013b) in an SRC poplar plantation of the same age and characterized by a similar SOC, and comparable to values

reported by Vande Walle et al. (2007) for an older SRC established on former agricultural land. The latter study also reported a low RDf (fine root density) in the surface layer, similar to that observed in our experimental area, which can be explained by the young age of the plantations.

In our study, only the fine root density was considered, as fine roots were predominant and had higher respiration rates compared to larger roots (Pregitzer et al. 1998, Desrochers et al. 2002). However, poor rooting was observed in croplands as well, likely suggesting a small autotrophic contribution to the total R_s over the whole experimental area. The relatively low root biomass and the short duration of live roots for annual crops usually results in a low contribution of root respiration in croplands (Raich & Tufekciogul 2000).

According to Verlinden et al. (2013a, 2013b), our SRCs may present characteristics more similar to croplands than to forest stands, where root contributions are generally higher. This could be the consequence of their young age and the tillage (although of low intensity) applied during the cultivation period. Indeed, we observed lower R_{cum} values at the VH-SRC stand, while for the H-SRC stand R_{cum} was similar to that recorded at the corn field.

The higher SOC content and the consequent lower bulk density (BD) in the upper soil layer of the alfalfa plot were mainly responsible for the highest R_{cum} values. As alfalfa has the ability to fix atmospheric N, it is possible that soil respiration (R_s) was also affected by the biological N fixation process, since that N-fixing bacteria oxidize the organic C to obtain energy, thus releasing large amounts of CO₂ (Hopkins 1999).

Nonetheless, our results showed that the differences in R_s among land uses may also be explained by differences in soil temperature (Rodeghiero & Cescatti 2005, Gaumont-Guay et al. 2006). Vegetation cover can deeply alter T_s and thereby significantly influence the R_s rates (Raich & Tufekciogul 2000). Accordingly, we found that the average T_s at the VH-SRC plot was lower than that detected at the other monitoring sites, probably due to the high density of poplar trees and their canopy shading. Moreover, T_s at the H-SRC plot was similar to that of alfalfa rather than VH-SRC, likely because the lower tree density, together with the limited canopy extension of the young stand, led to a higher proportion of bare soil.

Two years after VH-SRC establishment, irrigation and fertilization treatments did not appear to affect soil respiration. However, rows in VH-SRC showed a higher cumulative respiration (R_{cum}) than lanes, and the higher R_{cum} coincided with higher SOC and RDf and lower soil BD. The differences between the rows and lanes were likely due to management practices. Indeed, the lanes were subjected to tillage and periodically compacted by tractor transit, while no tillage was performed in the rows, thus

avoiding SOC losses through mineralization and allowing the root and soil structure to develop. Moreover, fresh input of organic matter, in addition to that stemming from root turnover, was derived from the litter, whose biomass was higher near the row than in the lane. This input of organic matter, together with a higher (although not statistically significant) root biomass, may have contributed to higher soil CO₂ emissions from the rows, through a higher heterotrophic and autotrophic contributions.

It is likely that the SOC enrichment observed only two years after the SRC establishment will increase over time as a result of the type of soil management. The intensity of SRC management is lower in comparison with the management of annual crops such as corn. In the experimental SRCs, trees are cut every 2 and 5 years at VH-SRC and H-SRC, respectively, and trees resprout from stumps that are permanently left on-site. The SRCs are characterized by regular leaf litter and root input and thus a higher SOC input compared to conventional crops, where soil ploughing is required annually (corn) or every 5 years (alfalfa), and the epigeic biomass is completely harvested. Furthermore, the tree canopies provide, at least in the VH-SRC, a significant shading throughout the summer season, limiting the increase in soil temperature and consequently soil CO₂ emissions.

Conclusions

Our study provided evidence that: (i) the spatial variability in soil properties is an important feature to be considered in the selection of monitoring points for soil respiration; (ii) the average cumulative respiration rate (R_{cum}) was lower in the very high-density short rotation coppices (VH-SRC) than in croplands, and higher in the SRC rows than in lanes; (iii) no significant differences in R_{cum} were found between the high-density coppice (H-SRC) and corn field; (iv) two years after VH-SRC establishment, agronomic treatments including irrigation and fertilization did not appear to affect soil respiration; and (v) land-use change affected the vertical SOC distribution, soil bulk density and soil surface temperature, which were reflected in the differences in soil respiration.

Further studies are needed and will be performed to confirm our results, through soil resampling over time to monitor long-term variation of soil properties in the study area, aimed at better understanding the effect of recent conversion of arable lands to SRC.

Acknowledgements

This work was supported by the SUSCACE project. Special thanks to G. Faccioto and S. Bergante of CREA-PLF (Casale Monferrato, Italy), for their field support and site management information, and to G. Seufert, A. Meijide, M. Duerr and C. Gruening of the Air and Climate Unit of the JRC of the European Commission (Ispra, Varese,

Italy) for their assistance in the selection of the sampling design and loaning the instrumentation for soil respiration monitoring.

References

- Arevalo CBM, Bhatti JS, Chang SX, Sidders D (2011). Land use change effects on ecosystem carbon balance: from agricultural to hybrid poplar plantation. *Agriculture, Ecosystems and Environment* 141: 342-349. - doi: [10.1016/j.agee.2011.03.013](https://doi.org/10.1016/j.agee.2011.03.013)
- Boehmel C, Lewandowski I, Claupein W (2008). Comparing annual and perennial energy cropping systems with different management intensities. *Agricultural Systems* 96: 224-236. - doi: [10.1016/j.agsy.2007.08.004](https://doi.org/10.1016/j.agsy.2007.08.004)
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JSS (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution* 24: 127-135. - doi: [10.1016/j.tree.2008.10.008](https://doi.org/10.1016/j.tree.2008.10.008)
- Chilès J-P, Delfiner P (1999). Discrete exact simulation by the Fourier method. In: "Geostatistics Wollongong 96". Wollongong, Australia, pp. 258-269.
- Desrochers A, Landhäusser SM, Lieffers VJ (2002). Coarse and fine root respiration in aspen (*Populus tremuloides*). *Tree Physiology* 22: 725-732. - doi: [10.1093/treephys/22.10.725](https://doi.org/10.1093/treephys/22.10.725)
- Ferré C, Comolli R, Leip A, Seufert G (2014). Forest conversion to poplar plantation in a Lombardy floodplain (Italy): effects on soil organic carbon stock. *Biogeosciences* 11: 6483-6493. - doi: [10.5194/bg-11-6483-2014](https://doi.org/10.5194/bg-11-6483-2014)
- Ferré C, Castrignanò A, Comolli R (2015). Assessment of multi-scale soil-plant interactions in a poplar plantation using geostatistical data fusion techniques: relationships to soil respiration. *Plant and Soil* 390: 95-109. - doi: [10.1007/s11104-014-2368-2](https://doi.org/10.1007/s11104-014-2368-2)
- Gasol CM, Brun F, Mosso A, Rieradevall J, Gabarrell X (2010). Economic assessment and comparison of acacia energy crop with annual traditional crops in Southern Europe. *Energy Policy* 38: 592-597. - doi: [10.1016/j.enpol.2009.10.011](https://doi.org/10.1016/j.enpol.2009.10.011)
- Gaumont-Guay D, Black TA, Griffis TJ, Barr AG, Jassal RS, Nesic Z (2006). Interpreting the dependence of soil respiration on soil temperature and water content in a boreal aspen stand. *Agricultural and Forest Meteorology* 140: 220-235. - doi: [10.1016/j.agrformet.2006.08.003](https://doi.org/10.1016/j.agrformet.2006.08.003)
- Géovariances (2013) Isatis Technical Ref., ver. 13.01. Géovariances & Ecole Des Mines De Paris, Avon Cedex, France, pp. 260.
- Goovaerts P (1997). *Geostatistics for natural resources evaluation*. Oxford University Press, New York, USA, pp. 487.
- Grigal DF, Berguson WE (1998). Soil carbon changes associated with short-rotation systems. *Biomass and Bioenergy* 14: 371-377. - doi: [10.1016/S0961-9534\(97\)10073-3](https://doi.org/10.1016/S0961-9534(97)10073-3)
- Hopkins WG (1999). *Introduction to plant physiology* (2nd edn). John Wiley and Sons, New York, USA, pp. 512. [online] URL: <http://www.cabdirect.org/cabdirect/abstract/19980712656>
- IUSS Working Group (2015). World reference base for soil resource 2014 (Update 2015). World Soil Resources Reports no. 103, FAO, Rome, Italy, pp. 192.
- Karp A, Shield I (2008). Bioenergy from plants and the sustainable yield challenge. *New Phytologist* 179: 15-32. - doi: [10.1111/j.1469-8137.2008.02432.x](https://doi.org/10.1111/j.1469-8137.2008.02432.x)
- Kauter D, Lewandowski I, Claupein W (2003). Quantity and quality of harvestable biomass from *Populus* short rotation coppice for solid fuel use - a review of the physiological basis and management influences. *Biomass and Bioenergy* 24: 411-427. - doi: [10.1016/S0961-9534\(02\)00177-0](https://doi.org/10.1016/S0961-9534(02)00177-0)
- Littell RC, Stroup WW, Milliken GA, Wolfinger RD, Schabenberger O (2006). *SAS for mixed models* (2nd edn). SAS Institute Inc., Cary, NC, USA, pp. 814.
- Nassi O, Di Nasso N, Guidi W, Ragagliani G, Tozzini C, Bonari E (2010). Biomass production and energy balance of a 12-year-old short-rotation coppice poplar stand under different cutting cycles. *GCB Bioenergy* 2: 89-97. - doi: [10.1111/j.1757-1707.2010.01043.x](https://doi.org/10.1111/j.1757-1707.2010.01043.x)
- Paustian K, Collins HP, Paul EA (1997). Management controls on soil carbon. In: "Soil Organic Matter in Temperate Agroecosystems: Long-Term Experiments in North America" (Paul EA, Paustian K, Elliott ET, Cole CV eds). CRC Press, Boca Raton, FL, USA, pp. 15-49. [online] URL: <http://books.google.com/books?id=sEyzRPzPxCCEC>
- Peveřill KI, La Sparrow Reuter DJ (1999). *Soil analysis - An interpretation manual*. CSIRO publishing, Collingwood, Victoria, Australia, pp. 369.
- Poeplau C, Don A (2013). Sensitivity of soil organic carbon stocks and fractions to different land-use changes across Europe. *Geoderma* 192: 189-201. - doi: [10.1016/j.geoderma.2012.08.003](https://doi.org/10.1016/j.geoderma.2012.08.003)
- Pregitzer KS, Laskowski MJ, Burton AJ, Lessard VC, Zak DR (1998). Variation in sugar maple root respiration with root diameter and soil depth. *Tree Physiology* 18: 665-670. - doi: [10.1093/treephys/18.10.665](https://doi.org/10.1093/treephys/18.10.665)
- Raich JW, Tufekcioglu A (2000). Vegetation and soil respiration: correlations and controls. *Biogeochemistry* 48: 71-90. - doi: [10.1023/A:1006112000616](https://doi.org/10.1023/A:1006112000616)
- Rodeghiero M, Cescatti A (2005). Main determinants of forest soil respiration along an elevation/temperature gradient in the Italian Alps. *Global Change Biology* 11: 1024-1041. - doi: [10.1111/j.1365-2486.2005.00963.x](https://doi.org/10.1111/j.1365-2486.2005.00963.x)
- Rodeghiero M, Cescatti A (2008). Spatial variability and optimal sampling strategy of soil respiration. *Forest Ecology and Management* 255: 106-112. - doi: [10.1016/j.foreco.2007.08.025](https://doi.org/10.1016/j.foreco.2007.08.025)
- Saiz G, Green C, Butterbach-Bahl K, Kiese R, Avitabile V, Farrell EP (2006). Seasonal and spatial variability of soil respiration in four Sitka spruce stands. *Plant and Soil* 287: 161-176. - doi: [10.1007/s11104-006-9052-0](https://doi.org/10.1007/s11104-006-9052-0)
- Sánchez-Moreno C (2002). Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food Science and Technology International* 8: 121-137. - doi: [10.1177/10282013202008003770](https://doi.org/10.1177/10282013202008003770)
- Searle SR, Casella G, McCulloch CE (2009). *Variance components*. John Wiley and Sons, Hoboken, NJ, USA, pp. 501. [online] URL: <http://books.google.com/books?id=mkCSOzHMngC>
- Schrumpf M, Schulze ED, Kaiser K, Schumacher J (2011). How accurately can soil organic carbon stocks and stock changes be quantified by soil inventories? *Biogeosciences Discussions* 8: 723-769. - doi: [10.5194/bgd-8-723-2011](https://doi.org/10.5194/bgd-8-723-2011)
- Smith P (2004). Carbon sequestration in croplands: the potential in Europe and the global context. *European Journal of Agronomy* 20: 229-236. - doi: [10.1016/j.eja.2003.08.002](https://doi.org/10.1016/j.eja.2003.08.002)
- Van Der Krift TAJ, Berendse F (2002). Root life spans of four grass species from habitats differing in nutrient availability. *Functional Ecology* 16: 198-203. - doi: [10.1046/j.1365-2435.2002.00611.x](https://doi.org/10.1046/j.1365-2435.2002.00611.x)
- Vande Walle I, Samson R, Looman B, Verheyen K, Lemeur R (2007). Temporal variation and high-resolution spatial heterogeneity in soil CO₂ efflux in a short-rotation tree plantation. *Tree Physiology* 27: 837-848. - doi: [10.1093/treephys/27.6.837](https://doi.org/10.1093/treephys/27.6.837)
- Verlinden MS, Broeckx LS, Bulcke J, Acker J, Ceulemans R (2013a). Comparative study of biomass determinants of 12 poplar (*Populus*) genotypes in a high-density short-rotation culture. *Forest Ecology and Management* 307: 101-111. - doi: [10.1016/j.foreco.2013.06.062](https://doi.org/10.1016/j.foreco.2013.06.062)
- Verlinden MS, Broeckx LS, Wei H, Ceulemans R (2013b). Soil CO₂ efflux in a bioenergy plantation with fast-growing *Populus* trees-influence of former land use, inter-row spacing and genotype. *Plant and Soil* 369: 631-644. - doi: [10.1007/s11104-013-1604-5](https://doi.org/10.1007/s11104-013-1604-5)
- Wackernagel H (2003). *Multivariate geostatistics: an introduction with applications*. Springer, Berlin, Germany, pp. 387.
- Webster R, Oliver MA (2007). *Geostatistics for environmental scientists* (2nd edn). John Wiley and Sons, Chichester, West Sussex, UK, pp. 330.
- Zabowski D, Whitney N, Gurung J, Hatten J (2011). Total soil carbon in the coarse fraction and at depth. *Forest Science* 57: 11-18.
- Zenone T, Zona D, Gelfand I, Gielen B, Camino-Serrano M, Ceulemans R (2015). CO₂ uptake is offset by CH₄ and N₂O emissions in a poplar short-rotation coppice. *GCB Bioenergy* 8: 524-538. - doi: [10.1111/gcbb.12269](https://doi.org/10.1111/gcbb.12269)

Supplementary Material

Fig. S1 - Representative soil profiles: (a) Calcaric Regosol (Epiloamic, Endoarenic); (b) Endoskeletal Calcaric Regosol (Epiloamic, Endoarenic).

Tab. S1 - Main characteristics of soil profiles: Calcaric Regosol (Epiloamic, Endoarenic).

Tab. S2 - Main characteristics of soil profiles: Endoskeletal Calcaric Regosol (Epiloamic, Endoarenic).

Link: Ferre_2621@suppl001.pdf