
Hosseini Z, Naghavi H, Latifi H, Bakhtiarvand-Bakhtiari S (2019).

Estimating biomass and carbon sequestration of plantations around industrial areas using very high resolution stereo satellite imagery

iForest – Biogeosciences and Forestry – doi: 10.3832/ifor3155-012

## **Supplementary Material**


**Fig. S1** - Observed vs. predicted values for *Pinus eldarica* DBH. Bold red line is the 1:1 line. Dashed black line is the fitted line.



Estimating biomass and carbon sequestration of plantations around industrial areas using very high resolution stereo satellite imagery

iForest – Biogeosciences and Forestry – doi: 10.3832/ifor3155-012


**Fig. S2** - Observed vs. predicted values for species-specific heights. Line colors descriptions follow those of Fig. S1.



Estimating biomass and carbon sequestration of plantations around industrial areas using very high resolution stereo satellite imagery

iForest – Biogeosciences and Forestry – doi: 10.3832/ifor3155-012

**Fig. S3** - Lines are regression model for 4 species, response (x-axis) and predicted (y-axis) biomass values. Line colors descriptions follow those of Fig. S1.



Estimating biomass and carbon sequestration of plantations around industrial areas using very high resolution stereo satellite imagery

iForest – Biogeosciences and Forestry – doi: 10.3832/ifor3155-012

**Fig. S4** - Lines are regression model for 4 species, response (x-axis) and predicted (y-axis) dry biomass values. Line colors descriptions follow those of Fig. S1.

