

Collection: Cost Action E29 Meeting 2008 - Istanbul (Turkey)  
 Future Monitoring and Research Needs for Forest Ecosystems  
 Guest Editor: Marcus Schaub (WSL, Birmensdorf, CH)

## Deposition measurements and critical loads calculations: monitoring data, results and perspective

Lorenz M, Granke O

This paper presents the variation of sulphur and nitrogen deposition and the exceedance of their critical loads on Level II ICP Forests plots. The fact that critical loads are still exceeded at many forest sites in Europe indicates a continuing need for further implementation of air pollution abatement strategies. Such results contribute to the scientific basis for the development and reviews of the effectiveness of clean air politics by the Convention on Long-range Transboundary Air Pollution.

**Keywords:** Forest monitoring, deposition, nitrogen, sulphur, critical loads

### Introduction

Forest condition in Europe is assessed by the International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests (ICP Forests). Established by the Convention on Long-range Transboundary Air Pollution (CLRTAP) under the United Nations Economic Commission for Europe (UNECE), ICP Forests has been monitoring forest condition in a close co-operation with the European Commission (EC) for 22 years. Large-scale variations of forest condition over space and time in relation to natural and anthropogenic factors are assessed on more than 6000 plots systematically spread across Europe. For studies of causal relationships about 860 intensive monitoring plots were installed covering the most important forest ecosystems in Europe. Intensive monitoring data were submitted for 774 of these plots in 2006. Intensive monitoring includes crown condition, phenology, litterfall, foliar chemistry, tree growth, soil

condition and soil solution chemistry, deposition, ambient air quality, and meteorology. Risks of atmospheric nitrogen and sulphur deposition are assessed by means of calculations of critical loads and their exceedances. This article aims at the presentation of the variation of sulphur and nitrogen deposition and the exceedance of their critical loads on Level II plots. Such results contribute to the scientific basis for the development and reviews of the effectiveness of clean air politics by CLRTAP.

### Summary of methods

The monitoring methods applied by ICP Forests are harmonised among its 40 participating countries (ICP Forests 2006). Bulk deposition is measured in the open field close to the Level II plots. Deposition under canopy is derived from throughfall measurements. For the present study, annual bulk and throughfall deposition of nitrate nitrogen ( $\text{N-NO}_3^-$ ), ammonium nitrogen ( $\text{N-NH}_4^+$ ), sulphate sulphur ( $\text{S-SO}_4^{2-}$ ), sodium ( $\text{Na}^+$ ), chlorine ( $\text{Cl}^-$ ) and calcium ( $\text{Ca}^{2+}$ ) were calculated as the arithmetic means of the yearly sums of the deposition in the years 2003 to 2005 for each Level II plot in  $\text{kg ha}^{-1} \text{ yr}^{-1}$ .

Changes over time were calculated over the period 2000-2005. This period permitted the inclusion of a maximum number of plots under the aspect of data availability. With the years of assessment as predictor and annual deposition as target variable, linear trends over time were calculated for each plot.

Only plots were involved on which deposition had been measured continuously from 2000 to 2005, with maximally 30 days of measurements missing per year. Data of missing days were replaced by the average daily deposition of the respective year. Tab. 1 shows the numbers of included plots. The temporal variation of throughfall and bulk deposition of these three compounds was derived from annual arithmetic means over the respective plots. Depending on the pollutant and the form of deposition (bulk or throughfall) considered, between 185 and 249 plots were available.

It must be mentioned that measured throughfall underestimates total deposition. One reason is that total deposition comprises wet and dry deposition, but only part of the dry deposition is washed off the leaves and included in the throughfall. Moreover, throughfall measurements do not take into account canopy exchange in the sense of uptake and leaching of the leaves which is particularly pronounced for N. For the separation between internal circulation and deposition, models have been developed (Draaijers et al. 1996). Deviations between the modelled total deposition and throughfall measurements by ICP Forests were shown by De Vries et al. (2001). Because of relatively high uncertainties of the models (Erisman et al. 2005), throughfall measurements were left uncorrected for canopy exchange in the present study. In this respect it must also be mentioned that stem flow was not included in the analysis which may lead to an underestimation at least in beech forests.

### Results and discussion

Throughfall deposition of nitrate ( $\text{N-NO}_3^-$ ), ammonium ( $\text{N-NH}_4^+$ ), and sulphate ( $\text{S-SO}_4^{2-}$ ) shows marked spatial patterns across Europe. For the majority of those plots on which both throughfall and bulk deposition were measured, throughfall deposition is clearly higher than bulk deposition, indicat-

✉ Institute for World Forestry,  
 Leuschnerstr. 91, D-21031 Hamburg,  
 Germany


@ Martin Lorenz  
 (martin.lorenz@vti.bund.de)

Received: Mar 13, 2008 - Accepted: Dec 09, 2008

**Citation:** Lorenz M, Granke O, 2009. Deposition measurements and critical loads calculations: monitoring data, results and perspective. *iForest* 2: 11-14 [online: 2009-01-21] URL: <http://www.sisef.it/iforest/show.php?id=478>

**Tab. 1** - Numbers of plots for which deposition data were available.

Variation	Deposition	Na <sup>+</sup>	Cl <sup>-</sup>	Ca <sup>2+</sup>	N- NH <sub>4</sub> <sup>+</sup>	N- NO <sub>3</sub> <sup>-</sup>	S- SO <sub>4</sub> <sup>2-</sup>
Temporal (2000-2005)	Bulk	193	193	193	192	193	185
	Throughfall	223	223	223	222	223	215
Spatial (2003-2005)	Bulk	216	216	216	216	216	216
	Throughfall	249	249	249	249	249	249



**Fig. 1** - Bulk deposition (left) and throughfall (right) of  $\text{N-NO}_3^-$  on the Level II plots involved in the study.

ing dry deposition filtered from the air by the canopy and washed off the leaves.

Fig. 1 shows two maps for bulk deposition (left) and throughfall (right) of  $\text{N-NO}_3^-$  on the Level II plots involved in the study. Throughfall is clearly higher and its spatial pattern is more pronounced than that of bulk deposition: Throughfall ranges from 0.2 to  $16.9 \text{ kg ha}^{-1} \text{ yr}^{-1}$ . The plots with the highest throughfall ( $6.3$  to  $16.9 \text{ kg ha}^{-1} \text{ yr}^{-1}$ ) are largely situated in areas of high vehicle exhaust, namely central Germany, Belgium and northern Italy. In contrast, plots with lowest throughfall ( $0.2$  to  $1.8 \text{ kg ha}^{-1} \text{ yr}^{-1}$ ) are located mainly in northern Europe and in the Alps. In comparison to  $\text{N-NO}_3^-$ , throughfall of  $\text{N-NH}_4^+$  shows a similar spatial pattern (not shown). But plots with highest throughfall of  $\text{N-NH}_4^+$  ( $7.5$  to  $23.8 \text{ kg ha}^{-1} \text{ yr}^{-1}$ ) are less frequent in central Germany and in northern Italy.

Throughfall of  $\text{S-SO}_4$  (not shown) ranges from  $0.7$  to  $42.7 \text{ kg ha}^{-1} \text{ yr}^{-1}$ . Plots with the highest throughfall ranging from  $8.0$  to  $42.7 \text{ kg ha}^{-1} \text{ yr}^{-1}$  are particularly frequent in central Europe. Sulphate deposition was not corrected for sea salt contribution. On plots in coastal areas (e.g., in Belgium, Cyprus, Greece, Italy, south-western Norway, north-


western Spain and in the United Kingdom) high sulphate deposition coincides with high sodium deposition, indicating sea salt as an origin. In the other regions the high deposition is likely to reflect regional emission situations. Similar to the nitrogen compounds, throughfall of  $\text{S-SO}_4^{2-}$  is particularly low ( $0.7$  to  $3.3 \text{ kg ha}^{-1} \text{ yr}^{-1}$ ) in northern Europe and in the Alps.

Temporal variation of bulk deposition and throughfall of the analysed substances (not shown) revealed a year by year (except 2004) decrease for  $\text{S-SO}_4^{2-}$ . From 2000 to 2005 bulk deposition decreased from  $6.1$  to  $4.6 \text{ kg ha}^{-1} \text{ yr}^{-1}$  and throughfall decreased from  $7.8$  to  $5.9 \text{ kg ha}^{-1} \text{ yr}^{-1}$ . This time series is too short for the quantification of a trend, but previous studies by ICP Forests show that the decrease in sulphur deposition has been ongoing for a longer time than the six years' period investigated in the present study (Lorenz et al. 2007, De Vries et al. 2001). It may therefore be stated that the results of ICP Forests give evidence of the reduction of sulphur emissions under CLRTAP politics over the last years and the less pronounced reduction of N emissions in Europe (Sliggers & Kakebeeke 2004).

Mere deposition measurements do not per-

mit any risk assessment for forest ecosystems. An approach for the assessment of the risk of forests damage by air pollution is the calculation of critical loads of deposition (Nilsson et al. 1986, Nilsson & Grennfelt 1988) and the determination of their exceedances. Therefore, the deposition data from Level II plots were used in previous studies of ICP Forests for calculations of exceedances of critical loads for each pollutant (Lorenz et al. 2008). Results of these studies show that on the investigated Level II plots critical load exceedances are smaller for sulphur than for nitrogen. Throughfall exceeds the critical loads for nitrogen on about two thirds of the plots (Fig. 2). Highest exceedances are found in The Netherlands, Belgium and in several parts of Germany. Hardly any exceedances are found in the United Kingdom, in Fennoscandia, in Greece and in the Alps. In the United Kingdom and Greece this reflects the low nitrogen deposition on Level II plots (Lorenz et al. 2007). Even in Fennoscandia, where critical loads are notoriously low, there are little or no exceedances because of low nitrogen deposition. However, the high precipitation in the Alps and lower Alps causes high leaching yielding high critical loads and low ex-





**Fig. 2** - Exceedances of critical loads for nitrogen assessed in 2004 (Lorenz et al. 2008).

ceedances, even if deposition is relatively high. This leads to the contradiction that low exceedances of critical loads for nitrogen deposition suggest a low risk on the one hand, while on the other hand high leaching of inorganic nitrogen constitutes a disturbance of the forest ecosystem. Critical loads for acidity were exceeded on less than a quarter of the plots. The plots are largely situated in The Netherlands, in southern Sweden, in several parts of Germany, and in Hungary.

The fact that critical loads are still exceeded at many forest sites in Europe indicates a continuing need for further implementation of air pollution abatement strategies.

In order to improve the reliability of the results, the potential of improving the modelling of canopy ion exchange should be considered.

As regards forest ecosystem response to deposition, it must be noted that the exceedance of critical loads indicates only that a pollutant flux is reached which may after a certain time lead to the exceedance of a critical limit, *i.e.*, to the exceedance of a pollutant concentration causing damage to the ecosystem. For the assessment of ecosystem response it is therefore crucial to know whether the critical limits are violated or not. Future studies of ICP Forests should therefore

focus on violations of critical limits and their relationships with forest ecosystem response.

### Acknowledgements

The data used for the described study were assessed and submitted within ICP Forests and under relevant EC Regulations by Austria, Belgium, Czech Republic, Denmark, Finland, France, Germany, Greece, Italy, Luxembourg, The Netherlands, Norway, Portugal, Slovakia, Spain, Sweden, Switzerland, and United Kingdom. Addresses of the National Focal Centres of ICP Forests are listed under <http://www.icp-forests.org>.

## References

- De Vries W, Reinds GJ, van der Salm C, Draaijers GPJ, Bleeker A, Erisman JW, Auee J, Gundersen P, Kristensen HL, Van Dobben H, De Zwart D, Derome J, Voogd JCH, Vel EM (2001). Intensive monitoring of forest ecosystems in Europe. Technical Report 2001. UNECE/EC, Geneva, Brussels.
- Draaijers G, Erisman J, Spranger T, Wyers G (1996). The application of throughfall measurements for atmospheric deposition monitoring. *Atmospheric Environment* 30: 3349-3361. - doi: [10.1016/1352-2310\(96\)00030-1](https://doi.org/10.1016/1352-2310(96)00030-1)
- Erisman J, Vermeulen A, Hensen A, Flechard C, Dämmgen U, Fowler D, Sutton M, Grünhage L, Tuovinen J (2005). Monitoring and modelling of biosphere/atmosphere exchange of gases and aerosols in Europe. *Environmental Pollution* 133 (3): 403-413. - doi: [10.1016/j.envpol.2004.07.-004](https://doi.org/10.1016/j.envpol.2004.07.-004)
- ICP Forests (2006). Manual on methodologies and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests. Federal Research Centre for Forestry and Forest Products (BFH), Hamburg, Germany. [online] URL: <http://www.icp-forests.org>
- Lorenz M (2004). Monitoring of forest condition in Europe: impact of nitrogen and sulphur deposition on forest ecosystems. In: "Towards the sustainable use of Europe's Forests - forest ecosystem and landscape research: scientific challenges and opportunities" (Andersson F, Birot Y, Päivinen R eds.). European Forest Institute 2004, S. 73-83, EFI-Proceedings Nr. 49, Joensuu, Finland.
- Lorenz M, Mues V (2007). Forest health status in Europe. *TheScientificWorldJournal* 7 (S1): 22-27. - doi: [10.1100/tsw.2007.17](https://doi.org/10.1100/tsw.2007.17)
- Lorenz M, Nagel H-D, Granke O, Kraft P (2008). Critical loads and their exceedances at intensive monitoring sites in Europe. *Environmental Pollution* 155: 426-435. - doi: [10.1016/j.envpol.2008.02.002](https://doi.org/10.1016/j.envpol.2008.02.002)
- Nilsson S, Andersen B, Dickson W, Eriksson E, Henriksen A, Kämeri J, Nilsson I (1986). Critical loads for nitrogen and sulphur. Report from a nordic working group. Nordic Council of Ministers, Miljø rapport 1986, 11, Stockholm, Sweden.
- Nilsson S, Grennfelt P (1988). Critical loads for sulphur and nitrogen. Report of a workshop held in Skogkloster. Nordic Council of Ministers, Miljø rapport 1988, 15, Stockholm, Sweden.
- Sliggers J, Kakebeeke W (2004). Clearing the air: 25 years of the Convention on Long-range Transboundary Air Pollution. United Nations, New York and Geneva.