

## The forest biodiversity artery: towards forest management for saproxylic conservation

Franco Mason<sup>(1-3)</sup>,  
Livia Zapponi<sup>(2-3)</sup>

One of the objectives of forest conservation is the set aside of unharvested areas. However, the fragmentation and lack of connectivity of protected areas make the integration of conservation measures in productive forests essential. Strategies to integrate conservation of saproxylic biodiversity in forest management have been developed, but often considering only specific aspects or remaining preliminary otherwise. As the impact of climate change and anthropogenic stresses increases, the development and the synthesis of this approach is crucial. We reviewed the key literature on forest management for biodiversity conservation, integrating forest science perspective to provide a practical management framework. Our goal is to present a management framework that could contribute to the effective preservation of forest insect biodiversity at the landscape scale, without high economic efforts, and addressing the conflicts that still jeopardize sustainable forest management. The results of our review support the creation of micro-reserves inside productive forests, to support large reserves in landscape conservation strategies. Micro-reserves increase the resilience of forest ecosystems to anthropogenic disturbances, through the development of a heterogeneous structure, maximizing microhabitat availability. Modeling forest management and harvest on local natural disturbance would extend the benefits of spatio-temporal heterogeneity in productive forests. Variable retention harvest systems, applied at the landscape scale, are a feasible and adaptable strategy to preserve and increase biodiversity, safeguarding structural legacies such as senescent trees and deadwood inside the productive matrix. The operational shift, from the stand to the forest landscape, is fundamental to extend the benefits of conservation measures. The Forest Biodiversity Artery, composed by several micro-reserves or *îlots de senescence*, connected by corridors of habitat trees and deadwood, constitutes a network that would deliver old-growth forests attributes to the productive matrix. This planning instrument would support forest connectivity, and socioeconomic constraints.

**Keywords:** Biodiversity, Deadwood, Gap, Habitat Tree, Integrative Conservation, Landscape, Microhabitat, Retention

### Introduction

Changes in land-use confined old-growth forest to less than 0.5% of the forested area in Europe, United States and China (Wirth et al. 2009). Structural and compositional changes that influenced forests in

the last centuries brought a heavy depletion of deadwood (Kaplan et al. 2009). Old-growth and deadwood-associated species (i.e., saproxylic) are among the most threatened in temperate forest ecosystems (Grove 2002). Species that require resour-

ces which have become rare in productive forests, such as large deadwood and open canopy areas (Seibold et al. 2015), may face a high risk of extinction. The uncertainties associated with future climate and anthropogenic stresses make the development of new forest conservation strategies imperative (Milad et al. 2011, Schaich & Milad 2013).

Traditionally conservation efforts brought the employment of natural forest as a model for restoration (Kuuluvainen 2002) and the set aside of unharvested areas rather than the restoration of harvested forests (Baker et al. 2013). Over the last 10 years the forest area allocated for biodiversity conservation in Europe increased of about half a million hectares every year (FOREST-EUROPE/UNECE/FAO 2011); however, these reserves tend to be isolated and surrounded by intensively managed landscapes (Branquart et al. 2008). Since state reserves cannot guarantee the survival of forest biodiversity (Miklín & Cížek 2014), the integration of conservation measures in pro-

---

□ (1) National Center for Study and Conservation of Forest Biodiversity “Bosco Fontana”, Laboratory for the Taxonomy of Invertebrates (CNBFVR-LANABIT). Via Tomaso Da Vico 1, 37123 Verona (Italy); (2) National Center for the Study and Conservation of Forest Biodiversity “Bosco Fontana”, Strada Mantova 29, 46045 Marmirolo, MN (Italy); (3) National Research Council (CNR), Agro-environmental and Forest Biology Institute, Via Salaria Km 29.300, 00015 Monterotondo Scalo, RM (Italy)

@ Franco Mason ([franco.mason.cnbfvr@gmail.com](mailto:franco.mason.cnbfvr@gmail.com))

Received: Apr 01, 2015 - Accepted: Aug 18, 2015

**Citation:** Mason F, Zapponi L (2015). The forest biodiversity artery: towards forest management for saproxylic conservation. *iForest* 9: 205-216. - doi: [10.3832/ifor1657-008](https://doi.org/10.3832/ifor1657-008) [online 2015-10-26]

Communicated by: Massimo Faccoli

**Tab. 1** - Used term combinations for the literature search. (\*): wildcard character used for boolean search.

Term 1	Term 2: Management	Term 2: Conservation
Forest	Certification	Biodiversity
Saproxylitic*	AND Disturbance	OR Biodiversity conservation
	Dynamic	Deadwood
	Eco unit	Habitat tree*
	Gap harvest	Hollow tree*
	Landscape approach	Îlot* de sénescence
	Landscape management	Old-growth
	Plantation	Structural legacies
	Retention harvest	Tree retention
	Sustainable management	Tree microhabitat*

ductive forests represents an urgent and vital intervention.

Since the establishment in 1811 of the Royal Saxon Academy of Forestry in Tharandt by Heinrich Cotta, the concept of sustainability, in its former sense of a sustained yield of wood, has been a leading principle for managing forests in Europe (Richter 1950). The intensity of wood harvest depended on forest growth potential, to ensure a continuous supply of timber for various uses. Ecosystem approaches have been part of European forestry since its origin as a science in the 18<sup>th</sup> century, through the comprehensive study of forest ecology (Sayer & Maginnis 2013). Still, there is often a disparity between traditional silviculture and the complexity of forest ecosystems (Boncina 2011). Our aim was to derive a framework to preserve saproxylitic diversity, relying on the last 20 years of literature on biodiversity conservation in productive forests. The proposed management strategies were derived answering the following questions: (1) which features of old-growth forests, fundamental for saproxylitic conservation, could be feasibly applied to productive forests? (2) which harvest methods could be better reconciled with saproxylitic preservation? (3) what are the current strategies to conserve saproxylitic biodiversity in productive forests? and (4) what should be the scale of intervention?

## Methods

For the selection of the reviewed studies, Oldeman (1990), Kohm & Franklin (1997) and Lindenmayer & Franklin (2002) works were used as guiding principles. Their works provided the terms listed in Tab. 1, which were used to interrogate three databases: (i) ISI Web of Science; (ii) Scopus; and (iii) Google Scholar. The obtained literature was then screened: no restrictions related to the time since publication were applied, whereas the selection was based on the relevance of the studies, focusing on European forests. The term “old-growth” is used in this article for both managed and unmanaged forests in Europe and generally applies to stands with trees older than 200 years, therefore “old-growth” is not synonymous of “virgin” (Peterken 1996).

## Forest management and conservation

### *Translating old-growth forest features into management principles*

After the 1970s, the idea that forests were constant and stable systems was abandoned, recognizing the complexity associated to the different kinds of disturbance (Kuuluvainen 2002). Oldeman (1990) seminal work highlighted how natural disturbances (e.g., wind storms, flooding, biotic-induced tree mortality) represent crucial processes and a permanent element of natural forest ecosystems. Far from being negative to the forest, these different types of disturbances are one of the main drivers of biological diversity. Stand heterogeneity is the result of gap dynamics, which is in turn influenced by small-scale (e.g., death of a single tree) and high-energy (e.g., wind throw) events (Bobiec et al. 2000). The interaction between climate, terrain and natural disturbance generates the spatial and structural heterogeneity of unmanaged forests (Lindenmayer & Franklin 2002). In shady and coarse woody debris (CWD) deprived managed forests, wind throw gaps act as biodiversity hotspots, providing deadwood and openings, which offer micro-habitats and trophic resources for a range of organisms (Bouget & Duelli 2004). The early-seral stages post disturbance, with the occurrence of natural regeneration, have a positive influence on saproxylitic composition (Horak 2015, Winter et al. 2015). Furthermore, the results presented by Beudert et al. (2015) show that the “benign neglect” approach towards pest outbreaks not only increases the overall biodiversity of a site but also does not threaten water quality in watersheds. Disturbance remnants, i.e., patches of surviving trees, may play a fundamental role in the recovery of large-scale and high-energy disturbances (Seidl et al. 2014a).

Oldeman’s view roots on the subdivision of forests in eco-units, forest areas whose development started at a unique moment in time, and “which architecture, eco-physiological functioning and species composition are ordained by one set of trees until the end”. Even if the pathways of forest development are countless and influenced

by chance, succession models have been derived to group them and summarize the involved processes (Donato et al. 2012). Oldeman (1990) identifies four phases forming the sylvatic mosaic: (i) innovation, (ii) aggradation, (iii) biostatic, and (iv) degradation. Likewise, Larrieu et al. (2014a) compared data from 32 unharvested European mountain mixed forests, and experimentally identified five forest development phases (FDPs): (i) regeneration, (ii) establishing, (iii) growing, (iv) culmination, and (v) disintegration. Hence, the forest is a mosaic of patches, characterized by their evolution and maturity (Schnitzler & Closeset 2003): this dynamics in space and time contribute to the development of a heterogeneous environment, characterized by multiple functional cycles (i.e., silvigenic, humus, pedofauna – Arpin et al. 1998). The sylvatic mosaic (*sensu* Oldeman 1990) of a temperate natural old-growth forest is characterized by a kaleidoscope of different kinds of eco-unit in dynamic equilibrium: the structural difference between even-aged (managed, mixed or pure) and uneven-aged forests (natural forest) is related to the size and spatial distribution of the tree cohorts (Oldeman 1990).

A silvicultural practice based on the occurrence of the different eco-units or FDPs, established considering reference conditions, would sustain structural biodiversity conservation (Winter & Brambach 2011). The explicit inclusion of natural eco-units or FDPs in forest management planning could favor the incorporation of the often missing culmination and disintegration phases, fundamental to guarantee high levels of microhabitats and deadwood (Larrieu et al. 2014b). Furthermore, natural early-seral stages can present old-growth forest structural attributes, thus where forest management focuses on ecological complexity this phase should be valued as well (Donato et al. 2012).

The maintenance of natural forest regeneration, together with the protection of soil productivity, represent the heart of nature-based silviculture (Boncina 2011). From the early conceptualizations of a close-to-nature silviculture (Otto 1993), it was clear that the application of a general panacea model was not appropriate: local, regional and temporal features of forests should act as guiding management principles. Models should favor the resilience of the forest, through the development of a heterogeneous structure, mimicking natural disturbances through gap harvest (Runkle 1991, Coates & Burton 1997, Mason 2002, 2004). As it has been stressed by Shorohova et al. (2011), the management and restoration of old growth-forests cannot be accomplished reproducing a static phase; conversely, it should focus on the emulation of natural disturbances and successional dynamics. Besides, silvicultural treatments that emulate natural disturbances, creating gaps in multi-aged forests, may increase resistance and resilience to future

disturbance (O'Hara & Ramage 2013).

Senescent and dead trees, together with snags and logs left on the forest floor, constitute what Franklin et al. (2000) defined "structural legacies", *i.e.*, elements that increase the post-disturbance complexity and provide habitat and food resources, promoting the survival and re-establishment of forest organisms (Ausden 2007). Regardless the considered geographic scale, the amount of available deadwood influences the functional composition of saproxylic organisms (Gossner et al. 2013). Despite the inclusion of deadwood volume among the improved Pan-European indicators for sustainable forest management (MCPFE 2003), "forest hygiene" still threatens its preservation. However, Stokland et al. (2012) stressed that both the risk for visitors and that of fires could be reduced by management, and that pest outbreaks (*e.g.*, *Ips typographus*) represent exceptional events related to lowland spruce introduction and introduced pest species.

Gossner et al. (2013) argue that close-to-nature approaches may fail the attempt of preserving saproxylic beetle biodiversity, disregarding the relevance of large diameter and late stage of decay deadwood availability. The concurring effects of increased harvest, biomass extraction, and assumed pest management reduced the volume of available deadwood by 90% (Jonsson & Siitonen 2012). Furthermore, the landscape scale removal of fuel wood endangers saproxylic organisms and other organisms affected by the physical and chemical changes of soil properties (see Bouget et al. 2012 for a review).

### The productive forest matrix and its controversial role for biodiversity preservation

Even if the threat represented by the management of a single forest stand is rather limited, the cumulative effect of these activities across a landscape may be significant (Villard & Jonsson 2009). Furthermore, the management intensity affects not only the present state of the stand but also its temporal sequence of states (Schall & Ammer 2013). The results obtained through the silvicultural management intensity indicator by Schall & Ammer (2013) highlighted that modern continuous-cover systems, such as selective logging, may preserve productivity without an intense environmental alteration. Nevertheless, canopy closure can affect the distribution of heliophilous species, thus the effect of the amount of retained trees on microclimate should be tested (Thorn et al. 2014).

### The variable retention harvest

In retention forestry, a part of trees and snags are left after the harvest, to create in young successional forests structures similar to those created by natural disturbance (Heikkala et al. 2014). The variable retention harvest (VRH), retaining structural, functional and compositional elements after the harvest, ensures: (1) lifeboating species that would otherwise disappear from a logged site, (2) the increase of stand structural complexity, and (3) the enhancement of forest landscape connectivity (Franklin et al. 1997, Gustafsson et al. 2012). The presence of a canopy closure gradient ensures the presence of a richer saproxylic fauna (Ulyshen et al. 2004, Gossner et al. 2006,

Lassauce et al. 2012). Together with agroforestry, retention forestry is considered an effective approach to reduce the conflict between biodiversity conservation and socio-economic needs at the landscape scale (Roberge et al. 2013). The suitability of the VRH in multifunctional forestry is guaranteed by its adaptability: it can be adjusted to meet different objectives and include distinctive local features, ranging from forest types to regional policy (Gustafsson et al. 2012). Gap-based approaches can be modeled on natural disturbance regimes: calibrating spatio-temporal components on local conditions in order to reach an effective and adaptive management (Runkle 1991, Coates & Burton 1997, Mercurio 1999, Lindenmayer & Franklin 2002, Mason 2004). There are several applications of this method (few examples are listed in Tab. 2) showing their diversified extent, aims and adaptability. Partial cuts may also have the advantage of ensuring the continual availability of deadwood, retaining large trees and snags through rotations (Fenton et al. 2013). The limits of retention forestry in preserving biodiversity include the lack of large-scale disturbances, as those generated by fire (Toivanen & Kotiaho 2007, Heikkala et al. 2014) and other chronic natural disturbances as windthrow, drought, heavy snowfalls, flooding or biotic factors as diseases and defoliations (Peterken 1996).


### The plantation opportunity

The general view on plantations is that they represent a sort of green desert. However, they can actually represent a potential habitat for biodiversity, since they re-

**Tab. 2** - Examples of gap based harvest with variable retention systems, applied to different forest types and various objectives.

Forest type	Aim	Harvest method	Unit size	Source
Boreal forest	Study the effect of green tree retention on forest biota	Retain 0, 10, 50 m <sup>3</sup> ha <sup>-1</sup> of the standing volume	3-5 tree groups	Hyvärinen et al. (2009)
Boreal mixed forests	Increase the density and vigour of spruce in aspen-dominated mixed woods	Alternate harvested and unharvested corridors	10 m wide	Man et al. (2010)
Boreal mixed forests	Evaluate regeneration after partial cutting and natural disturbance	Remove 47.9-63.2% of pre-harvest basal area	400 m <sup>2</sup>	Gendreau-Berthiaume et al. (2012)
Boreal mixed forests	Emulate natural disturbance, maintain beetle diversity	Remove 10-20% of the canopy in harvest gaps	0.1-0.2 ha	Thomas et al. (2009)
Boreal riparian forest	Emulate natural disturbance, increase habitat complexity through early succession forest regeneration	Partial harvest, up to 50% of basal area	10 - 400 m <sup>2</sup>	Mallik et al. (2014)
Floodplain forest	Natural reserve management emulating disturbances, saproxylic species conservation	Randomized small-gaps cutting with retention elements, waterlogging	250-300 m <sup>2</sup>	Mason (2004)
Mixed hardwood forest	Restore and maintain ground-layer vascular plant diversity in a second-growth forest	Dormant season timber harvesting in the gap and thinning outside	0 to 46 m of diameter	Kern et al. (2014)
Mixed temperate forests and plantations	Limit environmental impact, emulate disturbance, re-establish multi-layer structure and re-naturalization of conifer plantations	Eco-units with retention of deadwood	300-1000 m <sup>2</sup>	Mercurio et al. (2009)
Spruce-fir ( <i>Picea abies</i> - <i>Abies alba</i> )	Strengthen forest resilience to climate change increasing structural irregularity	Group selection with <50% of harvested trees (minimum diameter 52.5 cm)	500 m <sup>2</sup>	Lafond et al. (2014)
Lenga beech ( <i>Nothofagus pumilio</i> )	Assess regeneration under different microenvironmental conditions	Retain evenly distributed dominant trees between aggregates	30 m radius aggregates	Pastur et al. (2014)





**Fig. 1** - Representation of the proposed management system: the Forest Biodiversity Artery (FBA). The FBA is embedded in the productive matrix, showing the Deadwood Corridors (DC) and the *îlots de senescence* (IdSs). The productive forest matrix would be managed with gap harvest, with different eco-units or forest development phases. Landscape planning would facilitate the simultaneous presence of innovation/regeneration, aggradation/growing and biostatic/culmination phases (*sensu* Oldeman 1990 and Larrieu et al. 2014b, respectively). Deadwood thresholds and number of veteran trees in the productive matrix would be calibrated on the eco-unit/forest development phase, considering natural disturbance. The FBA would sustain saproxylic biodiversity providing high amounts of habitat trees and deadwood, diversified for decay and type. The landscape distribution of IdSs and DC would be established in order to ensure connectivity without affecting harvest operations (e.g., taking advantage of rides and glades, favoring sheerest areas).

semble the structural complexity of natural forests more than other more intense land uses (Gardner 2010). Their ecological value depends on the previous land use, alternative land uses, species involved and purpose (Brockhoff et al. 2008). Where reforestation entails the replacement of natural forests, plantations may be detrimental for conservation, whereas if established on anthropogenic grasslands that were once forested, they could confer environmental benefits to the landscape (Pawson et al. 2013). Plantations should be designed to deliver socio-economic benefits while providing ecological services, and should be included in a broad scale land-use planning (Paquette & Messier 2009). The extent and productivity of forest plantations are increasing worldwide, and the application of ecosystem approaches represents the vital and feasible strategy to support social and ecological outcomes (Sayer & Maginnis 2013). Koch Widerberg et al. (2012) suggest that the retention of oaks and deadwood in spruce plantations, along with the creation of small clearings, could represent a cost-effective intervention to preserve saproxylic beetle diversity. Plantations could contribute to the land sparing approach, segregating production in these areas and focusing conservation efforts in high quality natural habitats (Phalan et al. 2011, Tscharrntke et al. 2012). Land-sparing logging could be an effective strategy to preserve forest biodiversity in areas of large-scale interventions (Mayer et al. 2005, Edwards et al. 2014).

#### Current strategies: forest micro-reserves

Conservation efforts in productive forests include the protection of “tree islands”, where no harvest takes place (Aerts 2013). The *îlot de senescence* (IdS), described by Lachat & Büttler (2007), is a small and permanently unmanaged patch, distributed throughout managed forests. The IdS are a stepping stone that provide suitable habitat for saproxylic organisms. The retention of these groups of trees within a harvested forest ensures the availability of present and future deadwood, favoring both species with reduced dispersal and meta-community dynamics (Morrissey et al. 2014). Lachat & Büttler (2007, 2008) presented a practical design of forest conservation network, arguing that the conservation of saproxylic organisms in Switzerland could be achieved through the integration of large protected areas, IdSs and habitat trees. Further conceptualizations were presented by Cateau et al. (2013) for the Mont Ventoux (France), based on the balance between ecological, economic and social issues. For Fennoscandian countries, a similar setting aside of small habitat patches of productive forest to preserve landscape-level biodiversity was carried out from 1992 (Sverdrup-Thygeson et al. 2014a), naming these areas “Woodland Key Habitats” (Timonen et al. 2010). The small area of the IdS (ranging from one ha to a maximum of 20 ha) should not discourage on their effectiveness: small reserves can target high-quality remnants (Götmark & Thorell

2003), and their contribution to biodiversity conservation is increased by the fact that they are in non-wilderness areas (Lindenmayer & Franklin 2002). Where the establishment of protected areas is economically infeasible, biodiversity preservation could be achieved integrating reserves and managed matrix (Bengtsson et al. 2003, Shorohova et al. 2011). A further contribution could be represented by large marginal forests where management has been abandoned for socio-economic reasons (Sitzia et al. 2015). Nevertheless, the role of large reserves for the preservation of undisturbed ecosystems in optimum conditions is irreplaceable and cannot be granted by the matrix management alone (Lindenmayer & Franklin 2002): micro-reserves and natural reserves fulfill complementary functions and should be considered together in conservation strategies (Sverdrup-Thygeson et al. 2014a).

#### Scale of intervention: forest landscape conservation

Saproxylic species assemblages that characterize forest environments live in association with their surroundings, depending on heterogeneous conditions in time and space (Sverdrup-Thygeson et al. 2014b), and responding to substrate availability at various spatial scales (Bergman et al. 2012). The effectiveness of a micro-reserve depends on the intensity of the edge effect and the consequent disturbance (e.g., micro-climate alteration), as well as on the availability of “ecological memory” (Bengtsson et al. 2003), which ensures the resilience of a landscape through buffering and renewal. Planning and management need to be implemented at multiple spatial scales, from single trees to key habitats and finally to large reserves (Lindenmayer & Laurance 2012).

The general definition of stand, as an area characterized by its internal uniformity in terms of “composition, age, arrangement or condition” (Nieuwenhuis 2000) diverges deeply from Oldeman’s eco-unit. Forest ecosystem management at the stand level is a laudable objective, but alone it cannot ensure landscape biodiversity conservation (Bergeron et al. 1999). Hence, overcoming the concept of stand is fundamental to increase forest landscape permeability, favoring habitat diversity and reducing intensive land uses (Schaich & Milad 2013), as well as being flexible enough to accommodate wood market modifications (Seidl & Lexer 2013). The coordinated management of the components of a forest landscape ensures a more cost-effective and long-term use of conservation resources (Brumelis et al. 2011). Forest management should take into account that the spatial configuration of forest patches affects natural dynamics, ecosystem services and the sustainability of the extracted products (Kadiogullari et al. 2014). The extension of conservation measures in commercial and plantation forests needs a major operatio-

**Tab. 3** - Key elements favoring the preservation of forest biodiversity at the landscape scale.

Element	Function	Characteristics	Source
Îlot de senescence (IdS)	Conserving biodiversity in commercial forests	Part of the sylvatic mosaic, size established considering local disturbance dynamics. High availability of deadwood and veteran trees, never harvested	Lachat & Bütler (2007)
Deadwood corridor (DC)	Favoring the connections and meta-community dynamics	Approximate interior forest conditions	Warren & Fuller (1993)
Variable retention harvest (VRH)	Sustaining productive values without compromising ecosystem services	Gap harvest, with the retention of living structures and deadwood	See Tab. 2
Structural legacies	Lowering critical trophic and raising microhabitat resources	Coarse woody debris: chablis or uprooted trees, volis or snags, high and low stumps	Hedgren (2007), Abrahamsson & Lindbladh (2006)
Habitat tree	Life boating for species and ecosystem processes	Green-tree retention in the harvested gaps, preserving over-mature trees (DBH>90 cm) in the matrix and, indefinitely in the FBA	Franklin et al. (1997), Lachat & Bütler (2007)

nal shift: from stand-focused silviculture to a forest management that recognizes the landscape as the working unit (Pawson et al. 2013).

#### *A synthesis: the forest biodiversity artery and its incorporation in productive forests*

For the effective preservation of forest biodiversity, an extensive strategy integrating reserves and matrix management must be developed (Lindenmayer & Franklin 2002). The framework presented here aims at reaching an operational integration of biodiversity conservation in multi-functional forestry, and consists of two elements: a forest matrix managed with VRH in which the Forest Biodiversity Artery (FBA) is embedded. It represents an integrative conservation approach (*sensu* Bollmann & Braunisch 2013), which seeks to deliver old-growth forest attributes to the entire forest matrix. Structural legacies would be retained indefinitely inside the productive matrix, calibrating their number on the natural disturbance regimes and eco-unit/development phase. The FBA should be composed by several IdSs, connected each other by strips of similar habitats (habitat trees and deadwood corridors – DC). In the long term, the FBA system would constitute a network of old-growth forest patches integrated within productive forests (Fig. 1). The practical application of the proposed framework has been built up on the review presented in the previous chapters and is based on five main points, described below, which would ensure the simultaneous presence of the elements listed in Tab. 3.

#### **Incorporating natural disturbance**

Learning and inferring from natural models are key instruments to reach both economic production and biodiversity conservation (Lindenmayer & Franklin 2002). The amount and severity of disturbances associated with climate and land use changes has raised over the last decades, and understanding how these drivers interact is a prerequisite of their mitigation (Seidl et al. 2011). The frequency and severity of natural disturbance (both at regional and local

level) can constitute the preliminary data to establish retention approaches (Rosenvald & Löhms 2008) and small gap cutting treatments (Tab. 4). Disturbance regimes could be classified considering how probabilistic (e.g., fire intervals for a forest type) and random (e.g., where the fire happens) events interact (Angelstam 1998). The integration of these processes through spatio-temporal scales offers the means for the development of a more adaptive management (Nocentini 2011, Seidl et al. 2011), strengthening ecosystem resilience (Ciancio & Nocentini 2005, Milad et al. 2011). Seidl et al. (2014b) applied future scenario simulations and statistical modeling to show the upcoming intensification of disturbance regimes (due to fire, bark beetles and wind) under climate change, if management strategies remain unchanged. The deadwood created by these natural disturbances represents a valuable resource, which tends to host a higher biodiversity compared to deadwood artificially created (Komonen et al. 2014). Hence, policy measures should consider natural disturbance as a free conservation tool, allowing natural recovery and limiting salvage logging (Beudert et al. 2015). For example, the model LANDIS-II (<http://www.landis-ii.org/>) may simulate forest succession and natural/anthropogenic disturbances at the landscape level, and could be a useful tool for conservation planning (Leroux & Rayfield 2014). Managing a forest in terms of eco-units could represent a useful approximation that could allow the inclusion of the

diversity of compositional and structural phases. Among the selective cutting techniques, the most suitable treatments to restore the natural dynamics of the eco-units, characterizing the main part of the natural European forests (Peterken 1996), is represented by variable retention harvest (VRH).

#### **Criteria for the selection of retained structures**

Tree microhabitats are diverse (e.g., wounds, sap exudations, fungi fruiting bodies, cavities, cankers, bark pockets, etc.) and host a rich fauna (Siitonen 2012). Biodiversity could be preserved safeguarding valuable structures such as snags and veteran trees (Read 2000, Vuidot et al. 2011), which are particularly prone to develop such microhabitats. The same method adopted for selective logging (Giulietti et al. 2009) should be applied to identify the retained structures, focusing on the phenotypes showing age-related features, such as those identified by Lindenmayer & Franklin (2002): large-diameter trees (ensuring the development of deadwood cavities *per se*), complex canopies, rough bark. The criteria that could be applied for the selection of future habitat and hollow trees are listed in Tab. 5. Trees selected according to these characteristics would ensure a life-boating function (increasing microhabitat availability) and a structural enrichment (enhancing canopy and vegetation structure complexity – Franklin et al. 2000). The high value of a tree for biodiversity can

**Tab. 4** - Types of the small patch cutting as reviewed by Mercurio (1999). (D): larger diameter of the eco-unit; (H): predominant height of the trees in the eco-unit treatment; (NA): not available.

Eco-units		Source
Size (m <sup>2</sup> )	Shape	
500-1000	D = ½ H	Cappelli (1988)
< 400-500	Circular	De Philippis (1948)
600-1500	1-1.5 H	Del Favero et al. (1998)
NA	NA	Pavari (1956)
500-1500	NA	Perrin (1954)
1000-1500	D = 1-1.5 H	Piussi (1994)
NA	D = 2 H	Rousset (1972), Rojo Saiz (1977), Boudru (1989), Matthews (1989)

**Tab. 5** - Characteristics that increase the probability of microhabitat and hollow formation.

Tree species	Characteristics	Focus	Source
<i>Abies alba</i> , <i>Fagus sylvatica</i>	Minimum DBH: beech>90 cm, fir>100 cm	Montane forests	Larrieu & Cabanettes (2012)
<i>Eucalyptus</i> sp.	Stage of senescence, tree form and DBH determine the occurrence of hollows	Model probability of hollow occurrence	Rayner et al. (2014)
<i>Eucalyptus marginata</i>	Tree heights> 19 m, tree DBH>45 cm	Reptiles	Craig et al. (2011)
<i>Eucalyptus</i> sp., <i>Corymba</i> sp.	DBH>110 cm	Reptiles	Croak et al. (2013)
<i>Fagus sylvatica</i> , <i>Picea abies</i>	Minimum DBH: beech>72 cm, spruce>43 cm; the number of microhabitats increases for DBH>68 cm	Saproxylic beetles	Larrieu et al. (2012)
<i>Fagus sylvatica</i> , <i>Picea abies</i>	Minimum DBH: beech>50 cm, spruce>65 cm	Saproxylic beetles	Larrieu et al. (2014a)
<i>Fagus sylvatica</i> , <i>Quercus</i> sp., <i>Abies alba</i> , <i>Picea abies</i>	Snags host more microhabitats than living trees. More microhabitats on oaks than on the other species	Biodiversity	Vuidot et al. (2011)
<i>Nothofagus pumilio</i>	DBH 60.6 ± 16.4 cm	Birds	Diaz & Kitzberger (2013)
<i>Pinus sylvestris</i>	Minimum DBH for snags created from Scots pine to provide suitable cavity nest for birds and wildlife >40 cm	Birds and wildlife	Summers (2004)
<i>Pseudotsuga menziesii</i>	DBH≥70 cm	Birds and mammals	Michel & Winter (2009)
<i>Quercus ilex</i> , <i>Q. pubescens</i>	Good predictors: time since last cutting, diameter class. Non-coniferous species have more microhabitats	Birds, mammals and saproxylic beetles	Regnery et al. (2013)
<i>Quercus robur</i>	50% chance of hollow presence for trees 258 years old	Invertebrates, birds and mammals	Ranius et al. (2009)
Various	High DBH; balanced and pointed crown; straight trunk; dominant trees; local patterns related to altitude and slope	Wildlife conservation	Gibbons & Lindenmayer (2002)

also be determined focusing on its age, size and crown shape, or on the rarity of the species and its low mortality (Rosenvald & Löhmus 2008, Vuidot et al. 2011). Allowing a sub-population of trees to complete their natural cycle would ensure both the presence of microhabitats and the volume and diversity of deadwood (Larrieu et al. 2012). Large senescent trees could be excluded from the harvest applying DBH thresholds, thus the scarce high ecological value elements still present in productive

forests would never be logged (Aerts 2013). These elements have the key role of promoting the colonization of harvested sites, thus they should be as many as to sustain the species in this process (Baker et al. 2013). Management recommendations for the preservation of these structures are listed in Tab. 6.

#### Deadwood thresholds

The deadwood amount required to preserve biodiversity should be calibrated con-

sidering several taxonomical groups, taking into account the community composition rather than relying on the misleading species richness (Müller & Büttler 2010). Organisms that live in association with deadwood, a dynamic compound of living decaying trees and CWD (Bormann & Likens 1994) depend on a diverse and ephemeral substrate, and this inner stochasticity should be taken into account for its proper management (Jonsson et al. 2005). Müller & Büttler (2010) reviewed the dead-

**Tab. 6** - Management recommendations for the preservation and increase of habitat and hollow trees.

Tree species	Recommendations	Focus	Source
<i>Abies alba</i> , <i>Fagus sylvatica</i>	Favor mixed forests: differentiate microhabitats. Allow that a part of the trees complete their natural cycle	Montane forests	Larrieu & Cabanettes (2012)
<i>Eucalyptus</i> sp.	Time lag in the use of hollow-bearing trees in harvested areas: ensure landscape availability	Mammals	Cawthen & Munks (2011)
<i>Eucalyptus</i> sp.	Lag effects on the availability of hollows and conservation scenarios: model number of potential trees of high DBH/age	Birds	Manning et al. (2013)
<i>Fagus sylvatica</i>	Take into account spatial distribution, avoiding safety conflicts	Saproxylic beetles	Müller et al. (2014)
<i>Fagus sylvatica</i> , <i>Picea abies</i>	Include a sub-population (10-20% of the surface area) that will complete the natural silvigenetic cycle	Saproxylic beetles	Larrieu et al. (2012)
<i>Fagus sylvatica</i>	Preserve and create pollard trees	Saproxylics	Cantero et al. (2014)
<i>Pseudotsuga menziesii</i>	Microhabitat abundance is higher with low treatment history. Increase structural complexity: group harvest, small gap creation, retention of biological legacies	Birds and mammals	Michel & Winter (2009)
<i>Quercus ilex</i> , <i>Q. pubescens</i>	High stem density has a negative effect	Birds, mammals and saproxylic beetles	Regnery et al. (2013)
<i>Quercus robur</i>	Hollows generated early in fast-growing trees	Invertebrates, birds and mammals	Ranius et al. (2009)
<i>Quercus rubra</i> , <i>Platanus hybrida</i> , <i>Juglans nigra</i>	Guidelines for the artificial creation of microhabitats	Saproxylics	Mason et al. (2003)
<i>Salix alba</i>	Pollarding increases the probability of hollows formation	Saproxylic beetles	Sebek et al. (2013)
Broadleaf species	Hollow tree spatial distribution less than 100 m	Saproxylic beetles	Dubois et al. (2009)
Maturing hardwood forests	Uneven-aged harvest with group selection had a lower impact on hollow availability	Harvest regimes	Fan et al. (2004)
Various	Retain species with higher probability of developing hollows of survival	Wildlife	Gibbons & Lindenmayer (2002)
Various	Management guidelines in coppice	Insects	Fry & Lonsdale (1991)



wood threshold values recorded in the main forest types in Central Europe, suggesting that an amount of 20-50 m<sup>3</sup> ha<sup>-1</sup> should be guaranteed in several forest stands as part of a landscape network. This benchmark range could be used to calibrate the intervention according to productivity targets, higher wherever possible and lower to meet specific economic needs.

Forest restoration methods generally include tree girdling and felling to increase the availability of deadwood (Kuuluvainen et al. 2004) and tree microhabitats (Zapponi et al. 2015). “Morticulture”, the production of woody detritus to ensure ecosystem function (Harmon 2001), can be applied extending stand developmental models to deadwood dynamics in the considered forest management system (Jonsson & Siitonen 2012). The amount of deadwood could be artificially increased using the techniques presented by Cavalli & Mason (2003) and the retention of legacies (*sensu* Franklin et al. 2000). Among these legacies we can mention high stumps (Nordén et al. 2004, Abrahamsson & Lindbladh 2006, Brin et al. 2012), and *chablis* and *volis*, two French words full of forestry meaning, that indicate an uprooted tree and a broken tree-trunk, respectively (Oldeman 1990). The availability of snags has been increased topping trees with dynamite, chainsaw, girdling the canopy and with fungal inoculation (Bull et al. 1981, Lewis 1998, Campanaro et al. 2007). Models allow the simulation of CWD volumes per decay class that should be found in a stand as a function of its age (Ranius et al. 2003). Mortality and decomposition should be taken into account as well, since spatio-temporal dynamics are fundamental to understand the functionality and sustainability of deadwood compartment management (Morrissey et al. 2014). Additional factors to be considered include the stability and longevity of the structures, and how to optimize their spatial arrangement in order to reduce safety risks and do not affect forestry operations (Bauhus et al. 2009). The development of deadwood thresholds at the landscape scale represents a future challenge in the preservation of meta-population dynamics (Müller & Bütler 2010).

#### Landscape approach

Patch-based conservation networks have been designed specifying a minimum number of required patches, having certain habitat conditions (e.g., size, age distribution, structural complexity, composition, deadwood availability and decay rate – Lindenmayer & Franklin 2003, Schulte et al. 2006). For Fennoscandian and Baltic countries, identified criteria fell in three categories: (i) stand features (e.g., soil, slope, age, structure, and deadwood); (ii) habitat elements (e.g., habitat trees, boulders, streams, high stumps, pollard trees); and (iii) indicator species (i.e., specialized saproxylics – Timonen et al. 2010). The amount of retained habitat within a productive forest should

be above 5-10%, and possibly significantly higher, to achieve an ecological enrichment (Gustafsson et al. 2012). Managers should be aware that the benefits of these permanent small reserves tend to be tangible in the long-term (i.e., there could be a substantial lag time), and that these areas are sensitive to edge-effects and their effectiveness can be threatened by the surrounding harvest (Mönnkönen et al. 2011). The performance of the potential IdS can be evaluated maximizing the ecological value and minimizing their cost (Cateau et al. 2013). Concerning the area effect, Lindenmayer et al. (2015) performed the first experimental evaluation of the SLOSS (single large or several small areas for conservation) debate to VRH systems, and their results showed no significant area effect on bird species richness: thus IdS area could be adapted to local socio-economic requirements without compromising their value. Moreover, considering the extinction risk of saproxylic species, Ranius & Kindvall (2006) suggested that small reserves in landscapes characterized by commercial forestry have the advantage of containing higher quality fragments, which do not suffer the time-lag in deadwood restoration, as much as larger and younger areas.

The application of predictive forest models to establish the IdS minimum size in beech (*Fagus sylvatica* L.) forests proved that as little as 0.5 ha could ensure a high probability of continuous deadwood availability (Jakoby et al. 2010). The ecological connectivity of the IdS should be ensured by the presence of DC, made of veteran trees prone to develop microhabitats and to produce deadwood in its whole range of forms (e.g., position and decay stage). These two elements together constitute a network, continuous in both space and time, ensuring the preservation and dissemination of saproxylic species. The species' life-history traits, and in particular their mobility, could be considered to best define the balance between number and size of the IdS, thus between the life-boating function (smaller and more numerous IdSs) and habitat quality (larger, fewer and less affected by edge effect IdSs – Baker et al. 2013).

The management system offered by the FBA ensures habitat continuity and the presence of a more complete set of eco-units, to preserve the saproxylic species that occur either as habitat-tracking meta-populations (Schroeder et al. 2007) or as classical meta-population (Ranius 2000), according to their phylogenetic adaptation (Stokland et al. 2012). Further benefits of managing forests at the landscape scale include the combination of retention forestry with other land uses (Gustafsson et al. 2012) and *vice versa*, the integration of conservation actions in the existing land use practices (Hansbauer et al. 2010, Zapponi et al. 2013). The combination of statistical and remote sensing methods, such as

LIDAR, could allow the calibration of forest harvesting on local stochastic natural dynamics, overcoming artificial stand constraints. In addition, tools that allow the study of spatial patterns, like FRAGSTATS® (McGarigal et al. 2002), can increase the sustainability of forest products including spatio-temporal land cover changes (Radiogullari et al. 2014). Lastly, the experimental application of landscape retention forestry would allow the establishment of operative thresholds that could be used to reach specific conservation goals, according to disturbance regimes and site heterogeneity and composition (Rosenvald & Löhmus 2008). Combining different measures optimized according to site features and cost-effectiveness of interventions, can increase the feasibility of conservation strategies (Jonsson et al. 2006). Conservation planning software like MARXAN (Watts et al. 2009) offers further instruments to effectively allocate conservation efforts.

#### Bottom-up conservation strategies and certification

Nature-based silviculture relies on a thorough understanding of socio-economic and ecological components (Ciancio 2014), hence the collaboration and support of different stakeholders is strongly desirable (Boncina 2011). The effectiveness of landscape approaches, in both addressing the effect of growing anthropogenic pressures and finding ways to support the development of desirable conditions, has gained an extensive support (Sayer et al. 2013). Silvicultural practices integrating natural disturbances are especially suitable for public lands, where the management is generally not focused on timber production and/or traditional practices (O'Hara & Ramage 2013). On the other hand, since the conservation approaches associated with it (e.g., retention forestry) tend to be bottom-up, owners and license holders are those who generally bear the economic costs (Gustafsson et al. 2012). New policy frameworks should be developed to allow forest landscape planning through the incorporation of opportunity costs, to support landholders in the establishment of an “eco-eco” (economically and ecologically) viable productive forests. Examples of certifications for marketable products include labels that target species (e.g., bird-friendly shade-grown coffee), and their success depends on the consumers awareness of the biodiversity benefits and conversely on their willingness to pay for the additional value of the products (Pagiola & Ruthenberg 2002). To achieve the non-static landscape management, incentives and structures that coordinate the different stakeholders are essential (Bengtsson et al. 2003).

#### Conclusions

A heterogeneous range of substrate quality and management practices are essential to preserve saproxylic beetle diversity in productive forests (McGeoch et al.

2007). Preserving the functionality of forest ecosystems at the landscape-scale mitigates the detrimental effects of climate change (Sharma et al. 2013), and contributes to the connectivity, critical for protected area networks (Milad et al. 2011). Landscape approaches offer a widely accepted framework to deal with sustainable resource management and the growing pressures that threaten water, land and forests (Sayer et al. 2013). The implementation of a holistic system requires the combined effort of forestry technicians, conservation biologists and spatial statisticians, to understand disturbance regimes and design biodiversity networks. Quantitative approaches have long been applied in silviculture, providing fundamental instruments, unavailable in other environments, to establish conservation measures (Newton 2007). New standards for sustainable forest management are spreading, such as the certifications of the Forest Stewardship Council (FSC 2015) and the Program for the Endorsement of Forest Certification (PEFC 2010), that could feasibly take into account the key features of FBA. These planning instruments, considered together, would feasibly meet the critical targets identified by Lindenmayer & Franklin (2002): supporting species populations, allowing the movement of organisms, maintaining the reciprocal buffering between productive matrix and protected areas and, at the same time, ensuring production, commodities and services.

Since the expansion of protected areas is limited by competing socio-economic goals, the shift towards the integration of reserves and structural legacies in production and protection forests (Krumm et al. 2013) represents a promising approach. Recognizing that for biodiversity preservation in productive forests the benefits offered by the micro-reserves can exceed those of larger isolated reserves (Gustafsson et al. 2012), we believe that the FBA and similar approaches represent a cost-effective tool to work towards a truly multi-functional forestry.

## Acknowledgements

Franco Mason was supported by the MiPAAF, Ministry of Agricultural Policy and Forestry, Italian National Forest Service, Central Biodiversity Office, Rome, in the framework of the LIFE Project ManFor C.BD. (LIFE09 ENV/IT/000078); Livia Zapponi was supported by the LIFE Project ManFor C.BD. We would like to thank our colleagues Serena Corezzola and Davide Badano, who kindly read an earlier version of the manuscript. We would like to thank Randy Rollins for language revision, and Zarina Dalla Santa Brown for the final reviewing of grammar and style. We thank Giorgio Matteucci for his constant support in the framework of the Life Project ManFor C.BD., Alessandro Bottacci, Marco Pannella and Gianni Zanoni (Central Biodiversity Office of the Italian National Forest

Service), Paola Favero (local Biodiversity Office of Vittorio Veneto, Italy) for the keen interest in applying for the first time in Italy the *flot de senescence* concept in the Natural Biogenetic Reserve “Campo di Mezzo-Pian Parrocchia”, as an innovative action in the framework of the LIFE Project ManFor C.BD. Lastly, we thank Paolo Cantiani, Fabrizio Ferretti and Umberto Di Salvatore (CREA, Council for Agricultural Research and Economics – Arezzo, Italy) for the constructive discussions on the proposed method.

## References

- Abrahamsson M, Lindbladh M (2006). A comparison of saproxylic beetle occurrence between man-made high-and low-stumps of spruce (*Picea abies*). *Forest Ecology and Management* 226: 230-237. - doi: [10.1016/j.foreco.2006.01.046](https://doi.org/10.1016/j.foreco.2006.01.046)
- Aerts R (2013). Old trees: extraction, conservation can coexist. *Science* 339: 904-904. - doi: [10.1126/science.339.6122.904-a](https://doi.org/10.1126/science.339.6122.904-a)
- Angelstam PK (1998). Maintaining and restoring biodiversity in European boreal forests by developing natural disturbance regimes. *Journal of Vegetation Science* 9: 593-602. - doi: [10.2307/3237275](https://doi.org/10.2307/3237275)
- Arpin P, Ponge J-F, Faille A, Blandin P (1998). Diversity and dynamics of eco-units in the biological reserves of the Fontainebleau Forest (France): contribution of soil biology to a functional approach. *European Journal of Soil Biology* 34: 167-177. - doi: [10.1016/S1164-5563\(00\)86659-5](https://doi.org/10.1016/S1164-5563(00)86659-5)
- Ausden M (2007). *Habitat management for conservation. A handbook of techniques*. Oxford University Press, Oxford, UK, pp. 420.
- Baker SC, Spies TA, Wardlaw TJ, Balmer J, Franklin JF, Jordan GJ (2013). The harvested side of edges: effect of retained forests on the re-establishment of biodiversity in adjacent harvested areas. *Forest Ecology and Management* 302: 107-121. - doi: [10.1016/j.foreco.2013.03.024](https://doi.org/10.1016/j.foreco.2013.03.024)
- Bauhus J, Puettmann K, Messier C (2009). Silviculture for old-growth attributes. *Forest Ecology and Management* 258: 525-537. - doi: [10.1016/j.foreco.2009.01.053](https://doi.org/10.1016/j.foreco.2009.01.053)
- Bengtsson J, Angelstam P, Elmqvist T, Emanuelsson U, Folke C, Ihse M, Moberg F, Nyström M (2003). Reserves, resilience and dynamic landscapes. *AMBIO* 32: 389-396. - doi: [10.1579/0044-7447-32.6.389](https://doi.org/10.1579/0044-7447-32.6.389)
- Bergeron Y, Harvey B, Leduc A, Gauthier S (1999). Forest management guidelines based on natural disturbance dynamics: stand-and forest-level considerations. *The Forestry Chronicle* 75: 49-54. - doi: [10.5558/tfc75049-1](https://doi.org/10.5558/tfc75049-1)
- Bergman KO, Jansson N, Claesson K, Palmer MW, Milberg P (2012). How much and at what scale? Multiscale analyses as decision support for conservation of saproxylic oak beetles. *Forest Ecology and Management* 265: 133-141. - doi: [10.1016/j.foreco.2011.10.030](https://doi.org/10.1016/j.foreco.2011.10.030)
- Beudert B, Bässler C, Thorn S, Noss R, Schröder B, Dieffenbach-Fries H, Foulois N, Müller J (2015). Bark beetles increase biodiversity while maintaining drinking water quality. *Conservation Letters* 8 (4): 272-281. - doi: [10.1111/conl.12153](https://doi.org/10.1111/conl.12153)
- Bobiec A, Van Der Burgt H, Meijer K, Zuyderduyn

- C, Haga J, Vlaanderen B (2000). Rich deciduous forests in Bialowieza as a dynamic mosaic of developmental phases: premises for nature conservation and restoration management. *Forest Ecology and Management* 130: 159-175. - doi: [10.1016/S0378-1127\(99\)00181-4](https://doi.org/10.1016/S0378-1127(99)00181-4)
- Bollmann K, Braunisch V (2013). To integrate or to segregate: balancing commodity production and biodiversity conservation in European forests. In: “Integrative approaches as an opportunity for the conservation of forest biodiversity” (Kraus D, Krumm F eds). *European Forest Institute, Freiburg, Germany*, pp. 18-31. [online] URL: [http://cb.iew.unibe.ch/unibe/cb/content/e399625/Bollmann\\_BS2013.pdf](http://cb.iew.unibe.ch/unibe/cb/content/e399625/Bollmann_BS2013.pdf)
- Boncina A (2011). Conceptual approaches to integrate nature conservation into forest management: a Central European perspective. *International Forestry Review* 13: 13-22. - doi: [10.1505/146554811798201152](https://doi.org/10.1505/146554811798201152)
- Bormann FH, Likens GE (1994). *Pattern and process in a forested ecosystem*. Springer-Verlag, New York, USA, pp. 251.
- Boudru M (1989). *Forêt et sylviculture. Traitement des forêts*. [Forest and silviculture. Forest systems]. Les Presses Agronomiques, Gembloux, Belgium, pp. 356. [in French]
- Bouget C, Duelli P (2004). The effects of wind-throw on forest insect communities: a literature review. *Biological Conservation* 118: 281-299. - doi: [10.1016/j.biocon.2003.09.009](https://doi.org/10.1016/j.biocon.2003.09.009)
- Bouget C, Lassauce A, Jonsell M (2012). Effects of fuelwood harvesting on biodiversity - a review focused on the situation in Europe. *Canadian Journal of Forest Research* 42: 1421-1432. - doi: [10.1139/x2012-078](https://doi.org/10.1139/x2012-078)
- Branquart E, Verheyen K, Latham J (2008). Selection criteria of protected forest areas in Europe: the theory and the real world. *Biological Conservation* 141: 2795-2806. - doi: [10.1016/j.biocon.2008.08.015](https://doi.org/10.1016/j.biocon.2008.08.015)
- Brin A, Bouget C, Valladares L, Brustel H (2012). Are stumps important for the conservation of saproxylic beetles in managed forests? - Insights from a comparison of assemblages on logs and stumps in oak-dominated forests and pine plantations. *Insect Conservation and Diversity* 6: 255-264. - doi: [10.1111/j.1752-4598.2012.00209.x](https://doi.org/10.1111/j.1752-4598.2012.00209.x)
- Brockerhoff EG, Jactel H, Parrotta JA, Quine CP, Sayer JA (2008). Plantation forests and biodiversity: oxymoron or opportunity? *Biodiversity and Conservation* 17: 925-951. - doi: [10.1007/s10531-008-9380-x](https://doi.org/10.1007/s10531-008-9380-x)
- Brumelis G, Jonsson BG, Kouki J, Kuuluvainen T, Shorohova E (2011). Forest naturalness in northern Europe: perspectives on processes, structures and species diversity. *Silva Fennica* 45: 807-821. - doi: [10.14214/sf.446](https://doi.org/10.14214/sf.446)
- Bull E, Partridge W, Williams W (1981). *Creating snags with explosives*. Pacific Northwest Forest and Range Experiment Station, USDA Forest Service, Portland, OR, USA, pp. 4. [online] URL: <http://books.google.com/books?id=WFq1WzMUdqG>
- Campanaro A, Hardersen S, Mason F (2007). Piano di gestione della Riserva Naturale Statale e Sito Natura 2000 “Bosco della Fontana” [Management Plan of the State Natural Reserve and Natura 2000 Site “Bosco della Fontana”]. Cierre Edizioni, Verona, Italy, pp. 223. [in



- Italian]
- Cantero A, Passola G, Aragón A, De Francisco M, Mugarza V, Riaño P (2014). Apuntes sobre trasmochos. Guía práctica para el trasmocheo [Notes on pollards. Best practices' guide for pollarding]. Gipuzkoako Foru Aldundia, Fundación HAZI Fundazioa, Donostia, Spain, pp. 90. [in Spanish]
- Cappelli M (1988). Selvicoltura generale [General forestry]. Edagricole, Bologna, Italy, pp. 298. [in Italian]
- Cateau E, Parrot M, Reyna K, Roux A, Rossi M, Bruciamacchie M, Vallauri D (2013). Réseau d'îlots de vieux bois. Eléments de méthode et test dans les forêts publiques du Mont-Ventoux [Network of deadwood islands. Methodological approach and application to the Mont-Ventoux public forests]. In: "Naturalité vers un autre culture des eaux et des forêts". WWF, France, pp. 66. [in French]
- Cavalli R, Mason F (2003). Techniques for reestablishment of dead wood for saproxylic fauna conservation. LIFE Nature project NAT/IT/99/6245 "Bosco della Fontana" (Mantova, Italy). Scientific Reports 2, Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale di Verona - Bosco della Fontana. Gianluigi Arcari Editore, Mantova, Italy, pp. 112.
- Cawthen L, Munks S (2011). The use of hollow-bearing trees retained in multi-aged regenerating production forest by the Tasmanian common brushtail possum (*Trichosurus vulpecula fuliginosus*). *Wildlife Research* 38: 687-695. - doi: [10.1071/WR10125](https://doi.org/10.1071/WR10125)
- Ciancio O (2014). Storia del pensiero forestale. Selvicoltura, filosofia, etica [History of forestry thought. Forestry, philosophy, ethics]. Rubbettino Editore, Soveria Mannelli, Italy, pp. 560. [in Italian]
- Ciancio O, Nocentini S (2005). Biodiversity conservation in Mediterranean forest ecosystems: from theory to operationality. In: "Monitoring and indicators of forest biodiversity in Europe. From ideas to operationality" (Marchetti M ed). European Forest Institute, Joensuu, Finland, pp. 163-168. [online] URL: <http://www.iufro.org/download/file/4772/4508/>
- Coates KD, Burton PJ (1997). A gap-based approach for development of silvicultural systems to address ecosystem management objectives. *Forest Ecology and Management* 99: 337-354. - doi: [10.1016/S0378-1127\(97\)00113-8](https://doi.org/10.1016/S0378-1127(97)00113-8)
- Craig MD, Benkovic AM, Grigg AH, Hardy GESJ, Fleming PA, Hobbs RJ (2011). How many mature microhabitats does a slow-recolonising reptile require? Implications for restoration of bauxite minesites in south-western Australia. *Australian Journal of Zoology* 59: 9-17. - doi: [10.1071/ZO10046](https://doi.org/10.1071/ZO10046)
- Croak BM, Crowther MS, Webb JK, Shine R (2013). Movements and habitat use of an endangered snake, *Hoplocephalus bungaroides* (Elapidae): implications for Conservation. *PLoS One* 8 (4): e61711. - doi: [10.1371/journal.pone.0061711](https://doi.org/10.1371/journal.pone.0061711)
- Del Favero R, Poldini L, Bortoli P, Lasen C, Drossi G, Vanone G (1998). La vegetazione forestale e la selvicoltura nella regione Friuli-Venezia Giulia [Forest vegetation and silviculture in the region Friuli-Venezia Giulia]. Regione Autonoma Friuli Venezia Giulia, Udine, Italy, pp. 490. [in Italian]
- De Philippis A (1948). Su alcune forme di trattamento delle fustaie [Types of high forest treatment]. *L'Italia Forestale e Montana* 3: 3-10. [in Italian]
- Diaz S, Kitzberger T (2013). Nest habitat selection by the Austral parakeet in north-western Patagonia. *Austral Ecology* 38: 268-278. - doi: [10.1111/j.1442-9993.2012.02400.x](https://doi.org/10.1111/j.1442-9993.2012.02400.x)
- Donato DC, Campbell JL, Franklin JF (2012). Multiple successional pathways and precocity in forest development: can some forests be born complex? *Journal of Vegetation Science* 23: 576-584. - doi: [10.1111/j.1654-1103.2011.01362.x](https://doi.org/10.1111/j.1654-1103.2011.01362.x)
- Dubois GF, Vignon V, Delettre YR, Rantier Y, Vernon P, Burel F (2009). Factors affecting the occurrence of the endangered saproxylic beetle *Osmoderma eremita* (Scopoli, 1763) (Coleoptera: Cetoniidae) in an agricultural landscape. *Landscape and Urban Planning* 91: 152-159. - doi: [10.1016/j.landurbplan.2008.12.009](https://doi.org/10.1016/j.landurbplan.2008.12.009)
- Edwards DP, Gilroy JJ, Woodcock P, Edwards FA, Larsen TH, Andrews DJ, Derhé MA, Docherty TD, Hsu WW, Mitchell SL (2014). Land-sharing versus land-sparing logging: reconciling timber extraction with biodiversity conservation. *Global Change Biology* 20: 183-191. - doi: [10.1111/gcb.12353](https://doi.org/10.1111/gcb.12353)
- Fan Z, Shifley SR, Thompson III FR, Larsen DR (2004). Simulated cavity tree dynamics under alternative timber harvest regimes. *Forest Ecology and Management* 193: 399-412. - doi: [10.1016/j.foreco.2004.02.008](https://doi.org/10.1016/j.foreco.2004.02.008)
- Fenton NJ, Imbeau L, Work T, Jacobs J, Bescond H, Drapeau P, Bergeron Y (2013). Lessons learned from 12 years of ecological research on partial cuts in black spruce forests of northwestern Quebec. *Forestry Chronicle* 89: 350-359.
- FOREST-EUROPE/UNECE/FAO (2011). State of Europe's forests 2011. Status and trends in sustainable forest management in Europe. Ministerial Conference on the Protection of Forests in Europe, Forest Europe Liaison Unit Oslo, Norway, pp. 344.
- Franklin JF, Berg DR, Thornburgh DA, Tappeiner JC (1997). Alternative silvicultural approaches to timber harvesting: variable retention harvest systems. In: "Creating a forestry for the 21<sup>st</sup> century: the science of ecosystem management" (Kohm KA, Franklin JF eds). Island Press, Washington, DC, USA, pp. 111-139.
- Franklin JF, Lindenmayer D, MacMahon JA, McKee A, Magnuson J, Perry DA, Waide R, Foster D (2000). Threads of Continuity. Conservation in Practice 1: 8-17. - doi: [10.1111/j.1526-4629.2000.tb00155.x](https://doi.org/10.1111/j.1526-4629.2000.tb00155.x)
- Fry R, Lonsdale D (1991). Habitat conservation for insects: a neglected green issue. *Amateur Entomologists' Society*, Middlesex, UK, pp. 262. [online] URL: <http://library.wur.nl/WebQuery/cl/520073>
- FSC (2015). FSC principles and criteria for forest stewardship. FSC-STD-01-001 V5-2 EN, Forest Stewardship Council, Bonn, Germany, pp. 32.
- Gardner T (2010). Monitoring forest biodiversity. Improving conservation through ecologically responsible management. Earthscan, London, UK, pp. 652. [online] URL: <http://www.publish.csiro.au/nid/201/pid/6543.htm>
- Gendreau-Berthiaume B, Kneeshaw DD, Harvey BD (2012). Effects of partial cutting and partial disturbance by wind and insects on stand composition, structure and growth in boreal mixedwoods. *Forestry* 85: 551-565. - doi: [10.1093/forestry/cps051](https://doi.org/10.1093/forestry/cps051)
- Gibbons P, Lindenmayer DB (2002). Tree hollows and wildlife conservation in Australia. CSIRO, Collingwood, Australia, pp. 211. [online] URL: [http://books.google.com/books?id=6Q28RllsW\\_IC](http://books.google.com/books?id=6Q28RllsW_IC)
- Giulietti V, Ferretti F, Pelleri F (2009). Prove di diradamento in acero-frassinetti di neoformazione nella Comunità Montana Agno-Chiampo (VI): risultati dopo il secondo intervento [Thinning trials in secondary growth ash-maple stands carried out in the Agno-Chiampo district (VI): results after second thinning.]. *Annali CRA-Centro Ricerca per la Selvicoltura, Arezzo* 35: 2007-2008. [in Italian]
- Gossner M, Engel K, Ammer U (2006). Effects of selection felling and gap felling on forest arthropod communities: a case study in a spruce-beech stand of southern Bavaria. *European Journal of Forest Research* 125: 345-360. - doi: [10.1007/s10342-006-0126-6](https://doi.org/10.1007/s10342-006-0126-6)
- Gossner MM, Lachat T, Brunet J, Isacson G, Bouget C, Brustel H, Brandl R, Weisser WW, Mueller J (2013). Current near-to-nature forest management effects on functional trait composition of saproxylic beetles in beech forests. *Conservation Biology* 27: 605-614. - doi: [10.1111/cobi.12023](https://doi.org/10.1111/cobi.12023)
- Götmark F, Thorell M (2003). Size of nature reserves: densities of large trees and dead wood indicate high value of small conservation forests in southern Sweden. *Biodiversity and Conservation* 12: 1271-1285. - doi: [10.1023/A:1023000224642](https://doi.org/10.1023/A:1023000224642)
- Grove SJ (2002). Saproxylic insect ecology and the sustainable management of forests. *Annual Review of Ecology and Systematics* 33: 1-23. - doi: [10.1146/annurev.ecolsys.33.010802.150507](https://doi.org/10.1146/annurev.ecolsys.33.010802.150507)
- Gustafsson L, Baker SC, Bauhus J, Beese WJ, Brodie A, Kouki J, Lindenmayer DB, Löhmus A, Pastur GM, Messier C, Neyland M, Palik B, Sverdrup-Thygeson A, Volney WJA, Wayne A, Franklin JF (2012). Retention forestry to maintain multi-functional forests: a world perspective. *Bioscience* 62: 633-645. - doi: [10.1525/bio.2012.62.7.6](https://doi.org/10.1525/bio.2012.62.7.6)
- Hansbauer MM, Storch I, Knauer F, Pilz S, Kuchenhoff H, Vegvari Z, Pimentel RG, Metzger JP (2010). Landscape perception by forest understory birds in the Atlantic Rainforest: black-and-white versus shades of grey. *Landscape Ecology* 25: 407-417. - doi: [10.1007/s10980-009-9418-9](https://doi.org/10.1007/s10980-009-9418-9)
- Harmon ME (2001). Moving towards a new paradigm for woody detritus management. *Ecological Bulletins* 49: 269-278. [online] URL: <http://www.jstor.org/stable/20113283>
- Hedgren PO (2007). Early arriving saproxylic beetles (Coleoptera) and parasitoids (Hymenoptera) in low and high stumps of Norway spruce. *Forest Ecology and Management* 241: 155-161. - doi: [10.1016/j.foreco.2007.01.020](https://doi.org/10.1016/j.foreco.2007.01.020)
- Heikkala O, Suominen M, Junninen K, Hämäläinen A, Kouki J (2014). Effects of retention level and fire on retention tree dynamics in boreal forests. *Forest Ecology and Management* 328: 193-201. - doi: [10.1016/j.foreco.2014.05.022](https://doi.org/10.1016/j.foreco.2014.05.022)
- Horak J (2015). What is happening after an abiotic disturbance? Response of saproxylic beetles in the Primorsky Region woodlands (Far East-

- ern Russia). *Journal of Insect Conservation* 19: 97-103. - doi: [10.1007/s10841-014-9750-5](https://doi.org/10.1007/s10841-014-9750-5)
- Hyvärinen E, Kouki J, Martikainen P (2009). Prescribed fires and retention trees help to conserve beetle diversity in managed boreal forests despite their transient negative effects on some beetle groups. *Insect Conservation and Diversity* 2: 93-105. - doi: [10.1111/j.1752-4598.2009.00048.x](https://doi.org/10.1111/j.1752-4598.2009.00048.x)
- Jakoby O, Rademacher C, Grimm V (2010). Modelling dead wood islands in European beech forests: how much and how reliably would they provide dead wood? *European Journal of Forest Research* 129: 659-668. - doi: [10.1007/s10342-010-0366-3](https://doi.org/10.1007/s10342-010-0366-3)
- Jonsson BG, Krusys N, Ranius T (2005). Ecology of species living on dead wood-lessons for dead wood management. *Silva Fennica* 39: 289-309. - doi: [10.14214/sf.390](https://doi.org/10.14214/sf.390)
- Jonsson BG, Siitonen J (2012). Dead wood and sustainable forest management. In: "Biodiversity in dead wood" (Stokland J, Sittonen J, Jonsson B eds). Cambridge University Press, Cambridge, UK, pp. 302-337.
- Jonsson M, Ranius T, Ekvall H, Bostedt G, Dahlberg A, Ehnström B, Nordén B, Stokland JN (2006). Cost-effectiveness of silvicultural measures to increase substrate availability for red-listed wood-living organisms in Norway spruce forests. *Biological Conservation* 127: 443-462. - doi: [10.1016/j.biocon.2005.09.004](https://doi.org/10.1016/j.biocon.2005.09.004)
- Kadiogullari A, Sayin M, Çelîk D, Borucu S, Çil B, Bulut S (2014). Analysing land cover changes for understanding of forest dynamics using temporal forest management plans. *Environmental Monitoring and Assessment* 186: 2089-2110. - doi: [10.1007/s10661-013-3520-9](https://doi.org/10.1007/s10661-013-3520-9)
- Kaplan JO, Krumhardt KM, Zimmermann N (2009). The prehistoric and preindustrial deforestation of Europe. *Quaternary Science Reviews* 28: 3016-3034. - doi: [10.1016/j.quascirev.2009.09.028](https://doi.org/10.1016/j.quascirev.2009.09.028)
- Kern CC, Montgomery RA, Reich PB, Strong TF (2014). Harvest-created canopy gaps increase species and functional trait diversity of the forest ground-layer community. *Forest Science* 60: 335-344. - doi: [10.5849/forsci.13-015](https://doi.org/10.5849/forsci.13-015)
- Koch Widerberg M, Ranius T, Drobyshev I, Nilsson U, Lindbladh M (2012). Increased openness around retained oaks increases species richness of saproxylic beetles. *Biodiversity and Conservation* 21: 3035-3059. - doi: [10.1007/s10531-012-0353-8](https://doi.org/10.1007/s10531-012-0353-8)
- Kohm K, Franklin J (1997). *Creating a forestry for the 21<sup>st</sup> century: the science of ecosystem management*. Island Press, Washington, DC, USA, pp. 491. [online] URL: [http://books.google.com/books?id=XNiq\\_KJ-zpgC](http://books.google.com/books?id=XNiq_KJ-zpgC)
- Komonen A, Halme P, Jäntti M, Koskela T, Kotiaho JS, Toivanen T (2014). Created substrates do not fully mimic natural substrates in restoration: the occurrence of polypores on spruce logs. *Silva Fennica* 48: 1-12. - doi: [10.14214/sf.980](https://doi.org/10.14214/sf.980)
- Krumm F, Schuck A, Kraus D (2013). Integrative management approaches: a synthesis. In: "Integrative approaches as an opportunity for the conservation of forest biodiversity" (Kraus D, Krumm F eds). European Forest Institute, Freiburg, Germany, pp. 255-261. [online] URL: [http://www.researchgate.net/profile/Anne\\_Ox\\_brough/publication/258603988](http://www.researchgate.net/profile/Anne_Ox_brough/publication/258603988)
- Kuuluvainen T (2002). Natural variability of forests as a reference for restoring and managing biological diversity in boreal Fennoscandia. *Silva Fennica* 36: 97-125. [online] URL: <http://m.metla.eu/silvafennica/full/sf36/sf361097.pdf>
- Kuuluvainen T, Tukia H, Aapala K, (2004). Ecological restoration of forested ecosystems in Finland. In: "Restoration of boreal and temperate forests" (Stanturf J, Madsen P eds). CRC Press, Boca Raton, FL, USA, pp. 285-298.
- Lachat T, Büttler R (2007). Gestion des vieux arbres et du bois mort: îlots de sénescence, arbres-habitat et métapopulations saproxyliques [Veteran trees and deadwood management: senescence islands, habitat trees and saproxylic metacommunities]. Mandat de l'Office fédéral de l'environnement, OFEV, WSL, Lausanne, Switzerland, pp. 2. [in French] [online] URL: [http://www.slf.ch/fe/walddynamik/diversitaet/totholzmanagement/resume\\_rapport.pdf](http://www.slf.ch/fe/walddynamik/diversitaet/totholzmanagement/resume_rapport.pdf)
- Lachat T, Büttler R (2008). Îlots de sénescence et arbres-habitat pour augmenter la biodiversité en forêt [Senescence islands and habitat trees to increase forest biodiversity]. *La Forêt* 6: 20-21. [in French]
- Lafond V, Lagarrigues G, Cordonnier T, Courbaud B (2014). Uneven-aged management options to promote forest resilience for climate change adaptation: effects of group selection and harvesting intensity. *Annals of Forest Science* 71: 173-186. - doi: [10.1007/s13595-013-0291-y](https://doi.org/10.1007/s13595-013-0291-y)
- Larrieu L, Cabanettes A (2012). Species, life status, and diameter are important tree features for diversity and abundance of tree microhabitats in subnatural montane beech-fir forests. *Canadian Journal of Forest Research* 42: 1433-1445. - doi: [10.1139/x2012-077](https://doi.org/10.1139/x2012-077)
- Larrieu L, Cabanettes A, Brin A, Bouget C, Deconchat M (2014a). Tree microhabitats at the stand scale in montane beech-fir forests: practical information for taxa conservation in forestry. *European Journal of Forest Research* 133: 355-367. - doi: [10.1007/s10342-013-0767-1](https://doi.org/10.1007/s10342-013-0767-1)
- Larrieu L, Cabanettes A, Delarue A (2012). Impact of silviculture on dead wood and on the distribution and frequency of tree microhabitats in montane beech-fir forests of the Pyrenees. *European Journal of Forest Research* 131: 773-786. - doi: [10.1007/s10342-011-0551-z](https://doi.org/10.1007/s10342-011-0551-z)
- Larrieu L, Cabanettes A, Gonin P, Lachat T, Paillet Y, Winter S, Bouget C, Deconchat M (2014b). Deadwood and tree microhabitat dynamics in unharvested temperate mountain mixed forests: a life-cycle approach to biodiversity monitoring. *Forest Ecology and Management* 334: 163-173. - doi: [10.1016/j.foreco.2014.09.007](https://doi.org/10.1016/j.foreco.2014.09.007)
- Lassauce A, Anselle P, Lieutier F, Bouget C (2012). Coppice-with-standards with an overmature coppice component enhance saproxylic beetle biodiversity: a case study in French deciduous forests. *Forest Ecology and Management* 266: 273-285. - doi: [10.1016/j.foreco.2011.11.016](https://doi.org/10.1016/j.foreco.2011.11.016)
- Leroux SJ, Rayfield B (2014). Methods and tools for addressing natural disturbance dynamics in conservation planning for wilderness areas. *Diversity and Distributions* 20: 258-271. - doi: [10.1111/ddi.12155](https://doi.org/10.1111/ddi.12155)
- Lewis JC (1998). *Creating snags and wildlife trees in commercial forest landscapes*. Western Journal of Applied Forestry 13: 97-101. [online] URL: <http://www.ingentaconnect.com/content/saf/wjaf/1998/00000013/00000003/arto0006>
- Lindenmayer DB, Franklin JF (2002). *Conserving forest biodiversity: a comprehensive multi-scaled approach*. Island Press, Washington, DC, USA, pp. 351. [online] URL: <http://books.google.com/books?id=ODEFUM2WtAYC>
- Lindenmayer DB, Franklin JF (2003). *Towards forest sustainability*. CSIRO Publishing, Collingwood, Australia, pp. 244. [online] URL: <http://books.google.com/books?id=RVzMtnDAiuoC>
- Lindenmayer DB, Laurance WF (2012). *A history of hubris - cautionary lessons in ecologically sustainable forest management*. *Biological Conservation* 151: 11-16. - doi: [10.1016/j.biocon.2011.10.032](https://doi.org/10.1016/j.biocon.2011.10.032)
- Lindenmayer DB, Wood J, McBurney L, Blair D, Banks SC (2015). Single large versus several small: the SLOSS debate in the context of bird responses to a variable retention logging experiment. *Forest Ecology and Management* 339: 1-10. - doi: [10.1016/j.foreco.2014.11.027](https://doi.org/10.1016/j.foreco.2014.11.027)
- Mallik AU, Kreutzweiser DP, Spalvieri CM (2014). Forest regeneration in gaps seven years after partial harvesting in riparian buffers of boreal mixedwood streams. *Forest Ecology and Management* 312: 117-128. - doi: [10.1016/j.foreco.2013.10.015](https://doi.org/10.1016/j.foreco.2013.10.015)
- Man R, Rice JA, MacDonald GB (2010). Five-year light, vegetation, and regeneration dynamics of boreal mixedwoods following silvicultural treatments to establish productive aspen-spruce mixtures in northeastern Ontario. *Canadian Journal of Forest Research* 40: 1529-1541. - doi: [10.1139/X10-088](https://doi.org/10.1139/X10-088)
- Manning AD, Gibbons P, Fischer J, Oliver DL, Lindenmayer DB (2013). Hollow futures? Tree decline, lag effects and hollow-dependent species. *Animal Conservation* 16: 395-403. - doi: [10.1111/acv.12006](https://doi.org/10.1111/acv.12006)
- Mason F (2002). *Problems of conservation and management*. In: "Woodlands of the Po Plain" (Ruffo S ed). Italian Ministry of the Environment and Territory Protection, Friuli Museum of Natural History, Udine, Italy, pp. 91-137.
- Mason F (2004). *Dinamica di una foresta della Pianura Padana*. Bosco della Fontana. Primo contributo, monitoraggio 1995. Seconda edizione con linee di gestione forestale [Dynamics of a forest of the Po Valley. Bosco della Fontana. First contribution, 1995 monitoring. Second edition with guidelines of forest management]. Arcari Editore, Mantova, Italy, pp. 224. [in Italian]
- Mason F, Nardi G, Tisato M (2003). *Dead wood: a key to biodiversity*. Sherwood 95, Suppl. 2, pp. 100. [online] URL: <http://docplayer.it/1548729>
- Mayer AL, Kauppi PE, Angelstam PK, Zhang Y, Tikka PM (2005). Importing timber, exporting ecological impact. *Science* 308: 359-360. - doi: [10.1126/science.1109476](https://doi.org/10.1126/science.1109476)
- McGarigal K, Cushman SA, Neel MC, Ene E (2002). *FRAGSTATS: spatial pattern analysis program for categorical maps*. Computer software program, University of Massachusetts, Amherst, MA, USA. [online] URL: <http://www.umass.edu/landeco/research/fragstats/fragstats.html>
- McGeoch MA, Schroeder M, Ekbohm B, Larsson S (2007). Saproxylic beetle diversity in a mana-

- ged boreal forest: importance of stand characteristics and forestry conservation measures. *Diversity and Distributions* 13: 418-429. - doi: [10.1111/j.1472-4642.2007.00350.x](https://doi.org/10.1111/j.1472-4642.2007.00350.x)
- MCPFE (2003). Improved pan-European indicators for sustainable forest management. Ministerial Conference on the Protection of Forests, Europe Liaison Unit, Vienna, Austria, pp. 6.
- Mercurio R (1999). Il taglio a buche: una forma colturale ancora valida? [Is small-patch clear-cutting an up-to-date silvicultural system?] *L'Italia Forestale e Montana* 1: 4-17. [in Italian]
- Mercurio R, Mallamaci C, Muscolo A, Sidari M (2009). Gap size effects on tree regeneration in afforestations of Black pine (*Pinus nigra* Arn.). *Forest@* 6 (1): 312-319. [in Italian with English summary] - doi: [10.3832/eforo591-006](https://doi.org/10.3832/eforo591-006)
- Matthews JD (1989). *Sylvicultural systems*. Clarendon Press, Oxford, UK, pp. 284.
- Michel AK, Winter S (2009). Tree microhabitat structures as indicators of biodiversity in Douglas-fir forests of different stand ages and management histories in the Pacific Northwest, USA. *Forest Ecology and Management* 257: 1453-1464. - doi: [10.1016/j.foreco.2008.11.027](https://doi.org/10.1016/j.foreco.2008.11.027)
- Miklín J, Čížek L (2014). Erasing a European biodiversity hot-spot: open woodlands, veteran trees and mature forests succumb to forestry intensification, succession, and logging in a UNESCO Biosphere Reserve. *Journal for Nature Conservation* 22: 35-41. - doi: [10.1016/j.jnc.2013.08.002](https://doi.org/10.1016/j.jnc.2013.08.002)
- Milad M, Schaich H, Burgi M, Konold W (2011). Climate change and nature conservation in Central European forests: a review of consequences, concepts and challenges. *Forest Ecology and Management* 261: 829-843. - doi: [10.1016/j.foreco.2010.10.038](https://doi.org/10.1016/j.foreco.2010.10.038)
- Mönkkönen M, Reunanen P, Kotiaho JS, Juutinen A, Tikkanen O-P, Kouki J (2011). Cost-effective strategies to conserve boreal forest biodiversity and long-term landscape-level maintenance of habitats. *European Journal of Forest Research* 130: 717-727. - doi: [10.1007/s10342-010-0461-5](https://doi.org/10.1007/s10342-010-0461-5)
- Morrissey RC, Jenkins MA, Saunders MR (2014). Accumulation and connectivity of coarse woody debris in partial harvest and unmanaged relict forests. *PloS One* 9(11): e113323. - doi: [10.1371/journal.pone.0113323](https://doi.org/10.1371/journal.pone.0113323)
- Müller J, Büttler R (2010). A review of habitat thresholds for dead wood: a baseline for management recommendations in European forests. *European Journal of Forest Research* 129: 981-992. - doi: [10.1007/s10342-010-0400-5](https://doi.org/10.1007/s10342-010-0400-5)
- Müller J, Jarzabek-Müller A, Bussler H, Gossner MM (2014). Hollow beech trees identified as keystone structures for saproxylic beetles by analyses of functional and phylogenetic diversity. *Animal Conservation* 17: 154-162. - doi: [10.1111/acv.12075](https://doi.org/10.1111/acv.12075)
- Newton AC (2007). *Forest ecology and conservation*. Oxford University Press, Oxford, UK, pp. 471.
- Nieuwenhuis M (2000). Terminology of forest management planning. Terms and definitions in English - equivalent terms in German, French, Spanish, Italian, Portuguese, Hungarian and Japanese. IUFRO 4.04.07 Silvaplan and Silvavoc, IUFRO World Series, vol. 9-en, Wien.
- Nocentini S (2011). The forest as a complex biological system: theoretical and practical consequences. *L'Italia Forestale e Montana* 66: 191-196. - doi: [10.4129/jfm.2011.3.02](https://doi.org/10.4129/jfm.2011.3.02)
- Nordén B, Götmark F, Tönning M, Ryberg M (2004). Dead wood in semi-natural temperate broadleaved woodland: contribution of coarse and fine dead wood, attached dead wood and stumps. *Forest Ecology and Management* 194: 235-248. - doi: [10.1016/j.foreco.2004.02.043](https://doi.org/10.1016/j.foreco.2004.02.043)
- O'Hara KL, Ramage BS (2013). Silviculture in an uncertain world: utilizing multi-aged management systems to integrate disturbance. *Forestry* 86: 401-410. - doi: [10.1093/forestry/cpt012](https://doi.org/10.1093/forestry/cpt012)
- Oldeman RAA (1990). *Forests: elements of silvology*. Springer-Verlag, Berlin, Germany, pp. 624.
- Otto H (1993). La foresta dinamica: basi ecologiche della selvicoltura naturalistica [The dynamic forest: ecological basis of naturalistic silviculture]. *Dendronatura* 14: 17-26. [in Italian]
- Pagiola S, Ruthenberg IM (2002). Selling biodiversity in a coffee cup: shade-grown coffee and conservation in Mesoamerica. In: "Selling forest environmental services" (Pagiola S, Bishop J, Landell-Mills N eds). Earthscan, London, UK, pp. 103-126. [online] URL: <http://books.google.com/books?id=fBuXqru8Co4C>
- Paquette A, Messier C (2009). The role of plantations in managing the world's forests in the Anthropocene. *Frontiers in Ecology and the Environment* 8: 27-34. - doi: [10.1890/080116](https://doi.org/10.1890/080116)
- Pastur GM, Esteban R, Cellini J, Lencinas M, Peri P, Neyland M (2014). Survival and growth of *Nothofagus pumilio* seedlings under several microenvironments after variable retention harvesting in southern Patagonian forests. *Annals of Forest Science* 71: 349-362. - doi: [10.1007/s13595-013-0343-3](https://doi.org/10.1007/s13595-013-0343-3)
- Pavari A (1956). Governo e trattamento dei boschi [Forest systems and harvest methods]. Reda, Roma, Italy, pp. 81. [in Italian]
- Pawson S, Brin A, Brockerhoff E, Lamb D, Payn T, Paquette A, Parrotta J (2013). Plantation forests, climate change and biodiversity. *Biodiversity and Conservation* 22: 1203-1227. - doi: [10.1007/s10531-013-0458-8](https://doi.org/10.1007/s10531-013-0458-8)
- PEFC (2010). Sustainable forest management - Requirements. PEFC International Standard, Requirements for certification schemes, PEFC ST 1003:2010, Geneva, Switzerland, pp. 16. [online] URL: <http://www.pefc.org/standards/technical-documentation/pefc-international-standards-2010/676>
- Perrin H (1954). *Selvicoltura* [Forestry]. Accademia Italiana di Scienze Forestali, Firenze, Italy, pp. 430. [in Italian]
- Peterken GF (1996). *Natural woodland: ecology and conservation in northern temperate regions*. Cambridge University Press, Cambridge, UK, pp. 522.
- Phalan B, Onial M, Balmford A, Green RE (2011). Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Science* 333: 1289-1291. - doi: [10.1126/science.1208742](https://doi.org/10.1126/science.1208742)
- Piussi P (1994). *Selvicoltura generale* [General forestry]. UTET, Torino, Italy, pp. 421. [in Italian]
- Ranius T (2000). Minimum viable metapopulation size of a beetle, *Osmoderma eremita*, living in tree hollows. *Animal Conservation* 3: 37-43. - doi: [10.1111/j.1469-1795.2000.tb00085.x](https://doi.org/10.1111/j.1469-1795.2000.tb00085.x)
- Ranius T, Kindvall O (2006). Extinction risk of wood-living model species in forest landscapes as related to forest history and conservation strategy. *Landscape Ecology* 21: 687-698. - doi: [10.1007/s10980-005-5222-3](https://doi.org/10.1007/s10980-005-5222-3)
- Ranius T, Kindvall O, Krus N, Jonsson BG (2003). Modelling dead wood in Norway spruce stands subject to different management regimes. *Forest Ecology and Management* 182: 13-29. - doi: [10.1016/S0378-1127\(03\)00027-6](https://doi.org/10.1016/S0378-1127(03)00027-6)
- Ranius T, Niklasson M, Berg N (2009). Development of tree hollows in pedunculate oak (*Quercus robur*). *Forest Ecology and Management* 257: 303-310. - doi: [10.1016/j.foreco.2008.09.007](https://doi.org/10.1016/j.foreco.2008.09.007)
- Rayner L, Ellis M, Taylor JE (2014). Hollow occurrence and abundance varies with tree characteristics and among species in temperate woodland *Eucalyptus*. *Austral Ecology* 39: 145-157. - doi: [10.1111/aec.12052](https://doi.org/10.1111/aec.12052)
- Read H (2000). *Veteran trees: a guide to good management*. English Nature, Peterborough, UK, pp. 92.
- Regnery B, Paillet Y, Couvet D, Kerbirou C (2013). Which factors influence the occurrence and density of tree microhabitats in Mediterranean oak forests? *Forest Ecology and Management* 295: 118-125. - doi: [10.1016/j.foreco.2013.01.009](https://doi.org/10.1016/j.foreco.2013.01.009)
- Richter A (1950). *Heinrich Cotta. Leben und Werk eines deutschen Forstmannes* [Heinrich Cotta - life and work of a German forester]. Raabeul-Berlin Neumann Verlag, Berlin, Germany, pp. 125-166. [in German]
- Roberge JM, Mönkkönen M, Toivanen T, Kotiaho JS (2013). Retention forestry and biodiversity conservation: a parallel with agroforestry. *Nature Conservation* 4: 29-33. - doi: [10.3897/natureconservation.4.5116](https://doi.org/10.3897/natureconservation.4.5116)
- Rojo Saiz F (1977). *Fotologia forestal y ordenación* [Forest photology and ordination]. *Montes* 2: 31-34. [in Spanish]
- Rosenvald R, Löhms A (2008). For what, when, and where is green-tree retention better than clear-cutting? A review of the biodiversity aspects. *Forest Ecology and Management* 255: 1-15. - doi: [10.1016/j.foreco.2007.09.016](https://doi.org/10.1016/j.foreco.2007.09.016)
- Roussel L (1972). *Photologie forestière* [Forest photology]. Masson et Cie, Paris, France pp. 136. [in French]
- Runkle JR (1991). Gap dynamics of old-growth eastern forests: management implications. *Natural Areas Journal* 11: 19-25.
- Sayer J, Sunderland T, Ghazoul J, Pfund J-L, Sheil D, Meijaard E, Venter M, Boedihartono AK, Day M, Garcia C (2013). Ten principles for a landscape approach to reconciling agriculture, conservation, and other competing land uses. *Proceedings of the National Academy of Sciences USA* 110: 8349-8356. - doi: [10.1073/pnas.1210595110](https://doi.org/10.1073/pnas.1210595110)
- Sayer JA, Maginnis S (2013). Forests in landscapes: expanding horizons for ecosystem forestry. In: "Forests in landscapes: ecosystem approaches to sustainability" (Maginnis S, Sayer JA eds). Earthscan, London, UK, pp. 177-191. [online] URL: <http://books.google.com/books?id=4oQgCkH3v28C>
- Schaich H, Milad M (2013). Forest biodiversity in a changing climate: which logic for conservation strategies? *Biodiversity and Conservation* 22: 1107-1114. - doi: [10.1007/s10531-013-0491-7](https://doi.org/10.1007/s10531-013-0491-7)
- Schall P, Ammer C (2013). How to quantify forest



- management intensity in Central European forests. *European Journal of Forest Research* 132: 379-396. - doi: [10.1007/s10342-013-0681-6](https://doi.org/10.1007/s10342-013-0681-6)
- Schnitzler A, Closset D (2003). Forest dynamics in unexploited birch (*Betula pendula*) stands in the Vosges (France): structure, architecture and light patterns. *Forest Ecology and Management* 183: 205-220. - doi: [10.1016/S0378-1127\(03\)00118-X](https://doi.org/10.1016/S0378-1127(03)00118-X)
- Schroeder LM, Ranius T, Ekblom B, Larsson S (2007). Spatial occurrence of a habitat-tracking saproxylic beetle inhabiting a managed forest landscape. *Ecological Applications* 17: 900-909. - doi: [10.1890/06-0090](https://doi.org/10.1890/06-0090)
- Schulte LA, Mitchell RJ, Hunter Jr ML, Franklin JF, Kevin McIntyre R, Palik BJ (2006). Evaluating the conceptual tools for forest biodiversity conservation and their implementation in the US. *Forest Ecology and Management* 232: 1-11. - doi: [10.1016/j.foreco.2006.05.009](https://doi.org/10.1016/j.foreco.2006.05.009)
- Sebek P, Altman J, Platak M, Cizek L (2013). Is active management the key to the conservation of saproxylic biodiversity? Pollarding promotes the formation of tree hollows. *PLoS One* 8 (3): e60456. - doi: [10.1371/journal.pone.0060456](https://doi.org/10.1371/journal.pone.0060456)
- Seibold S, Brandl R, Buse J, Hothorn T, Schmidl J, Thorn S, Müller J (2015). Association of extinction risk of saproxylic beetles with ecological degradation of forests in Europe. *Conservation Biology* 29: 382-390. - doi: [10.1111/cobi.12427](https://doi.org/10.1111/cobi.12427)
- Seidl R, Fernandes PM, Fonseca TF, Gillet F, Jönsson AM, Merganičová K, Netherer S, Arpacı A, Bontemps J-D, Bugmann H, González-Olabarria JR, Lasch P, Meredieu C, Moreira F, Schelhaas M-J, Mohren F (2011). Modelling natural disturbances in forest ecosystems: a review. *Ecological Modelling* 222: 903-924. - doi: [10.1016/j.ecolmodel.2010.09.040](https://doi.org/10.1016/j.ecolmodel.2010.09.040)
- Seidl R, Lexer MJ (2013). Forest management under climatic and social uncertainty: trade-offs between reducing climate change impacts and fostering adaptive capacity. *Journal of Environmental Management* 114: 461-469. - doi: [10.1016/j.jenvman.2012.09.028](https://doi.org/10.1016/j.jenvman.2012.09.028)
- Seidl R, Rammer W, Spies TA (2014a). Disturbance legacies increase the resilience of forest ecosystem structure, composition, and functioning. *Ecological Applications* 24: 2063-2077. - doi: [10.1890/14-0255.1](https://doi.org/10.1890/14-0255.1)
- Seidl R, Schelhaas M-J, Rammer W, Verkerk PJ (2014b). Increasing forest disturbances in Europe and their impact on carbon storage. *Nature Climate Change* 4: 806-810. - doi: [10.1038/nclimate2318](https://doi.org/10.1038/nclimate2318)
- Sharma T, Kurz WA, Stinson G, Pellatt MG, Li Q (2013). A 100-year conservation experiment: impacts on forest carbon stocks and fluxes. *Forest Ecology and Management* 310: 242-255. - doi: [10.1016/j.foreco.2013.06.048](https://doi.org/10.1016/j.foreco.2013.06.048)
- Shorohova E, Kneeshaw D, Kuuluvainen T, Gauthier S (2011). Variability and dynamics of old-growth forests in the circumboreal zone: implications for conservation, restoration and management. *Silva Fennica* 45: 785-806. - doi: [10.14214/sf.72](https://doi.org/10.14214/sf.72)
- Siitonen J (2012). Microhabitats. In: "Biodiversity in Dead Wood" (Stokland JN, Siitonen J, Jonsson BG eds). Cambridge University Press, Cambridge, UK, pp. 151-182.
- Sitzia T, Campagnaro T, Gatti E, Sommacal M, Kotze DJ (2015). Wildlife conservation through forestry abandonment: responses of beetle communities to habitat change in the Eastern Alps. *European Journal of Forest Research* 134 (3): 511-524. - doi: [10.1007/s10342-015-0868-0](https://doi.org/10.1007/s10342-015-0868-0)
- Stokland JN, Siitonen J, Jonsson BG, (2012). Biodiversity in dead wood. Cambridge University Press, Cambridge, UK, pp. 509. [online] URL: <http://books.google.com/books?id=2jcgAwAAQBAJ>
- Summers RW (2004). Use of pine snags by birds in different stand types of Scots Pine *Pinus sylvestris*: capsule hole-nesting birds tended to breed in the largest pine snags (standing dead trees) predominating in ancient native pine-wood and scarce in plantations. *Bird Study* 51: 212-221. - doi: [10.1080/00063650409461356](https://doi.org/10.1080/00063650409461356)
- Sverdrup-Thygeson A, Bendiksen E, Birkemoe T, Larsson KH (2014a). Do conservation measures in forest work? A comparison of three area-based conservation tools for wood-living species in boreal forests. *Forest Ecology and Management* 330: 8-16. - doi: [10.1016/j.foreco.2014.06.036](https://doi.org/10.1016/j.foreco.2014.06.036)
- Sverdrup-Thygeson A, Gustafsson L, Kouki J (2014b). Spatial and temporal scales relevant for conservation of dead-wood associated species: current status and perspectives. *Biodiversity and Conservation* 23: 513-535. - doi: [10.1007/s10531-014-0628-3](https://doi.org/10.1007/s10531-014-0628-3)
- Thomas S, Wagner R, Halteman W (2009). Influence of harvest gaps and coarse woody material on click beetles (Coleoptera: Elateridae) in Maine's Acadian forest. *Biodiversity and Conservation* 18: 2405-2419. - doi: [10.1007/s10531-009-9597-3](https://doi.org/10.1007/s10531-009-9597-3)
- Thorn S, Bässler C, Gottschalk T, Hothorn T, Busler H, Raffa K, Müller J (2014). New insights into the consequences of post-windthrow salvage logging revealed by functional structure of saproxylic beetles assemblages. *PLoS One* 9 (7): e101757. - doi: [10.1371/journal.pone.0101757](https://doi.org/10.1371/journal.pone.0101757)
- Timonen J, Siitonen J, Gustafsson L, Kotiaho JS, Stokland JN, Sverdrup-Thygeson A, Mönkkönen M (2010). Woodland key habitats in northern Europe: concepts, inventory and protection. *Scandinavian Journal of Forest Research* 25: 309-324. - doi: [10.1080/02827581.2010.497160](https://doi.org/10.1080/02827581.2010.497160)
- Toivanen T, Kotiaho JS (2007). Mimicking natural disturbances of boreal forests: the effects of controlled burning and creating dead wood on beetle diversity. *Biodiversity and Conservation* 16: 3193-3211. - doi: [10.1007/s10531-007-9172-8](https://doi.org/10.1007/s10531-007-9172-8)
- Tscharntke T, Clough Y, Wanger TC, Jackson L, Motzke I, Perfecto I, Vandermeer J, Whitbread A (2012). Global food security, biodiversity conservation and the future of agricultural intensification. *Biological Conservation* 151: 53-59. - doi: [10.1016/j.biocon.2012.01.068](https://doi.org/10.1016/j.biocon.2012.01.068)
- Ulyshen MD, Hanula JL, Horn S, Kilgo JC, Moorman CE (2004). Spatial and temporal patterns of beetles associated with coarse woody debris in managed bottomland hardwood forests. *Forest Ecology and Management* 199: 259-272. - doi: [10.1016/j.foreco.2004.05.046](https://doi.org/10.1016/j.foreco.2004.05.046)
- Villard MA, Jonsson BG (2009). A plea for quantitative targets in biodiversity conservation. In: "Setting conservation targets for managed landscapes" (Villard MA, Jonsson BG eds). Cambridge University Press, Cambridge, UK, pp. 1-8. [online] URL: [http://www.newbooks-services.de/MediaFiles/Texts/1/9780521877091\\_Excerpt\\_001.pdf](http://www.newbooks-services.de/MediaFiles/Texts/1/9780521877091_Excerpt_001.pdf)
- Vuidot A, Paillet Y, Archaux F, Gosselin F (2011). Influence of tree characteristics and forest management on tree microhabitats. *Biological Conservation* 144: 441-450. - doi: [10.1016/j.biocon.2010.09.030](https://doi.org/10.1016/j.biocon.2010.09.030)
- Warren M, Fuller R (1993). Woodland rides and glades their management for wildlife. Joint Nature Conservation Committee, Peterborough, UK, pp. 31.
- Watts ME, Ball IR, Stewart RS, Klein CJ, Wilson K, Steinback C, Lourival R, Kircher L, Possingham HP (2009). Marxan with Zones: software for optimal conservation based land-and sea-use zoning. *Environmental Modelling and Software* 24: 1513-1521. - doi: [10.1016/j.envsoft.2009.06.005](https://doi.org/10.1016/j.envsoft.2009.06.005)
- Winter M-B, Ammer C, Baier R, Donato DC, Seibold S, Müller J (2015). Multi-taxon alpha diversity following bark beetle disturbance: evaluating multi-decade persistence of a diverse early-seral phase. *Forest Ecology and Management* 338: 32-45. - doi: [10.1016/j.foreco.2014.11.019](https://doi.org/10.1016/j.foreco.2014.11.019)
- Winter S, Brambach F (2011). Determination of a common forest life cycle assessment method for biodiversity evaluation. *Forest Ecology and Management* 262: 2120-2132. - doi: [10.1016/j.for-eco.2011.07.036](https://doi.org/10.1016/j.for-eco.2011.07.036)
- Wirth C, Messier C, Bergeron Y, Frank D, Fankhänel A, (2009). Old-growth forest definitions: a pragmatic view. In: "Old-Growth Forests" (Wirth C, Gleixner G, Heimann M eds). Springer Berlin, Heidelberg, Germany, pp. 11-33. - doi: [10.1007/978-3-540-92706-8\\_2](https://doi.org/10.1007/978-3-540-92706-8_2)
- Zapponi L, Minari E, Longo L, Toni I, Mason F, Campanaro A (2015). The Habitat-Trees experiment: using exotic tree species as new microhabitats for the native fauna. *iForest - Biogeosciences and Forestry* 8: 464-470. - doi: [10.3832/ifor1281-007](https://doi.org/10.3832/ifor1281-007)
- Zapponi L, Del Bianco M, Luiselli L, Catorci A, Bologna MA (2013). Assessing environmental requirements effects on forest fragmentation sensitivity in two arboreal rodents. *Mammalian Biology - Zeitschrift für Säugetierkunde* 78: 157-163. - doi: [10.1016/j.mambio.2012.08.005](https://doi.org/10.1016/j.mambio.2012.08.005)