

## Species-specific morphological and physiological characteristics and progressive nitrogen limitation under elevated CO<sub>2</sub> concentration

Woo Kyung Song<sup>(1-2)</sup>,  
Si Yeon Byeon<sup>(2)</sup>,  
HoonTaek Lee<sup>(1-2)</sup>,  
Min Su Lee<sup>(2)</sup>,  
Daun Ryu<sup>(3)</sup>,  
Jun Won Kang<sup>(4)</sup>,  
Sim Hee Han<sup>(5)</sup>,  
Chang Young Oh<sup>(6)</sup>,  
Hyun Seok Kim<sup>(2-3-7-8)</sup>

Elevated atmospheric CO<sub>2</sub> (eCO<sub>2</sub>) concentration initially enhances photosynthesis, growth and ecosystem productivity, but the excessive use of nitrogen due to the increased productivity causes uncertainty in long-term ecosystem responses. We exposed Korean red pine, Chinese ash, and Korean mountain ash to current atmospheric CO<sub>2</sub> concentration (aCO<sub>2</sub>), 1.4 times higher CO<sub>2</sub> concentration (eCO<sub>2</sub>1.4), and 1.8 times higher CO<sub>2</sub> concentration (eCO<sub>2</sub>1.8) in an Open-Top Chamber (OTC) experiment for eight years (2010-2017) to investigate the effect on the morphological and physiological properties of trees. We also assessed whether nitrogen limitation occurred with time by comparing leaf and soil nitrogen concentration. CO<sub>2</sub> fertilization effect was observed on tree growth for the first two years ( $p < 0.05$ ), but there was no difference thereafter. For photosynthetic properties, CO<sub>2</sub> effects were species-specific; no effects on Korean red pine and Chinese ash vs. significant effect on Korean mountain ash. However, maximum photosynthetic and carboxylation rates significantly decreased by 24.3% and 31.3% from 2013 to 2017, respectively. Leaf nitrogen significantly decreased by 21.0 % at eCO<sub>2</sub>1.4 and 18.5 % at eCO<sub>2</sub>1.8 compared with aCO<sub>2</sub> treatment. This study showed the decline of leaf nitrogen and species-specific responses to long-term high CO<sub>2</sub> concentration, which will effect on species competition and ecosystem succession.

**Keywords:** Elevated CO<sub>2</sub>, Photosynthetic Properties, Down-regulation, Progressive Nitrogen Limitation, Carbon dioxide

### Introduction

Elevated CO<sub>2</sub> concentration has become a common phenomenon in the Earth's atmosphere over the last half-century (IPCC 2013). The indiscreet use of fossil fuels and deforestation has raised the atmospheric CO<sub>2</sub> concentration, thus promoting photosynthesis and growth (Drake et al. 1997, Ainsworth & Long 2005, Sang et al. 2019) and affecting the productivity of plants (McCarthy et al. 2010, Norby et al. 2010), which is referred to as the CO<sub>2</sub> fertilization

effect (Norby et al. 2010). However, there are conflicting opinions on whether such an effect will continue with increasing CO<sub>2</sub> concentration (Hungate et al. 2003). In general, the productivity of terrestrial ecosystems is affected by nutrient availability in the soil (Terrer et al. 2018). It should be taken into account that, if CO<sub>2</sub> fertilization promotes plant growth, the excessive nutrient use will result in insufficient nutrients in the soil, failing to meet the nutrient requirements needed for plant growth (Hun-

gate et al. 2003, Wang & Houlton 2009) and decreasing the CO<sub>2</sub> fertilization effect. Therefore, the expected increase of ecosystem productivity and carbon storage due to CO<sub>2</sub> fertilization is still uncertain (Terrer et al. 2018).

Early studies using pots and growth chambers hardly reflected the real forest ecosystem conditions (Curtis & Wang 1998, Norby et al. 2010), therefore experiments with Open-Top chamber (OTC) and Free Air CO<sub>2</sub> Enrichment (FACE) techniques have been widely employed (Norby et al. 2010). Many OTC and FACE experiments reported an increased growth and photosynthesis under elevated CO<sub>2</sub> (Hungate et al. 2013, Talhelm et al. 2014). However, whether these effects would last for a long time is questioned. In particular, experiments on elevated CO<sub>2</sub> and soil nitrogen fertilization (Finzi et al. 2007), which were conducted at the Duke-FACE and Oak Ridge-FACE, confirmed that nitrogen can directly limit the increase in forest productivity due to the carbon fertilization response, highlighting the relevance of nitrogen on forest productivity.

Nitrogen is an important component of plant photosynthetic organs and of functional and structural proteins (Kwon et al. 2019). In particular, nitrogen in the leaves constitutes the photosynthetic enzymes, such as chlorophyll and rubisco (ribulose-1,5-biphosphate carboxylase/oxygenase), which determines the maximum carboxyla-

□ (1) Forest Ecology and Climate Change Division, National Institute of Forest Science, Seoul 02455 (Republic of Korea); (2) Department of Forest Sciences, Seoul National University, Seoul 08826 (Republic of Korea); (3) Interdisciplinary Program in Agricultural and Forest Meteorology, Seoul National University, Seoul 08826 (Republic of Korea); (4) Department of Forestry, Kyungpook National University, Daegu 41566 (Republic of Korea); (5) Forest Biotechnology Division, National Institute of Forest Science, Gyeonggi 16631 (Republic of Korea); (6) Urban Forests Research Center, National Institute of Forest Science, Seoul 02455 (Republic of Korea); (7) National Center for AgroMeteorology, Seoul 08826 (Republic of Korea); (8) Research Institute for Agriculture and Life Sciences, Seoul National University, Seoul 08826 (Republic of Korea)

@ Hyun Seok Kim ([cameroncrazies@snu.ac.kr](mailto:cameroncrazies@snu.ac.kr))

Received: Nov 08, 2019 - Accepted: May 06, 2020

**Citation:** Song WK, Byeon SY, Lee HT, Lee MS, Ryu D, Kang JW, Han SH, Oh CY, Kim HS (2020). Species-specific morphological and physiological characteristics and progressive nitrogen limitation under elevated CO<sub>2</sub> concentration. *iForest* 13: 270-278. - doi: [10.3832/for3288-013](https://doi.org/10.3832/for3288-013) [online 2020-07-03]

Communicated by: *Silvano Fares*

tion rate ( $V_{cmax}$  – Makino & Osmond 1991, Hikosaka 2004), with a decisive influence on forest productivity (Harley et al. 1992, Harrison et al. 2009). However, owing to increased productivity by long-term exposure to a high  $CO_2$  concentrations, the consumption of nutrients may overcome the nutrients input, leading to a reduction in effective nutrients in the soil, and consequently to a reduction of plants' photosynthesis ability. In particular, as the nitrogen concentration absorbed by plants decreases, its concentration in the leaves is lowered, thus decreasing photosynthesis. This in turn results in a progressive nitrogen limitation (PNL), which offsets the  $CO_2$  fertilization effect (Drake et al. 1997, Luo et al. 2004, Cai et al. 2018). This phenomenon is more common under nitrogen deficiency in the soil (Pettersson & McDonald 1994, McCarthy et al. 2010).

Under PNL, an increase in productivity due to the  $CO_2$  fertilization effect is a short-term phenomenon, whereas in the long term, the ecosystem productivity might decrease again and return to its original state (Luo et al. 2004, Finzi et al. 2007). This suggests that increased forest productivity due to a high atmospheric concentration of  $CO_2$  might be inhibited by soil nitrogen deficiency (Feng et al. 2015). Reich et al. (2006) and Feng et al. (2015) confirmed that the response of plants to  $CO_2$  fertilization was limited by nitrogen deficiency. Moreover, PNL is expected to be more prominent when the high  $CO_2$  concentration environment is maintained for a long time (Rütting 2017).

The main objective of this study was to determine the occurrence of nitrogen limitation under long-term  $CO_2$  fertilization on three major species of the Korean temperate region: Korean red pine (*Pinus densiflora*), Chinese ash (*Fraxinus rhynchophylla*), and Korean mountain ash (*Sorbus alnifolia*). The effect of the increase in atmospheric  $CO_2$  concentration on seedling diameter and height, as well as photosynthetic properties, such as maximum photosynthetic rate, maximum carboxylation rate, and maximum electron transfer rate, was evaluated over a period of 8 years. Moreover, we compared the nitrogen concentration in the leaves and soils to investigate whether the increased productivity due to  $CO_2$  fertilization is a short-term effect owing to a progressive nitrogen limitation (PNL) in the soil.

## Materials and methods

### Experimental system and design

This study was carried out in an Open Top Chamber (OTC) of the Forest Biotechnology Division at National Institute of Forest Science in Suwon-si, Gyeonggi-do, Republic of Korea (37° 15' 04" N, 126° 57' 29" E). The OTC was made of a decagon structure of diameter 10 m and height 7 m (see Fig. S1 in Supplementary material). The angle of the roof was 45° in order to maintain an open-

ing ratio of 75% or higher. A polyolefin film of thickness 0.15 mm with a light transmittance of approximately 88%, low specific gravity, and excellent chemical and water resistance was used as an external covering material (Lee et al. 2012).

The atmospheric  $CO_2$  concentrations kept in the chambers during the experiment were: (i) the current atmospheric  $CO_2$  concentration (Chamber 1, a $CO_2$ ); (ii) 1.4 times higher  $CO_2$  concentration than the current concentration (Chamber 2, e $CO_2$ 1.4), which is expected to be the atmospheric  $CO_2$  concentration by 2050, according to the IPCC scenario (IPCC 2013); and (iii) 1.8 times higher  $CO_2$  concentration than the current concentration, which is expected to be the atmospheric  $CO_2$  concentration by 2070 (Chamber 3, e $CO_2$ 1.8). The exposure of  $CO_2$  were conducted for 10 hours (08:00-18:00) a day during the growing season (April-November). Pest controls were conducted every year and weed controls were conducted for first four years of the experiment. The annual average, minimum and maximum monthly temperature in chambers were  $12.4 \pm 0.2$  °C,  $-1.3 \pm 0.4$  °C,  $22.5 \pm 0.2$  °C, respectively. The annual rainfall range was 751.16 mm yr<sup>-1</sup> to 1975.96 mm yr<sup>-1</sup> over the 9 years of the experimental period.

In each chamber, the same three clones for each species were tested. Four-year-old seedlings of Korean red pine (*Pinus densiflora*), two-year-old seedlings of Chinese ash (*Fraxinus rhynchophylla*), and two-year-old seedlings of Korean mountain ash (*Sorbus alnifolia*) were planted in September 2009. The density within the OTC was 2547 seedlings ha<sup>-1</sup>.

### Growth measurements

Growth measurements (height and diameter) on trees started in May 2010 and were repeated every year until 2017 at the beginning (mid-April, mean daily temperature: ~10 °C) and the end (end of October, mean daily temperature: ~5 °C) of the growing season. The annual increase in growth was obtained as the difference in these two measures. Also, the lack of differences in diameter between the last measurement of the previous year and the first of the subsequent year was verified.

Tree height was recorded using a measuring rod (A-15, SENSHIN Industry Co., LTD., Osaka, Japan), while the tree diameter was measured twice in orthogonal directions 10 cm above the root collar using a digital Vernier callipers (CD-10CPX, Mitutoyo, Kawasaki, Japan), and the average value was used. Exceptionally, since 2016, Korean red pine and Chinese ash trees were measured using a diameter tape (F10-02DM, KDS, Malaysia) because of their large diameters.

### Gas exchange measurements

The photosynthetic parameters were measured on tree leaves using a portable device LI-6400 (LI-COR Inc., Lincoln, NE,

USA). From 2013 to 2017, except for 2015, the parameters were measured 1-3 times a year from June to August on 3 sunlit leaves for each species. The temperature of the leaves was set at 25 °C and the relative humidity at 55%-60%. The leaves were stabilized before measurements. The area of leaves was set at 6 cm<sup>2</sup> for all the species except Korean red pine. For this species the leaf area was recalculated by measuring the actual leaf area using a scanner after all measurements.

The light response curve was obtained using the photosynthetic rate recorded by sequentially varying the intensity of light irradiated on the leaves, using the following sequence in all the chambers: 1400, 1200, 1000, 800, 600, 400, 200, 100, 75, 50, 25, 0, and 1200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . The reference  $CO_2$  supplied in each LI-6400 chamber was set at the atmospheric  $CO_2$  in each OTC. The maximum photosynthetic rate ( $A_{max}$ ) was estimated as the photosynthetic rate at light saturation (1200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) under the  $CO_2$  concentration in the chambers, which was measured using the light response curve (Taiz & Zeiger 2010).

The A/C<sub>i</sub> curve was achieved by varying the reference  $CO_2$  concentration at light saturation point in the following order: (i) a $CO_2$ : 400, 300, 200, 100, 75, 50, 25, 0, 400, 400, 600, 800, 1000, and 1200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ; (ii) e $CO_2$ 1.4: 560, 400, 300, 200, 100, 75, 50, 25, 0, 560, 560, 800, 1000, and 1200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ; and (iii) e $CO_2$ 1.8: 720, 600, 400, 300, 200, 100, 75, 50, 25, 0, 720, 720, 1000, and 1200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . The parameters  $V_{cmax}$  and  $J_{max}$  were derived using the model proposed by Sharkey (2016) and estimated in Excel™ spreadsheet version 2.0 (Microsoft, Redmond, WA, USA).


### Leaf total nitrogen analysis

Three sunlit leaves were chosen on the same branch and used for photosynthetic measurements ( $n = 9$  for each species, three repetitions for three species), and 1 cm<sup>2</sup> leaf disk was collected in 2017. The leaves were dried in a 70 °C in the lab for more than 72 h, then crushed using a Fast-Prep-24® crusher (MP Biomedicals, Solon, OH, USA), homogenized, and finally analysed for nitrogen content using CHNS Analyzer Flash EA 1112® (Thermo Electron Corporation, USA) at the National Instrumentation Center for Environmental Management (NICEM).

### Soil nitrogen analysis

Before planting trees in 2009, the soil in the OTC was excavated to a depth of 1 m and replaced with forest soil to control the soil characteristics of all the treatment groups. Soil at depth < 30 cm was composed of forest soil and sand with a ratio of 1:1 (Lee et al. 2012). For all treatment groups, samples of soil from 0 to 5 cm depth were collected at five randomly selected points in September 2017, air-dried for 3 days at room temperature, and then analysed using the CHNS Analyzer Flash EA

**Fig. 1** - Effects of elevated CO<sub>2</sub> concentration (aCO<sub>2</sub>, Chamber 1; eCO<sub>2</sub>1.4, Chamber 2; eCO<sub>2</sub>1.8, Chamber 3) on average tree diameter (a, b, c) and height (d, e, f) growth per year from 2010 to 2017. There were no significant differences among CO<sub>2</sub> treatments ( $p > 0.05$ ).



1112 described above at the NICEM.

**Statistical analysis**

All statistical analyses were carried using the software R ver. 3.3.2 (R Core Team 2016). Growth, photosynthesis variables and leaf total nitrogen over time were analyzed by repeated-measures ANOVA with CO<sub>2</sub> treatments as a fixed factor and the recording year as the repeated-measure. When significant CO<sub>2</sub> × Year interactions and CO<sub>2</sub> or Year effect were detected, means were compared using Tukey post-hoc comparisons. In addition, one-way ANOVA of individual species and Tukey post-hoc comparisons were conducted to assess the CO<sub>2</sub> effect for each year.

**Results**

To investigate the CO<sub>2</sub> fertilization effect over the period 2010-2017, diameter and height increments of trees of different treatments and species were recorded (Tab. 1, Tab. 2). The cumulative increments for height and diameter over the 8 years of the experiment are presented in Fig. 1.

**Diameter growth**

Tab. 1 summarizes the annual diameter in-

crease measured after CO<sub>2</sub> exposure. The results showed a significant increment in diameter growth under high CO<sub>2</sub> concentration due to the CO<sub>2</sub> fertilization effect ( $p < 0.001$  – Tab. 3). On average, the difference between aCO<sub>2</sub> and eCO<sub>2</sub>1.8 was 47.7%, 25.3%, and 3.1% in Korean mountain ash ( $41.4 \pm 6.2$  vs.  $61.1 \pm 7.9$  mm), Korean red pine ( $144.7 \pm 15.1$  vs.  $181.3 \pm 13.5$  mm), and Chinese ash ( $70.9 \pm 9.2$  vs.  $73.1 \pm 5.5$  mm), respectively.

The annual diameter increment of individual species under eCO<sub>2</sub> was significantly enhanced in all the species in 2010-2011 (maximum  $p = 0.029$  – Tab. 3). However, since 2012, there was no significant difference in diameter increment under eCO<sub>2</sub>. Korean red pine showed a significant difference in diameter growth under eCO<sub>2</sub> in 2011 in the order of aCO<sub>2</sub> < eCO<sub>2</sub>1.4 < eCO<sub>2</sub>1.8, an increase of about 100.0% (aCO<sub>2</sub> vs. eCO<sub>2</sub>1.8,  $p = 0.015$  – Tab. 1). Korean mountain ash showed significant difference under eCO<sub>2</sub> in 2010, an increase of about 57.9% (aCO<sub>2</sub>:  $5.7 \pm 0.7$  vs. eCO<sub>2</sub>1.8:  $9.0 \pm 0.5$  mm;  $p = 0.02$ ).

**Height growth**

Similarly to growth in diameter, annual in-

crease in tree height was recorded each year from 2010 until 2017 (Fig. 1). For all the species, the increase in height was high under eCO<sub>2</sub> ( $p < 0.01$  – Tab. 2). On average, the difference between aCO<sub>2</sub> and eCO<sub>2</sub>1.8 was 22.01%, 18.3% and 14.3% in Korean red pine ( $470.7 \pm 22.8$  vs.  $574.3 \pm 72.5$  cm), Chinese ash ( $585.3 \pm 39.1$  vs.  $692.3 \pm 48.1$  cm) and Korean mountain ash ( $299.3 \pm 40.5$  vs.  $342.0 \pm 8.2$  cm), respectively.

Tab. 2 summarizes the annual growth of tree height measured after CO<sub>2</sub> exposure. For all the species, results were similar to the diameter growth. Chinese ash showed a significant difference in height growth under eCO<sub>2</sub> in 2010 ( $p = 0.029$ ). Differences were observed under eCO<sub>2</sub>1.8 and aCO<sub>2</sub>, which was increased about 57.9%. Korean red pine showed a significant increase under eCO<sub>2</sub> in 2011, an increase of about 110.7% ( $p = 0.006$ ). However, after 2012, there was no difference in the increase in height due to CO<sub>2</sub> concentration for all the species.

**Maximum photosynthetic rate (A<sub>max</sub>)**

The average A<sub>max</sub> under eCO<sub>2</sub> measured from 2013 increased in the following order: aCO<sub>2</sub> ( $12.7 \pm 0.7 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) < eCO<sub>2</sub>1.4 ( $13.6$

**Tab. 1** - Annual growth in diameter (mm, mean ± SE) of Korean red pine (*Pinus densiflora*, Pd), Chinese ash (*Fraxinus rhynchophylla*, Fr) and Korean mountain ash (*Sorbus alnifolia*, Sa) in response to aCO<sub>2</sub> (Chamber 1), eCO<sub>2</sub>1.4 (Chamber 2) and eCO<sub>2</sub>1.8 (Chamber 3) during 8 years. Different lowercase letters indicate significant ( $p < 0.05$ ) multiple comparison results among CO<sub>2</sub> treatments (one-way ANOVA followed by Tukey post-hoc test). (ns): not significant.

Species	Treatments	Diameter growth (mm)							
		2010	2011	2012	2013	2014	2015	2016	2017
Pd	aCO <sub>2</sub>	11.2 ± 1.4 <sup>ns</sup>	8.5 ± 1.4 <sup>b</sup>	11.6 ± 10.1 <sup>ns</sup>	19.0 ± 3.5 <sup>ns</sup>	9.3 ± 3.3 <sup>ns</sup>	18.8 ± 7.7 <sup>ns</sup>	25.8 ± 7.7 <sup>ns</sup>	16.7 ± 5.2 <sup>ns</sup>
	eCO <sub>2</sub> 1.4	11.9 ± 0.2 <sup>ns</sup>	11.4 ± 1.1 <sup>ab</sup>	20.3 ± 4.2 <sup>ns</sup>	18.0 ± 6.7 <sup>ns</sup>	12.3 ± 3.8 <sup>ns</sup>	26.3 ± 6.3 <sup>ns</sup>	19.3 ± 9.1 <sup>ns</sup>	16.5 ± 3.8 <sup>ns</sup>
	CeCO <sub>2</sub> 1.8	12.5 ± 2.9 <sup>ns</sup>	17.0 ± 1.7 <sup>a</sup>	8.9 ± 3.4 <sup>ns</sup>	40.0 ± 24.1 <sup>ns</sup>	20.1 ± 4.8 <sup>ns</sup>	25.2 ± 0.9 <sup>ns</sup>	31.0 ± 5.4 <sup>ns</sup>	19.2 ± 3.0 <sup>ns</sup>
Fr	aCO <sub>2</sub>	11.7 ± 0.8 <sup>ns</sup>	8.6 ± 2.3 <sup>ns</sup>	8.8 ± 3.4 <sup>ns</sup>	2.0 ± 1.2 <sup>ns</sup>	12.6 ± 5.6 <sup>ns</sup>	13.8 ± 1.1 <sup>ns</sup>	6.8 ± 1.7 <sup>ns</sup>	5.5 ± 1.0 <sup>ns</sup>
	eCO <sub>2</sub> 1.4	14.4 ± 2.0 <sup>ns</sup>	7.6 ± 1.4 <sup>ns</sup>	4.7 ± 6.0 <sup>ns</sup>	3.0 ± 1.7 <sup>ns</sup>	8.4 ± 4.6 <sup>ns</sup>	11.0 ± 1.9 <sup>ns</sup>	8.2 ± 2.0 <sup>ns</sup>	5.0 ± 2.2 <sup>ns</sup>
	CeCO <sub>2</sub> 1.8	18.5 ± 2.5 <sup>ns</sup>	5.3 ± 2.9 <sup>ns</sup>	3.4 ± 6.0 <sup>ns</sup>	5.0 ± 3.1 <sup>ns</sup>	7.0 ± 0.7 <sup>ns</sup>	5.5 ± 0.1 <sup>ns</sup>	10.5 ± 1.9 <sup>ns</sup>	10.6 ± 2.7 <sup>ns</sup>
Sa	aCO <sub>2</sub>	5.7 ± 0.7 <sup>b</sup>	4.5 ± 0.5 <sup>ns</sup>	3.8 ± 4.9 <sup>ns</sup>	1.7 ± 0.7 <sup>ns</sup>	3.0 ± 0.6 <sup>ns</sup>	6.1 ± 2.5 <sup>ns</sup>	6.0 ± 3.3 <sup>ns</sup>	4.9 ± 0.9 <sup>ns</sup>
	CeCO <sub>2</sub> 1.4	7.0 ± 0.5 <sup>ab</sup>	6.3 ± 0.2 <sup>ns</sup>	5.1 ± 2.7 <sup>ns</sup>	2.7 ± 0.8 <sup>ns</sup>	4.7 ± 0.9 <sup>ns</sup>	9.0 ± 1.9 <sup>ns</sup>	8.0 ± 0.9 <sup>ns</sup>	7.6 ± 2.9 <sup>ns</sup>
	eCO <sub>2</sub> 1.8	9.0 ± 0.5 <sup>a</sup>	6.9 ± 0.6 <sup>ns</sup>	3.2 ± 2.6 <sup>ns</sup>	7.7 ± 4.7 <sup>ns</sup>	2.3 ± 2.0 <sup>ns</sup>	9.6 ± 1.2 <sup>ns</sup>	7.4 ± 1.7 <sup>ns</sup>	3.5 ± 2.3 <sup>ns</sup>



**Tab. 2** - Annual growth in height (cm, mean ± SE) of Korean red pine (*Pinus densiflora*, Pd), Chinese ash (*Fraxinus rhynchophylla*, Fr) and Korean mountain ash (*Sorbus alnifolia*, Sa) in response to aCO<sub>2</sub> (Chamber 1), eCO<sub>2</sub>1.4 (Chamber 2) and eCO<sub>2</sub>1.8 (Chamber 3) during 8 years. Different lowercase letters indicate significant ( $p < 0.05$ ) multiple comparison results among CO<sub>2</sub> treatments (one-way ANOVA followed by Tukey post-hoc test). (ns): not significant.

Species	Treatment	Height growth (cm)							
		2010	2011	2012	2013	2014	2015	2016	2017
Pd	aCO <sub>2</sub>	10.5 ± 1.6 <sup>ns</sup>	27.0 ± 0.9 <sup>b</sup>	65.2 ± 6.6 <sup>ns</sup>	52.3 ± 13.0 <sup>ns</sup>	41.3 ± 18.3 <sup>ns</sup>	72.0 ± 5.1 <sup>ns</sup>	35.3 ± 2.7 <sup>ns</sup>	73.0 ± 22.1 <sup>ns</sup>
	eCO <sub>2</sub> 1.4	11.5 ± 1.5 <sup>ns</sup>	11.4 ± 1.1 <sup>b</sup>	57.6 ± 19.1 <sup>ns</sup>	59.3 ± 9.5 <sup>ns</sup>	65.7 ± 8.7 <sup>ns</sup>	94.7 ± 49.6 <sup>ns</sup>	50.0 ± 46.0 <sup>ns</sup>	17.7 ± 24.6 <sup>ns</sup>
	eCO <sub>2</sub> 1.8	6.3 ± 1.6 <sup>ns</sup>	56.9 ± 7.5 <sup>a</sup>	70.5 ± 34.6 <sup>ns</sup>	77.0 ± 5.3 <sup>ns</sup>	63.0 ± 6.1 <sup>ns</sup>	60.0 ± 20.0 <sup>ns</sup>	99.0 ± 21.7 <sup>ns</sup>	48.0 ± 17.1 <sup>ns</sup>
Fr	aCO <sub>2</sub>	136.2 ± 23.3 <sup>b</sup>	86.5 ± 11.9 <sup>ns</sup>	29.0 ± 16.0 <sup>ns</sup>	13.0 ± 2.1 <sup>ns</sup>	90.7 ± 34.7 <sup>ns</sup>	89.3 ± 44.9 <sup>ns</sup>	30.2 ± 22.3 <sup>ns</sup>	78.3 ± 2.6 <sup>ns</sup>
	eCO <sub>2</sub> 1.4	145.4 ± 14.8 <sup>ab</sup>	7.6 ± 1.3 <sup>ns</sup>	22.9 ± 3.3 <sup>ns</sup>	27.7 ± 12.8 <sup>ns</sup>	60.3 ± 29.5 <sup>ns</sup>	73.7 ± 18.2 <sup>ns</sup>	29.3 ± 20.1 <sup>ns</sup>	86.7 ± 0.9 <sup>ns</sup>
	eCO <sub>2</sub> 1.8	215.0 ± 8.3 <sup>a</sup>	84.4 ± 18.4 <sup>ns</sup>	9.0 ± 3.3 <sup>ns</sup>	75.7 ± 33.1 <sup>ns</sup>	92.0 ± 40.8 <sup>ns</sup>	81.7 ± 10.9 <sup>ns</sup>	70.3 ± 19.7 <sup>ns</sup>	82.7 ± 39.8 <sup>ns</sup>
Sa	aCO <sub>2</sub>	39.1 ± 7.0 <sup>ns</sup>	27.0 ± 4.3 <sup>ns</sup>	18.2 ± 12.8 <sup>ns</sup>	51.0 ± 16.8 <sup>ns</sup>	42.3 ± 20.3 <sup>ns</sup>	38.0 ± 3.21 <sup>ns</sup>	43.0 ± 2.1 <sup>ns</sup>	18.7 ± 9.8 <sup>ns</sup>
	eCO <sub>2</sub> 1.4	55.7 ± 15.9 <sup>ns</sup>	33.4 ± 9.7 <sup>ns</sup>	10.6 ± 5.5 <sup>ns</sup>	53.7 ± 33.7 <sup>ns</sup>	17.7 ± 7.2 <sup>ns</sup>	46.6 ± 13.1 <sup>ns</sup>	59.0 ± 11.5 <sup>ns</sup>	50.0 ± 8.0 <sup>ns</sup>
	eCO <sub>2</sub> 1.8	85.8 ± 11.4 <sup>ns</sup>	17.3 ± 6.5 <sup>ns</sup>	22.3 ± 15.1 <sup>ns</sup>	16.0 ± 0.7 <sup>ns</sup>	29.3 ± 14.4 <sup>ns</sup>	45.3 ± 7.8 <sup>ns</sup>	35.7 ± 6.0 <sup>ns</sup>	36.0 ± 18.6 <sup>ns</sup>

**Tab. 3** - Results statistics for diameter, height, A<sub>max</sub>, V<sub>cmax</sub>, J<sub>max</sub> and leaf total N of all species with F statistics and p-values from two-way repeated-measure ANOVA.

Parameter	Factor	F	p
Diameter	CO <sub>2</sub>	58.03	< 0.001
	Year	19.08	< 0.001
	CO <sub>2</sub> × Year	1.04	0.982
Height	CO <sub>2</sub>	17.727	< 0.001
	Year	9.201	< 0.001
	CO <sub>2</sub> × Year	0.367	0.416
A <sub>max</sub>	CO <sub>2</sub>	2.942	0.06
	Year	13.474	< 0.001
	CO <sub>2</sub> × Year	0.771	0.594
V <sub>cmax</sub>	CO <sub>2</sub>	1.662	0.194
	Year	2.791	0.030
	CO <sub>2</sub> × Year	0.545	0.819
J <sub>max</sub>	CO <sub>2</sub>	0.192	0.825
	Year	0.628	0.644
	CO <sub>2</sub> × Year	0.251	0.979
Leaf Total N	CO <sub>2</sub>	3.701	0.026
	Year	5.295	0.002
	CO <sub>2</sub> × Year	0.945	0.464

age A<sub>max</sub> (minimum  $p = 0.094$  – Tab. 4), but Korean mountain ash showed significant CO<sub>2</sub> effect. Korean red pine in all the years under eCO<sub>2</sub> showed no specific tendency. The A<sub>max</sub> of Korean mountain ash was  $14.0 \pm 1.3 \mu\text{mol m}^{-2} \text{s}^{-1}$  under eCO<sub>2</sub>1.8, which showed an enhancement from  $10.7 \pm 0.7 \mu\text{mol m}^{-2} \text{s}^{-1}$  under aCO<sub>2</sub> ( $p = 0.014$ ). The difference in the A<sub>max</sub> due to CO<sub>2</sub> exposure in each year did not show significant differences (Fig. 2c). The A<sub>max</sub> of Chinese ash decreased in the following order: aCO<sub>2</sub> ( $12.4 \pm 1.2 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) < eCO<sub>2</sub>1.4 ( $15.0 \pm 1.7 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) < eCO<sub>2</sub>1.8 ( $17.4 \pm 1.6 \mu\text{mol m}^{-2} \text{s}^{-1}$ ), and also decreased steadily from 2013 ( $19.4 \pm 1.1 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to 2017 ( $11.7 \pm 0.8 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) by about 39.7%, though not significantly ( $p = 0.094$ ,  $p = 0.095$ , respectively – Fig. 2b). However, the enhancement of A<sub>max</sub> due to elevated CO<sub>2</sub> was 27.7% at eCO<sub>2</sub>1.8, and the increase was constantly maintained at 28.6% under eCO<sub>2</sub>1.8 in 2013 and 27.5% in 2017 relative to the rate under aCO<sub>2</sub>.

**Maximum rate of carboxylation (V<sub>cmax</sub>) and electron transport (J<sub>max</sub>)**


Unlike the maximum photosynthetic rate, the V<sub>cmax</sub> was decreased of about 13.6% from eCO<sub>2</sub>1.8 ( $49.6 \pm 3.4 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to aCO<sub>2</sub> ( $57.4 \pm 3.5 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) but did not dif-

$\pm 0.8 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) < eCO<sub>2</sub>1.8 ( $14.9 \pm 0.8 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) but did not differ significantly ( $p = 0.06$  – Tab. 3). In terms of response by year, the average A<sub>max</sub> decreased by about 28.9%, from  $15.9 \pm 0.8 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2013 to  $11.3 \pm 0.9 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2016 ( $p < 0.001$ ). However, these results showed differences among species and year (Fig. 2, Tab. 4). Korean red pine and Chinese ash did not show a significant difference in the aver-

**Tab. 4** - Results statistics of diameter, height, A<sub>max</sub>, V<sub>cmax</sub>, J<sub>max</sub> and leaf total N of the studied species with F statistics and p-values from two-way repeated measured ANOVA. (Pd): Korean red pine (*Pinus densiflora*); (Fr): Chinese ash (*Fraxinus rhynchophylla*); (Sa): Korean mountain ash (*Sorbus alnifolia*).

Species	Factors	Diameter		Height		A <sub>max</sub>		V <sub>cmax</sub>		J <sub>max</sub>		Leaf Total N	
		F	p	F	p	F	p	F	p	F	p	F	p
Pd	CO <sub>2</sub>	0.632	0.540	14.449	< 0.001	0.266	0.771	0.618	0.554	0.719	0.506	1.459	0.247
	Year	5.827	< 0.001	9.673	< 0.001	0.840	0.498	0.979	0.452	1.957	0.161	7.837	< 0.001
	CO <sub>2</sub> ×Year	0.249	0.995	1.250	0.306	1.189	0.374	2.817	0.051	0.511	0.810	3.793	0.006
Fr	CO <sub>2</sub>	30.737	< 0.001	5.527	0.010	2.891	0.094	1.594	0.238	0.486	0.625	4.025	0.021
	Year	10.920	< 0.001	0.403	0.890	2.661	0.095	2.232	0.118	2.364	0.103	13.145	< 0.001
	CO <sub>2</sub> ×Year	2.537	0.021	0.456	0.935	0.545	0.765	1.368	0.292	0.507	0.815	2.646	0.021
Sa	CO <sub>2</sub>	11.368	< 0.001	13.344	< 0.001	6.983	0.014	2.890	0.094	1.782	0.210	0.178	0.838
	Year	0.179	0.987	0.583	0.762	0.918	0.433	5.243	0.015	2.931	0.0768	0.765	0.522
	CO <sub>2</sub> ×Year	2.611	0.018	1.809	0.097	1.123	0.404	0.632	0.702	0.708	0.649	1.288	0.289

**Fig. 2** - Effects of CO<sub>2</sub> concentration (aCO<sub>2</sub>, eCO<sub>2</sub>1.4, eCO<sub>2</sub>1.8) on maximum photosynthetic rate (A<sub>max</sub>), maximum rate of carboxylation (V<sub>cmax</sub>), and maximum rate of electron transport (J<sub>max</sub>) in: (a, d, g) Korean red pine (*Pinus densiflora*); (b, e, h) Chinese ash (*Fraxinus rhynchophylla*); and (c, f, i) Korean mountain ash (*Sorbus alnifolia*). The values shown are mean ± SE. Different lowercase letters above the bars indicate significant (p < 0.05) differences from multiple comparison among CO<sub>2</sub> treatments. (ns): not significant.



fer significantly ( $p = 0.194$  – Tab. 3). Similar to the A<sub>max</sub>, the V<sub>cmax</sub> value by year was  $68.0 \pm 5.2 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2013 and it significantly decreased to  $46.7 \pm 2.6 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2017 ( $p = 0.03$ ).

In terms of response by species and year, Korean red pine showed no significant differences with CO<sub>2</sub> and year ( $p = 0.554$  and  $p = 0.452$ , respectively – Tab. 4). Contrastingly, V<sub>cmax</sub> showed a decreasing tendency of about 46.4% from 2013 ( $98.1 \pm 8.1 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to 2017 ( $52.6 \pm 5.9 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Depending on the CO<sub>2</sub> concentration, it also showed same decreasing trend by about 14.8% from aCO<sub>2</sub> ( $70.9 \pm 7.1 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to eCO<sub>2</sub>1.8 ( $60.4 \pm 7.5 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) and the difference between eCO<sub>2</sub>1.8 and aCO<sub>2</sub> was the highest and significantly different in 2017 ( $p = 0.002$  – Fig. 2d). The annual V<sub>cmax</sub> of Korean mountain ash decreased significantly by 23.5% from  $55.0 \pm 4.5 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2013 to  $42.1 \pm 2.0 \mu\text{mol m}^{-2} \text{s}^{-1}$  in 2017 ( $p=0.015$ ).

For Chinese ash, no statistical differences were found. However, its annual V<sub>cmax</sub> decreased under eCO<sub>2</sub>1.8 by 55.7% compared with that under aCO<sub>2</sub> in 2015 ( $p = 0.03$ , Fig. 2e), but there was no difference thereafter. All species showed no significant differences in V<sub>cmax</sub> due to CO<sub>2</sub> enhancement (minimum  $p = 0.094$ ).

The J<sub>max</sub> showed no significant differences under eCO<sub>2</sub> and year (Tab. 3, Tab. 4). However, it had a decreasing tendency of about 23.4% from 2013 ( $100.2 \pm 7.2 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to 2017 ( $76.8 \pm 3.1 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) with no significance. Depending on the CO<sub>2</sub> concentration, there was no significant differences and tendency (minimum  $p = 0.210$ ). In terms of response by species, all species showed no statistical differences with CO<sub>2</sub> and year (Fig. 2g, Fig. 2h, Fig. 2i).


**Leaf and soil N**

The leaf total nitrogen was significantly

different under eCO<sub>2</sub> and year (maximum  $p = 0.026$  – Tab. 3). The leaf total N exposed to CO<sub>2</sub> significantly ( $p = 0.026$ ) decreased in the following order: aCO<sub>2</sub> ( $1.57 \pm 0.05 \%$ ) > eCO<sub>2</sub>1.8 ( $1.28 \pm 0.04 \%$ ) > eCO<sub>2</sub>1.4 ( $1.24 \pm 0.03 \%$ ).


Korean red pine showed significant differences in leaf total N under CO<sub>2</sub> exposure ( $p < 0.001$  – Tab. 4). The leaf total N decreased significantly by about 24.4% from 2013 ( $1.31 \pm 0.08 \%$ ) to 2017 ( $0.99 \pm 0.03 \%$ ). The leaf total N under eCO<sub>2</sub> decreased in the following order: aCO<sub>2</sub> ( $1.45 \pm 0.15 \%$ ) > eCO<sub>2</sub>1.8 ( $1.21 \pm 0.09 \%$ ) > eCO<sub>2</sub>1.4 ( $1.07 \pm 0.04 \%$ ) but it was not significant ( $p = 0.247$ ). The eCO<sub>2</sub>1.4 treatment showed the lowest leaf total N and the difference between aCO<sub>2</sub> and eCO<sub>2</sub>1.4 were 31.0% in 2013, 27.9% in 2015, and 10% in 2017 (Fig. 3a).

Chinese ash showed a significant decrease in leaf total N under eCO<sub>2</sub> and year ( $p = 0.021$  and  $p < 0.001$ , respectively). The



**Fig. 3** - Effects of CO<sub>2</sub> concentration (aCO<sub>2</sub>, eCO<sub>2</sub>1.4, eCO<sub>2</sub>1.8) on leaf total nitrogen content (%) in (a) Korean red pine (*Pinus densiflora*), (b) Chinese ash (*Fraxinus rhynchophylla*), (c) Korean mountain ash (*Sorbus alnifolia*). The values shown are mean ± SE. Different lowercase letters above the bars indicate significant (p < 0.05) differences from multiple comparison among CO<sub>2</sub> treatments. (ns): not significant.

**Fig. 4** - Effects of elevated CO<sub>2</sub> concentration (aCO<sub>2</sub>, eCO<sub>2</sub>1.4, eCO<sub>2</sub>1.8) on the soil total nitrogen (%) in 2017. Values are mean ± SE. There were no significant differences among CO<sub>2</sub> treatments ( $p > 0.05$ ).



leaf total N under eCO<sub>2</sub> decreased significantly ( $p = 0.021$ ) in the following order: aCO<sub>2</sub> ( $1.78 \pm 0.04\%$ ) > eCO<sub>2</sub>1.4 ( $1.41 \pm 0.04\%$ ) > eCO<sub>2</sub>1.8 ( $1.36 \pm 0.05\%$ ). In particular, the leaf total N was higher under eCO<sub>2</sub>1.4 and eCO<sub>2</sub>1.8 than under aCO<sub>2</sub> in 2013 ( $p < 0.001$ ), but it became significantly lower under eCO<sub>2</sub>1.4 and eCO<sub>2</sub>1.8 than under aCO<sub>2</sub> over time (Fig. 3b). The difference between aCO<sub>2</sub> and eCO<sub>2</sub>1.8 was 27.8% in 2016, and 18.8% in 2017 (maximum  $p = 0.04$ ).

Korean mountain ash showed no significant differences under eCO<sub>2</sub> and year (minimum  $p = 0.522$ ). But it decreased significantly by 15.4% under eCO<sub>2</sub>1.4 compared with aCO<sub>2</sub> only in 2017 ( $p = 0.03$  – Fig. 3c).

There was no significant difference in soil total N under eCO<sub>2</sub> in 2017 ( $p = 0.125$ ). The absolute concentration of soil total N was very low, and therefore, the variation depending on CO<sub>2</sub> concentration was negligible (Fig. 4).

## Discussion

### Relationship between the CO<sub>2</sub> fertilization effect and growth

Several studies have reported that elevated CO<sub>2</sub> concentration due to climate change directly affects photosynthesis, suggesting that eCO<sub>2</sub> promotes ecosystem production and biomass accumulation (McCarthy et al. 2010, Norby et al. 2010, Reich et al. 2018). For instances, the net primary production (NPP) increased by 22-32% in the Duke-FACE experiment, which conducted the CO<sub>2</sub> exposure for more than 10 years. In the Pop-FACE experiment, the diameter increased about 5% and the 12 species leaf area increased about 8% in seven FACE (Ainsworth & Long 2005, Finzi et al. 2007, McCarthy et al. 2010). In this study, the annual growth of the three species was analysed, confirming a significant enhancement in height and diameter under eCO<sub>2</sub> for the first two years. However, the CO<sub>2</sub> fertilization effect was not sustained thereafter (Tab. 1, Tab. 2, Fig. 1). Temporary CO<sub>2</sub> fertilization effect was also reported by other studies. In the Oak-Ridge FACE and BioCON experiments, the reduction of CO<sub>2</sub> effect was offset after 6 and 3

years, respectively, and nitrogen fertilization showed an immediate increase in NPP (Reich et al. 2006, Norby et al. 2010). In addition, there was no CO<sub>2</sub> effect in *Picea abies* growing on a very low-nutrient soil (Ward et al. 2008). Moreover, in the Duke FACE experiment, where the CO<sub>2</sub> fertilization effect remained for more than 10 years, soil N positively correlated with productivity increment under eCO<sub>2</sub> (McCarthy et al. 2010). Thus, soil nutrients and CO<sub>2</sub> fertilization effect appeared to be closely related. In particular, in our study site, the soil total N was very low because the nearby forest soil was mixed with sand and leaf total N decreased over time. For this reason, we argue that growth promotion through the CO<sub>2</sub> fertilization effect was not sustained.

### Changes in photosynthetic properties under elevated CO<sub>2</sub> concentration

Elevated atmospheric CO<sub>2</sub> concentration increases the difference in the partial pressure of CO<sub>2</sub> between the atmosphere and leaf mesophyll tissues, increasing the maximum photosynthetic rate (Curtis & Wang 1998, Ainsworth & Rogers 2007). Twelve FACE experiments showed that the average A<sub>max</sub> of all the tested species increased by about 31%. Depending on species, it was increased by about 37% in *Pinus australis* (Mozdzer & Caplan 2018), 67% in *P. taeda*, and 62% in *Liquidambar styraciflua* (Ellsworth et al. 2012). However, over time, the increase of A<sub>max</sub> under eCO<sub>2</sub> was reduced by the acclimation of photosynthetic properties (Ainsworth & Rogers 2007). In our study, the reduction of A<sub>max</sub> increase under eCO<sub>2</sub> varied depending on the species (Fig. 2). Korean red pine and Korean mountain ash did not show a significant difference since 2013, which was only three years after eCO<sub>2</sub> exposure. Chinese ash showed the enhancement in 2013, but the increase in A<sub>max</sub> was reduced over time. A<sub>max</sub> of Korean mountain ash under aCO<sub>2</sub> was maintained over time, but it was decreased in Chinese ash, which might have been due to the low soil nutrient (Fig. 2, Fig. 4). This decrease in the A<sub>max</sub> resulted in the down-regulation of the V<sub>Cmax</sub> and J<sub>max</sub>. In other words, the A<sub>max</sub>

is usually measured under the CO<sub>2</sub> concentration of each treatment, therefore no difference or a decrease indicates that photosynthesis enhancement is not maintained, rather lowered over time. Similar to previous studies (Drake et al. 1997, Martínez-Carrasco et al. 2005), we observed a reduction in the V<sub>Cmax</sub> and J<sub>max</sub> under eCO<sub>2</sub>. This reduction usually occurs after long term CO<sub>2</sub> exposure (Ainsworth & Rogers 2007, Norby et al. 2010). In *P. ponderosa* exposed to eCO<sub>2</sub> for 6 years, the V<sub>Cmax</sub> and J<sub>max</sub> were decreased by about 36% and 21%, respectively (Tissue et al. 1999), and the V<sub>Cmax</sub> was decreased by about 19% in *P. abies* (Uddling & Wallin 2012). However, in some cases, the reduction of photosynthetic ability under eCO<sub>2</sub> did not occur even in the long term (more than 6 years – Bader et al. 2010, Darbah et al. 2010, Warren et al. 2015). In an experiment on *Fagus sylvatica*, *Quercus petraea*, *Carpinus betulus*, *Acer campestre*, and *Tilia platyphyllos* in mature deciduous forests, the increase in the A<sub>max</sub> was maintained even after 8 years of CO<sub>2</sub> treatment, and the down-regulation of V<sub>Cmax</sub> and J<sub>max</sub> did not occur. Similarly, in the Aspen FACE experiment, the increases in V<sub>Cmax</sub> (48%-85%) and J<sub>max</sub> (23%-34%) were sustained for 11 years, and no reduction occurred. Warren et al. (2015) also showed that the increase in photosynthesis was maintained during the first 8 years when the leaf N was 0.2 mg cm<sup>-2</sup> or more, but after a decrease of leaf N, the CO<sub>2</sub> fertilization effect disappeared. Thus, the increase in photosynthesis and the decrease in photosynthetic ability are largely affected by the nitrogen available in the environment. These studies were conducted in well-developed organic layer of forest floor, such as mature stands (Bader et al. 2010), or soil N of 3% or more (Darbah et al. 2010), i.e., environments with considerably higher N than normal forest soil. On the contrary, the long-term exposure under eCO<sub>2</sub> leads to nitrogen limitation in common forest soil, which results in PNL that decreases photosynthesis and growth increment (Reich et al. 2006, Norby et al. 2010, Feng et al. 2015, Rütting 2017).

### Relationship between photosynthetic properties (V<sub>Cmax</sub> and J<sub>max</sub>) and leaf nitrogen

In general, the leaf total N is positively correlated with the photosynthesis ability (Evans & Seemann 1989). Several studies have shown that the leaf N decreased in response to long-term eCO<sub>2</sub> condition (Ellsworth et al. 2012). The elevated CO<sub>2</sub> concentration increases the capacity of the rubisco enzyme to adsorb carbon dioxide, thereby causing carboxylation and a decrease in the demand of nitrogen for rubisco (Nowak et al. 2004, Liu et al. 2012). Similarly, in this study, the leaf N was significantly lowered under eCO<sub>2</sub> over time (Fig. 3). In addition, studies have shown that the V<sub>Cmax</sub> and photosynthesis ability, decreased as the leaf N decreased (Ainsworth & Long



2005, Warren et al. 2015). Thus, fertilization at the time of nitrogen limitation increased the CO<sub>2</sub> fertilization effect again or the decreasing rate was reduced (Liberloo et al. 2007, Crous et al. 2008, Terrer et al. 2018).

In general, there is a strong correlation between  $V_{cmax}$  and  $J_{max}$ , with  $J_{max}/V_{cmax} = 1.5$ – $2.0$  at 25 °C (Medlyn et al. 2002, Walker et al. 2014, Cho et al. 2019). Similarly,  $V_{cmax}$ ,  $J_{max}$  and leaf N showed correlation under eCO<sub>2</sub> (Crous et al. 2008, Warren et al. 2015). Therefore, when the leaf N and  $V_{cmax}$  were decreased after long-term eCO<sub>2</sub> treatment, the  $J_{max}$  also decreased in some cases (Herrick & Thomas 2001, Medlyn et al. 2002, Sholtis et al. 2004, Crous et al. 2008, Bader et al. 2010). As in our study, the decrease in  $J_{max}$  (Tab. 4) was relatively lower than that of  $V_{cmax}$  in some cases (Liberloo et al. 2007, Warren et al. 2015). Moreover, these changes are different depending on species. Chinese ash and Korean red pine showed a decrease only in the  $V_{cmax}$ , while Korean mountain ash did not show any decrease in both the above photosynthetic parameters (Fig. 2). These differences due to the change in intracellular nitrogen distribution under eCO<sub>2</sub> are dependent not only on the environment but also on the species. Further studies are needed to examine how species change the intercellular nitrogen distribution under elevated CO<sub>2</sub> (Evans & Seemann 1989, Hikosaka 2004).

#### CO<sub>2</sub> fertilization effect and PNL

The enhanced productivity of forest ecosystem due to the CO<sub>2</sub> fertilization is heavily influenced by the nutrients available in the soil (Finzi et al. 2007). PNL is a hypothesis that the increased forest productivity due to eCO<sub>2</sub> decreases over time because of the increased N accumulation in the biomass, resulting in a decrease of soil N availability, an increase of N immobilization and a decrease of N mineralization (Luo et al. 2004). In the long-term eCO<sub>2</sub> at DukeFACE, Oak-Ridge FACE, and BioCON experiments, N fertilization led to an immediate increase in NPP. Therefore, there seemed to be an interaction between eCO<sub>2</sub> and N in terms of increased productivity (Reich et al. 2006, McCarthy et al. 2010, Norby et al. 2010). In particular, the soil N is important and N deficiency leads to a decline in production due to PNL even under eCO<sub>2</sub> (Norby et al. 2016). Such N deficiency changes the N distribution in the plants' organs for effective nitrogen utilization, resulting in a decrease of the N used for above-ground photosynthesis (Pettersson & McDonald 1994). In addition, N distribution in photosynthetic apparatus such as Rubisco is greatly reduced. Therefore, nitrogen-use efficiency is increased generally (Vicente et al. 2016, Sharwood et al. 2017), but the persistence of such an increase is controversial (Ainsworth & Rogers 2007, Leakey et al. 2009). There is limited information about N distribution in the above- and below-ground under eCO<sub>2</sub>. Therefore, it is necessary to determine and quantify the

distribution of N in the above- and below-ground through follow-up studies.

#### Conclusions

In this study, we investigated the physiological and morphological characteristics of Korean red pine, Chinese ash, and Korean mountain ash, which are native tree species in Korea, under eCO<sub>2</sub> over a period of 8 years. We also examined the longevity of CO<sub>2</sub> fertilization effect in our study sites.

The CO<sub>2</sub> fertilization effect caused by eCO<sub>2</sub> led to an increase in growth in the early stage of exposure, but there was no significant difference thereafter. Photosynthetic properties showed a decreasing tendency in all species and a down-regulation of photosynthetic capacity increase with time, especially in Korean red pine and Korean mountain ash. The analysis of leaf and soil N to identify the cause revealed a significant decrease of leaf N under eCO<sub>2</sub>. We argue that the enhancement of productivity might have decreased due to low soil N.

In conclusion, progressive nitrogen limitation (PNL) caused by the N reduction might have already started or is about to start in our study sites. Further investigation is needed to clarify N use efficiency and nitrogen distribution according to species.

#### List of abbreviations

(OTC): Open Top Chamber; (aCO<sub>2</sub>): current atmospheric CO<sub>2</sub> concentration; (eCO<sub>2</sub>,1.4): 1.4 times higher CO<sub>2</sub> concentration than the current concentration (Chamber 2); (eCO<sub>2</sub>,1.8): 1.8 times higher CO<sub>2</sub> concentration than the current concentration (Chamber 3); ( $A_{max}$ ): maximum photosynthetic rate; ( $V_{cmax}$ ): maximum rate of carboxylation; ( $J_{max}$ ): maximum electron transport rate.

#### Acknowledgments

This research was funded by the National Research Foundation of Korea, by the projects “Evaluation of climate change adaptation mechanisms of native temperate tree species and its ecological effects using open-top chambers” (2014R1A1A2055127) and “Testing progressive nitrogen limitation and nitrogen use efficiency increment through the quantification of leaf nitrogen allocation” (2017R1A2B2012605). We thank members of forest ecophysiology laboratory for their assistance with field sampling.

#### References

Ainsworth EA, Long SP (2005). What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytologist* 165: 351–372. - doi: [10.1111/j.1469-8137.2004.01224.x](https://doi.org/10.1111/j.1469-8137.2004.01224.x)

Ainsworth EA, Rogers A (2007). The response of photosynthesis and stomatal conductance to rising [CO<sub>2</sub>]: mechanisms and environmental interactions. *Plant, Cell and Environment* 30: 258–270. - doi: [10.1111/j.1365-3040.2007.01641.x](https://doi.org/10.1111/j.1365-3040.2007.01641.x)

Bader MKF, Siegwolf R, Körner C (2010). Sustained enhancement of photosynthesis in mature deciduous forest trees after 8 years of free air CO<sub>2</sub> enrichment. *Planta* 232: 1115–1125. - doi: [10.1007/s00425-010-1240-8](https://doi.org/10.1007/s00425-010-1240-8)

Cai C, Li G, Yang H, Yang J, Liu H, Struik PC, Luo W, Yin X, Di L, Guo X (2018). Do all leaf photosynthesis parameters of rice acclimate to elevated CO<sub>2</sub>, elevated temperature, and their combination in FACE environments? *Global Change Biology* 24: 1685–1707. - doi: [10.1111/gcb.13961](https://doi.org/10.1111/gcb.13961)

Cho S, Ser-Oddamba B, Batkhuu N, Kim HS (2019). Comparison of water use efficiency and biomass production in 10-year-old *Populus sibirica* and *Ulmus pumila* plantations in Lun soum, Mongolia. *Forest Science and Technology* 15: 147–158. - doi: [10.1080/21580103.2019.1634646](https://doi.org/10.1080/21580103.2019.1634646)

Crous KY, Walters MB, Ellsworth DS (2008). Elevated CO<sub>2</sub> concentration affects leaf photosynthesis-nitrogen relationships in *Pinus taeda* over nine years in FACE. *Tree Physiology* 28: 607–614. - doi: [10.1093/treephys/28.4.607](https://doi.org/10.1093/treephys/28.4.607)

Curtis PS, Wang X (1998). A meta-analysis of elevated CO<sub>2</sub> effects on woody plant mass, form, and physiology. *Oecologia* 113: 299–313. - doi: [10.1007/s004420050381](https://doi.org/10.1007/s004420050381)

Darbah JN, Kubiske ME, Nelson N, Kets K, Riikonen J, Sober A, Rouse L, Karnosky DF (2010). Will photosynthetic capacity of aspen trees acclimate after long-term exposure to elevated CO<sub>2</sub> and O<sub>3</sub>? *Environmental Pollution* 158: 983–991. - doi: [10.1016/j.envpol.2009.10.022](https://doi.org/10.1016/j.envpol.2009.10.022)

Drake BG, González-Meler MA, Long SP (1997). More efficient plants: a consequence of rising atmospheric CO<sub>2</sub>? *Annual Review of Plant Biology* 48: 609–639. - doi: [10.1146/annurev.arplant.48.1.609](https://doi.org/10.1146/annurev.arplant.48.1.609)

Ellsworth DS, Thomas R, Crous KY, Palmroth S, Ward E, Maier C, DeLucia E, Oren R (2012). Elevated CO<sub>2</sub> affects photosynthetic responses in canopy pine and subcanopy deciduous trees over 10 years: a synthesis from Duke FACE. *Global Change Biology* 18: 223–242. - doi: [10.1111/j.1365-2486.2011.02505.x](https://doi.org/10.1111/j.1365-2486.2011.02505.x)

Evans JR, Seemann JR (1989). The allocation of protein nitrogen in the photosynthetic apparatus: costs, consequences, and control. *Photosynthesis* 183: 205. [online] URL: <http://www.researchgate.net/publication/285010849>

Feng Z, Rütting T, Pleijel H, Wallin G, Reich PB, Kammann CI, Newton PC, Kobayashi K, Luo Y, Uddling J (2015). Constraints to nitrogen acquisition of terrestrial plants under elevated CO<sub>2</sub>. *Global Change Biology* 21: 3152–3168. - doi: [10.1111/gcb.12938](https://doi.org/10.1111/gcb.12938)

Finzi AC, Norby RJ, Calfapietra C, Gallet-Budynek A, Gielen B, Holmes WE, Hoosbeek MR, Iversen CM, Jackson RB, Kubiske ME (2007). Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated CO<sub>2</sub>. *Proceedings of the National Academy of Sciences USA* 104: 14014–14019. - doi: [10.1073/pnas.0706518104](https://doi.org/10.1073/pnas.0706518104)

Harley PC, Loreto F, Di Marco G, Sharkey TD (1992). Theoretical considerations when estimating the mesophyll conductance to CO<sub>2</sub> flux by analysis of the response of photosynthesis to CO<sub>2</sub>. *Plant Physiology* 98: 1429–1436. - doi: [10.1104/pp.98.4.1429](https://doi.org/10.1104/pp.98.4.1429)

Harrison MT, Edwards EJ, Farquhar GD, Nicotra

- AB, Evans JR (2009). Nitrogen in cell walls of sclerophyllous leaves accounts for little of the variation in photosynthetic nitrogen-use efficiency. *Plant, Cell and Environment* 32: 259-270. - doi: [10.1111/j.1365-3040.2008.01918.x](https://doi.org/10.1111/j.1365-3040.2008.01918.x)
- Herrick J, Thomas R (2001). No photosynthetic down-regulation in sweetgum trees (*Liquidambar styraciflua* L.) after three years of CO<sub>2</sub> enrichment at the Duke Forest FACE experiment. *Plant, Cell and Environment* 24: 53-64. - doi: [10.1046/j.1365-3040.2001.00652.x](https://doi.org/10.1046/j.1365-3040.2001.00652.x)
- Hikosaka K (2004). Interspecific difference in the photosynthesis-nitrogen relationship: patterns, physiological causes, and ecological importance. *Journal of Plant Research* 117: 481-494. - doi: [10.1007/s10265-004-0174-2](https://doi.org/10.1007/s10265-004-0174-2)
- Hungate BA, Dijkstra P, Wu Z, Duval BD, Day FP, Johnson DW, Megonigal JP, Brown AL, Garland JL (2013). Cumulative response of ecosystem carbon and nitrogen stocks to chronic CO<sub>2</sub> exposure in a subtropical oak woodland. *New Phytologist* 200: 753-766. - doi: [10.1111/nph.12333](https://doi.org/10.1111/nph.12333)
- Hungate BA, Dukes JS, Shaw MR, Luo Y, Field CB (2003). Nitrogen and climate change. *Science* 302: 1512-1513. - doi: [10.1126/science.1091390](https://doi.org/10.1126/science.1091390)
- IPCC (2013). Summary for policymakers. climate change: the physical science basis. Contribution of Working Group to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK, pp. 1-33.
- Kwon B, Kim HS, Yi MJ (2019). The quantity and pattern of leaf fall and nitrogen resorption strategy by leaf-litter in the Gwangneung natural broadleaved forest. *Korean Journal of Agricultural and Forest Meteorology* 21: 208-220. [online] URL: <http://www.koreascience.or.kr/article/JAKO201929064695163.page>
- Leakey AD, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR (2009). Elevated CO<sub>2</sub> effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* 60: 2859-2876. - doi: [10.1093/jxb/erp096](https://doi.org/10.1093/jxb/erp096)
- Lee JC, Kim DH, Kim GN, Kim PG, Han S-H (2012). Long-term climate change research facility for trees: CO<sub>2</sub>-enriched open top chamber system. *Journal of Agricultural and Forest Meteorology* 14: 19-27. - doi: [10.5532/KJAFM.2012.14.1.019](https://doi.org/10.5532/KJAFM.2012.14.1.019)
- Liberloo M, Tulva I, Raim O, Kull O, Ceulemans R (2007). Photosynthetic stimulation under long-term CO<sub>2</sub> enrichment and fertilization is sustained across a closed *Populus* canopy profile (EUROFACE). *New Phytologist* 173: 537-549. - doi: [10.1111/j.1469-8137.2006.01926.x](https://doi.org/10.1111/j.1469-8137.2006.01926.x)
- Liu J, Zhang D, Zhou G, Duan H (2012). Changes in leaf nutrient traits and photosynthesis of four tree species: effects of elevated [CO<sub>2</sub>], N fertilization and canopy positions. *Journal of Plant Ecology* 5: 376-390. - doi: [10.1093/jpe/rtso06](https://doi.org/10.1093/jpe/rtso06)
- Luo Y, Su B, Currie WS, Dukes JS, Finzi A, Hartwig U, Hungate B, McMurtrie RE, Oren R, Parton W (2004). Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *Bioscience* 54: 731-739. - doi: [10.1641/0006-3568\(2004\)054\[0731:PNLOER\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0731:PNLOER]2.0.CO;2)
- Makino A, Osmond B (1991). Effects of nitrogen nutrition on nitrogen partitioning between chloroplasts and mitochondria in pea and wheat. *Plant Physiology* 96: 355-362. - doi: [10.1104/pp.96.2.355](https://doi.org/10.1104/pp.96.2.355)
- Martínez-Carrasco R, Pérez P, Morcuende R (2005). Interactive effects of elevated CO<sub>2</sub>, temperature and nitrogen on photosynthesis of wheat grown under temperature gradient tunnels. *Environmental Experimental Botany* 54: 49-59. - doi: [10.1016/j.envexpbot.2004.05.004](https://doi.org/10.1016/j.envexpbot.2004.05.004)
- McCarthy HR, Oren R, Johnsen KH, Gallet-Budynek A, Pritchard SG, Cook CW, LaDeau SL, Jackson RB, Finzi AC (2010). Re-assessment of plant carbon dynamics at the Duke free-air CO<sub>2</sub> enrichment site: interactions of atmospheric [CO<sub>2</sub>] with nitrogen and water availability over stand development. *New Phytologist* 185: 514-528. - doi: [10.1111/j.1469-8137.2009.03078.x](https://doi.org/10.1111/j.1469-8137.2009.03078.x)
- Medlyn B, Dreyer E, Ellsworth D, Forstreuter M, Harley P, Kirschbaum M, Le Roux X, Montpied P, Strassmeyer J, Walcroft A (2002). Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data. *Plant, Cell and Environment* 25: 1167-1179. - doi: [10.1046/j.1365-3040.2002.00891.x](https://doi.org/10.1046/j.1365-3040.2002.00891.x)
- Mozdzer TJ, Caplan JS (2018). Complementary responses of morphology and physiology enhance the stand-scale production of a model invasive species under elevated CO<sub>2</sub> and nitrogen. *Functional Ecology* 32: 1784-1796. - doi: [10.1111/1365-2435.13106](https://doi.org/10.1111/1365-2435.13106)
- Norby RJ, Warren JM, Iversen CM, Medlyn BE, McMurtrie RE (2010). CO<sub>2</sub> enhancement of forest productivity constrained by limited nitrogen availability. *Proceedings of the National Academy of Sciences USA* 107: 19368-19373. - doi: [10.1073/pnas.1006463107](https://doi.org/10.1073/pnas.1006463107)
- Norby RJ, De Kauwe MG, Domingues TF, Durusma RA, Ellsworth DS, Goll DS, Lapola DM, Luus KA, MacKenzie AR, Medlyn BE (2016). Model-data synthesis for the next generation of forest free-air CO<sub>2</sub> enrichment (FACE) experiments. *New Phytologist* 209: 17-28. - doi: [10.1111/nph.13593](https://doi.org/10.1111/nph.13593)
- Nowak RS, Ellsworth DS, Smith SD (2004). Functional responses of plants to elevated atmospheric CO<sub>2</sub> - do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist* 162: 253-280. - doi: [10.1111/j.1469-8137.2004.01033.x](https://doi.org/10.1111/j.1469-8137.2004.01033.x)
- Pettersson R, McDonald AJS (1994). Effects of nitrogen supply on the acclimation of photosynthesis to elevated CO<sub>2</sub>. *Photosynthesis Research* 39: 389-400. - doi: [10.1007/BF00014593](https://doi.org/10.1007/BF00014593)
- R Core Team (2016). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [online] URL: <http://www.r-project.org/>
- Reich PB, Hungate BA, Luo Y (2006). Carbon-nitrogen interactions in terrestrial ecosystems in response to rising atmospheric carbon dioxide. *Annual Review of Ecology, Evolution, and Systematics* 37: 611-636. - doi: [10.1146/annurev.ecolsys.37.091305.110039](https://doi.org/10.1146/annurev.ecolsys.37.091305.110039)
- Reich PB, Hobbie SE, Lee TD, Pastore MA (2018). Unexpected reversal of C<sub>3</sub> versus C<sub>4</sub> grass response to elevated CO<sub>2</sub> during a 20-year field experiment. *Science* 360: 317-320. - doi: [10.1126/science.aas9313](https://doi.org/10.1126/science.aas9313)
- Rütting T (2017). Nitrogen mineralization, not N<sub>2</sub> fixation, alleviates progressive nitrogen limitation - Comment on "Processes regulating progressive nitrogen limitation under elevated carbon dioxide: a meta-analysis" by Liang et al. (2016). *Biogeosciences* 14: 751-754. - doi: [10.5194/bg-14-751-2017](https://doi.org/10.5194/bg-14-751-2017)
- Sang WG, Kim JH, Shin P, Baek JK, Lee YH, Cho JI, Seo MC (2019). Evaluation of Photochemical Reflectance Index (PRI) response to soybean drought stress under climate change conditions. *Korean Journal of Agricultural and Forest Meteorology* 21: 261-268. [online] URL: <http://www.koreascience.or.kr/article/JAKO201904858009849.page>
- Sharkey TD (2016). What gas exchange data can tell us about photosynthesis. *Plant, Cell and Environment* 39: 1161-1163. - doi: [10.1111/pce.12641](https://doi.org/10.1111/pce.12641)
- Sharwood RE, Crous KY, Whitney SM, Ellsworth DS, Ghannoum O (2017). Linking photosynthesis and leaf N allocation under future elevated CO<sub>2</sub> and climate warming in *Eucalyptus globulus*. *Journal of Experimental Botany* 68: 1157-1167. - doi: [10.1093/jxb/erw484](https://doi.org/10.1093/jxb/erw484)
- Sholtis JD, Gunderson CA, Norby RJ, Tissue DT (2004). Persistent stimulation of photosynthesis by elevated CO<sub>2</sub> in a sweetgum (*Liquidambar styraciflua*) forest stand. *New Phytologist* 162: 343-354. - doi: [10.1111/j.1469-8137.2004.01028.x](https://doi.org/10.1111/j.1469-8137.2004.01028.x)
- Taiz L, Zeiger E (2010). Photosynthesis: the light reactions. In: "Plant Physiology" (5<sup>th</sup> edn). Sinauer Associates Inc, Sunderland, USA, pp. 163-198. [online] URL: <http://www.sinauer.com/media/wysiwyg/tocs/PlantPhysiology5.pdf>
- Talhelm AF, Pregitzer KS, Kubiske ME, Zak DR, Company CE, Burton AJ, Dickson RE, Hendrey GR, Isebrands JG, Lewin KF (2014). Elevated carbon dioxide and ozone alter productivity and ecosystem carbon content in northern temperate forests. *Global Change Biology* 20: 2492-2504. - doi: [10.1111/gcb.12564](https://doi.org/10.1111/gcb.12564)
- Terrer C, Vicca S, Stocker BD, Hungate BA, Phillips RP, Reich PB, Finzi AC, Prentice IC (2018). Ecosystem responses to elevated CO<sub>2</sub> governed by plant-soil interactions and the cost of nitrogen acquisition. *New Phytologist* 217: 507-522. - doi: [10.1111/nph.14872](https://doi.org/10.1111/nph.14872)
- Tissue DT, Griffin KL, Ball JT (1999). Photosynthetic adjustment in field-grown ponderosa pine trees after six years of exposure to elevated CO<sub>2</sub>. *Tree Physiology* 19: 221-228. - doi: [10.1093/treephys/19.4-5.221](https://doi.org/10.1093/treephys/19.4-5.221)
- Uddling J, Wallin G (2012). Interacting effects of elevated CO<sub>2</sub> and weather variability on photosynthesis of mature boreal Norway spruce agree with biochemical model predictions. *Tree Physiology* 32: 1509-1521. - doi: [10.1093/treephys/tps086](https://doi.org/10.1093/treephys/tps086)
- Vicente R, Pérez P, Martínez-Carrasco R, Feil R, Lunn JE, Watanabe M, Arrivault S, Stitt M, Höfgen R, Morcuende R (2016). Metabolic and transcriptional analysis of durum wheat responses to elevated CO<sub>2</sub> at low and high nitrate supply. *Plant and Cell Physiology* 57: 2133-2146. - doi: [10.1093/pcp/pcw131](https://doi.org/10.1093/pcp/pcw131)
- Walker AP, Beckerman AP, Gu L, Kattge J, Cernusak LA, Domingues TF, Scales JC, Wohlfahrt G, Wullschlegel SD, Woodward FI (2014). The relationship of leaf photosynthetic traits - V<sub>cm</sub> and J<sub>max</sub> - to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis and modeling



study. *Ecology and Evolution* 4: 3218-3235. - doi: [10.1002/ece3.1173](https://doi.org/10.1002/ece3.1173)  
Wang YP, Houlton BZ (2009). Nitrogen constraints on terrestrial carbon uptake: Implications for the global carbon-climate feedback. *Geophysical Research Letters* 36 (24): 623. - doi: [10.1029/2009GL041009](https://doi.org/10.1029/2009GL041009)  
Ward EJ, Oren R, Sigurdsson BD, Jarvis PG, Linder S (2008). Fertilization effects on mean sto-

matal conductance are mediated through changes in the hydraulic attributes of mature Norway spruce trees. *Tree Physiology* 28: 579-596. - doi: [10.1093/treephys/28.4.579](https://doi.org/10.1093/treephys/28.4.579)  
Warren JM, Jensen AM, Medlyn BE, Norby RJ, Tissue DT (2015). Carbon dioxide stimulation of photosynthesis in *Liquidambar styraciflua* is not sustained during a 12-year field experiment. *AoB Plants* 7: 351. - doi: [10.1093/aobpla/plu074](https://doi.org/10.1093/aobpla/plu074)

## Supplementary Material

**Fig. S1** – The Open Top Chamber (OTC) user in the experiment.

**Link:** [Song\\_3288@suppl001.pdf](mailto:Song_3288@suppl001.pdf)